

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 February 2002 (14.02.2002)

PCT

(10) International Publication Number
WO 02/12340 A2

Published:
— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guide
once Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 02/12340 A2



(51) International Patent Classification: C07K 14/705

(21) International Application Number: PCT/US01/24217

(22) International Filing Date: 1 August 2001 (01.08.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Date:
6/02/23, 3/69 3 August 2000 (03.08.2000) US
6/02/24, 4/56 10 August 2000 (10.08.2000) US
6/02/26, 4/10 18 August 2000 (18.08.2000) US
6/02/28, 1/40 22 August 2000 (25.08.2000) US
6/02/30, 0/57 31 August 2000 (31.08.2000) US
6/02/31, 4/34 8 September 2000 (08.09.2000) US

(71) Applicant (for all designated States except US): INCTTE
GENOMICS, INC. (US/US): 3160 Porter Drive, Palo
Alto, CA 94304 (US).

(72) Inventors: and
(75) Inventor/Applicants (for US only): YUE, Henry
(US/US): 826 Luis Avenue, Sunnyvale, CA 94087 (US);
THORNTON, Michael (US/US): 9 Melbury Road,
Woodside, CA 94062 (US); RANKUNAKI, Jayahani
(US/US): 34359 Mayfield Circle, Fremont, CA 94535
(US); TANG, Y., Tom (US/US): 4230 Kanwick Court,
San Jose, CA 95118 (US); AZIMZAI, Yalda (US/US):
5518 Boulder Canyon Drive, Castro Valley, CA 94522
(US); BAUGLIN, Mariah, R. (US/US): 14244 Santiago
Road, San Leandro, CA 94577 (US); YANG, Junliang
(US/US): 7125 Buck Lane, San Jose, CA 95129 (US);
YAO, Miquel, G. (US/US): 111 Frederick Court, Mon-
tain View, CA 94043 (US); LAI, Preeti (US/US): P.O. Box
5142, Santa Clara, CA 95056 (US); WALIA, Nandini, K.
(US/US): 800 Robie Avenue #1, San Leandro, CA 94577
(US); GANDHI, Ameeta, R. (US/US): 837 Robie Av-
enue, #1, Menlo Park, CA 94025 (US); HARVALIA, April,
J., A. (US/US): 2227 Calle de Primavera, Santa Clara,
CA 95054 (US); NGUYEN, Danielle, B. (US/US): 1403
Ridgewood Drive, San Jose, CA 95118 (US); PATTER-
SON, Chandrea (US/US): 490 Sherwood Way #1, Menlo
Park, CA 94025 (US); ELLIOTT, Vicki, S. (US/US):
3770 Hollen Place Way, San Jose, CA 95121 (US); TRI-
BOULEY, Catherine, M. (FR/US): 1121 Tennessee,

#1, San Francisco, CA 94107 (US); LU, DYUNG, Alina,
M. (US/US): 233 Coy Drive, San Jose, CA 95123 (US);
XU, Yaming (US/US): 1739 Walnut Drive, Mountain
View, CA 94040 (US); REDDY, Roopa (US/US): 1233 W.
McKinley Avenue #3, Sunnyvale, CA 94086 (US); HER-
NANDEZ, Roberto (US/US): 160 Pine Tree Avenue,
Camelburg Kent, CT 07822 (US); BOROWSKY, Mark,
L. (US/US): 122 Orchard Avenue, Redwood City, CA
94061 (US); LO, Terence, P. (US/US): 1451 Beach Park
Boulevard, Apt. 115, Foster City, CA 94024 (US); LU,
Yao (US/US): 3885 Cortina Way, Palo Alto, CA 94303
(US); POLICKY, Jennifer, L. (US/US): 1511 Jarvis
Court, San Jose, CA 95118 (US); GREENE, Barrie,
D. (US/US): 1332 10th Avenue #104, San Francisco, CA
94122 (US); SANJAYWALA, Madhu, S. (US/US): 210
Sylvia Court, Los Altos, CA 94024 (US); RAUIMANN,
Brigitte, E. (US/US): 5801 South Dorchester Avenue #3B,
Chicago, IL 60637 (US); BURFORD, Neil (GB/US): 105
Withwind Circle, Durham, CT 06422 (US); ISON, Craig,
H. (US/US): 1243 Weatherford Way, San Jose, CA 95118
(US); LEE, Ernestine, A. (US/US): 624 Kains Street,
Albany, CA 94706 (US); DING, LI (CN/US): 3351 Alamo
Street #146, Palo Alto, CA 94306 (US); DAS, Debopriya
(US/US): 1179 Bonita Avenue Apt. 3, Mountain View,
CA 94040 (US); KALLICK, Deborah, A. (US/US):
900 Olive Street, Menlo Park, CA 94025 (US); KHAN,
Farrah, A. (US/US): 3617 Central Road #102, Glenview,
IL 60025 (US); SELHAMER, Jeffrey, J. (US/US):
12555 La Cresta, Los Altos, CA 94022 (US).

(74) Agents: HANLETT-COX, Diana et al.: Invey Pharma-
ceuticals, Inc., 3160 Porter Drive, Palo Alto, CA 94304
(US).

(81) Designated States (nationally): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, FI, GB, GD, GR, GT, GU,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MY, NZ, NI, NL, NO, NZ, PT, RO, RU, SD, SE, SG, SI, SK,
TH, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW,
TG).
(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GN, GQ, GW, ML, MR, NE, NG, SN, TD,
TG).

(Continued on next page)

WO 02/12340 A2

(54) Title: TRANSPORTERS AND ION CHANNELS
(57) Abstract: The invention provides human transporters and ion channels (TRIC1) and polynucleotides which identify and en-
code TRIC1. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also
provides methods for diagnosing, treating, or preventing disorders associated with aberrant expression of TRIC1.

TRANSPORTERS AND ION CHANNELS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of transporters and ion channels and to the use of these sequences in the diagnosis, treatment, and prevention of transport, neurological, muscle, immunological, and cell proliferative disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of transporters and ion channels.

BACKGROUND OF THE INVENTION

Eukaryotic cells are surrounded and subdivided into functionally distinct organelles by hydrophobic lipid bilayer membranes which are highly impermeable to most polar molecules. Cells and organelles require transport proteins to import and export essential nutrients and metal ions including K^+ , NH_4^+ , P_i , SO_4^{2-} , sugars, and vitamins, as well as various metabolic waste products. Transport proteins also play roles in antibiotic resistance, toxin secretion, ion balance, synaptic neurotransmission, kidney function, intestinal absorption, tumor growth, and other diverse cell functions (Griffith, J. and C. Sansom (1998) *The Transporter Facts Book*, Academic Press, San Diego CA, pp. 3-29). Transport can occur by a passive concentration-dependent mechanism, or can be linked to an energy source such as ATP hydrolysis or an ion gradient. Proteins that function in transport include carrier proteins, which bind to a specific solute and undergo a conformational change that translocates the bound solute across the membrane, and channel proteins, which form hydrophilic pores that allow specific solutes to diffuse through the membrane down an electrochemical solute gradient.

Carrier proteins which transport a single solute from one side of the membrane to the other are called uniporters. In contrast, coupled transporters link the transfer of one solute with simultaneous or sequential transfer of a second solute, either in the same direction (symport) or in the opposite direction (antiport). For example, intestinal and kidney epithelium contains a variety of symporter systems driven by the sodium gradient that exists across the plasma membrane. Sodium moves into the cell down its electrochemical gradient and brings the solute into the cell with it. The sodium gradient that provides the driving force for solute uptake is maintained by the ubiquitous Na^+/K^+ ATPase system. Sodium-coupled transporters include the mammalian glucose transporter (SGLT1), iodide transporter (NIS), and multivitamin transporter (SMVT). All three transporters have twelve putative transmembrane segments, extracellular glycosylation sites, and cytoplasmically-oriented N- and C-termini. NIS plays a crucial role in the evaluation, diagnosis, and treatment of various thyroid pathologies because it is the molecular basis for radioiodide thyroid-imaging

techniques and for specific targeting of radioisotopes to the thyroid gland (Levy, O. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:5568-5573). SMVT is expressed in the intestinal mucosa, kidney, and placenta, and is implicated in the transport of the water-soluble vitamins, e.g., biotin and pantothenate (Prasad, P.D. et al. (1998) *J. Biol. Chem.* 273:7501-7506).

One of the largest families of transporters is the major facilitator superfamily (MFS), also called the uniporter-symporter-antiporter family. MFS transporters are single polypeptide carriers that transport small solutes in response to ion gradients. Members of the MFS are found in all classes of living organisms, and include transporters for sugars, oligosaccharides, phosphates, nitrates, nucleosides, monocarboxylates, and drugs. MFS transporters found in eukaryotes all have a structure comprising 12 transmembrane segments (Pao, S.S. et al. (1998) *Microbiol. Molec. Biol. Rev.* 62:1-34). The largest family of MFS transporters is the sugar transporter family, which includes the seven glucose transporters (GLUT1-GLUT7) found in humans that are required for the transport of glucose and other hexose sugars. These glucose transport proteins have unique tissue distributions and physiological functions. GLUT1 provides many cell types with their basal glucose requirements and transports glucose across epithelial and endothelial barrier tissues; GLUT2 facilitates glucose uptake or efflux from the liver; GLUT3 regulates glucose supply to neurons; GLUT4 is responsible for insulin-regulated glucose disposal; and GLUT5 regulates fructose uptake into skeletal muscle. Defects in glucose transporters are involved in a recently identified neurological syndrome causing infantile seizures and developmental delay, as well as glycogen storage disease, Fanconi-Bickel syndrome, and non-insulin-dependent diabetes mellitus (Mueckler, M. (1994) *Eur. J. Biochem.* 219:713-725; Longo, N. and L.J. Elsas (1998) *Adv. Pediatr.* 45:293-313).

Monocarboxylate anion transporters are proton-coupled symporters with a broad substrate specificity that includes L-lactate, pyruvate, and the ketone bodies acetate, acetoacetate, and beta-hydroxybutyrate. At least seven isoforms have been identified to date. The isoforms are predicted to have twelve transmembrane (TM) helical domains with a large intracellular loop between TM6 and TM7, and play a critical role in maintaining intracellular pH by removing the protons that are produced stoichiometrically with lactate during glycolysis. The best characterized H^+ -monocarboxylate transporter is that of the erythrocyte membrane, which transports L-lactate and a wide range of other aliphatic monocarboxylates. Other cells possess H^+ -linked monocarboxylate transporters with differing substrate and inhibitor selectivities. In particular, cardiac muscle and tumor cells have transporters that differ in their K_m values for certain substrates, including stereoselectivity for L- over D-lactate, and in their sensitivity to inhibitors. There are Na^+ -monocarboxylate cotransporters on the luminal surface of intestinal and kidney epithelia, which allow the uptake of lactate, pyruvate, and ketone bodies in these tissues. In addition, there are specific and selective transporters for organic cations and organic anions in organs including the

kidney, intestine and liver. Organic anion transporters are selective for hydrophobic charged molecules with electron-attracting side groups. Organic cation transporters, such as the ammonium transporter, mediate the secretion of a variety of drugs and endogenous metabolites, and contribute to the maintenance of intercellular pH (Poole, R.C. and A.P. Halestrap (1993) *Am. J. Physiol.* 264:C761-C782; Price, N.T. et al. (1998) *Biochem. J.* 329:321-328; and Martinelle, K. and I. Haggström (1993) *J. Bioelectrol.* 30:339-350).

ATP-binding cassette (ABC) transporters are members of a superfamily of membrane proteins that transport substances ranging from small molecules such as ions, sugars, amino acids, peptides, and phospholipids, to lipopeptides, large proteins, and complex hydrophobic drugs. ABC transporters consist of four modules: two nucleotide-binding domains (NBD), which hydrolyze ATP to supply the energy required for transport, and two membrane-spanning domains (MSD), each containing six putative transmembrane segments. These four modules may be encoded by a single gene, as is the case for the cystic fibrosis transmembrane regulator (CFTR), or by separate genes. When encoded by separate genes, each gene product contains a single NBD and MSD. These "half-molecules" form homo- and heterodimers, such as Tap1 and Tap2, the endoplasmic reticulum-based major histocompatibility (MHC) peptide transport system. Several genetic diseases are attributed to defects in ABC transporters, such as the following diseases and their corresponding proteins: cystic fibrosis (CFTR, an ion channel), adrenoleukodystrophy (adrenoleukodystrophy protein, ALDP), Zellweger syndrome (peroxisomal membrane protein-70, PMP70), and hypervitaminosis hypoglycemia (sulfonyleurea receptor, SUR). Overexpression of the multidrug resistance (MDR) protein, another ABC transporter, in human cancer cells makes the cells resistant to a variety of cytotoxic drugs used in chemotherapy (Taglicht, D. and S. Michaelis (1998) *Meth. Enzymol.* 292:130-162).

A number of metal ions such as iron, zinc, copper, cobalt, manganese, molybdenum, selenium, nickel, and chromium are important as cofactors for a number of enzymes. For example, copper is involved in hemoglobin synthesis, connective tissue metabolism, and bone development, by acting as a cofactor in oxidoreductases such as superoxide dismutase, ferroxidase (ceruloplasmin), and lysyl oxidase. Copper and other metal ions must be provided in the diet, and are absorbed by transporters in the gastrointestinal tract. Plasma proteins transport the metal ions to the liver and other target organs, where specific transporters move the ions into cells and cellular organelles as needed. Imbalances in metal ion metabolism have been associated with a number of disease states (Danks, D.M. (1986) *J. Med. Genet.* 23:99-106).

Transport of fatty acids across the plasma membrane can occur by diffusion, a high capacity, low affinity process. However, under normal physiological conditions a significant fraction of fatty acid transport appears to occur via a high affinity, low capacity protein-mediated transport process.

Fatty acid transport protein (FATP), an integral membrane protein with four transmembrane segments, is expressed in tissues exhibiting high levels of plasma membrane fatty acid flux, such as muscle, heart, and adipose. Expression of FATP is upregulated in 3T3-L1 cells during adipose conversion, and expression in COS7 fibroblasts elevates uptake of long-chain fatty acids (Hui, T.Y. et al. (1998) *J. Biol. Chem.* 273:27420-27429).

Mitochondrial carrier proteins are transmembrane-spanning proteins which transport ions and charged metabolites between the cytosol and the mitochondrial matrix. Examples include the ADP, ATP carrier protein; the 2-oxoglutarate/malate carrier; the phosphate carrier protein; the pyruvate carrier; the dicarboxylate carrier which transports malate, succinate, fumarate, and phosphate; the tricarboxylate carrier which transports citrate and malate; and the Grave's disease carrier protein, a protein recognized by IgG in patients with active Grave's disease, an autoimmune disorder resulting in hyperthyroidism. Proteins in this family consist of three tandem repeats of an approximately 100 amino acid domain, each of which contains two transmembrane regions (Stryer, L. (1995) *Biochemistry*, W.H. Freeman and Company, New York, NY, p. 551; PROSITE PD000189).

Mitochondrial energy transfer proteins signature; Online Mendelian Inheritance in Man (OMIM) *275000 Graves Disease).

This class of transporters also includes the mitochondrial uncoupling proteins, which create proton leaks across the inner mitochondrial membrane, thus uncoupling oxidative phosphorylation from ATP synthesis. The result is energy dissipation in the form of heat. Mitochondrial uncoupling proteins have been implicated as modulators of thermoregulation and metabolic rate, and have been proposed as potential targets for drugs against metabolic diseases such as obesity (Riquier, D. et al. (1999) *J. Int. Med.* 245:637-642).

Ion Channels

The electrical potential of a cell is generated and maintained by controlling the movement of ions across the plasma membrane. The movement of ions requires ion channels, which form ion-selective pores within the membrane. There are two basic types of ion channels, ion transporters and gated ion channels. Ion transporters utilize the energy obtained from ATP hydrolysis to actively transport an ion against the ion's concentration gradient. Gated ion channels allow passive flow of an ion down the ion's electrochemical gradient under restricted conditions. Together, these types of ion channels generate, maintain, and utilize an electrochemical gradient that is used in 1) electrical impulse conduction down the axon of a nerve cell, 2) transport of molecules into cells against concentration gradients, 3) initiation of muscle contraction, and 4) endocrine cell secretion.

Ion Transporters

Ion transporters generate and maintain the resting electrical potential of a cell. Utilizing the energy derived from ATP hydrolysis, they transport ions against the ion's concentration gradient.

These transmembrane ATPases are divided into three families. The phosphorylated (P) class ion transporters, including Na⁺-K⁺ ATPase, Ca²⁺-ATPase, and H⁺-ATPase, are activated by a phosphorylation event. P-class ion transporters are responsible for maintaining resting potential distributions such that cytosolic concentrations of Na⁺ and Ca²⁺ are low and cytosolic concentration of K⁺ is high. The vacuolar (V) class of ion transporters includes H⁺ pumps on intracellular organelles, such as lysosomes and Golgi. V-class ion transporters are responsible for generating the low pH within the lumen of these organelles that is required for function. The coupling factor (F) class consists of H⁺ pumps in the mitochondria. F-class ion transporters utilize a proton gradient to generate ATP from ADP and inorganic phosphate (P_i).

The P-ATPases are hexamers of a 100 kD subunit with ten transmembrane domains and several large cytoplasmic regions that may play a role in ion binding (Scarborough, G.A. (1999) *Curr. Opin. Cell Biol.* 11:517-522). The V-ATPases are composed of two functional domains: the V₁ domain, a peripheral complex responsible for ATP hydrolysis; and the V₀ domain, an integral complex responsible for proton translocation across the membrane. The F-ATPases are structurally and evolutionarily related to the V-ATPases. The F-ATPase F₀ domain contains 12 copies of the c subunit, a highly hydrophobic protein composed of two transmembrane domains and containing a single buried carboxyl group in TM2 that is essential for proton transport. The V-ATPase V₀ domain contains three types of homologous c subunits with four or five transmembrane domains and the essential carboxyl group in TM4 or TM3. Both types of complex also contain a single a subunit that may be involved in regulating the pH dependence of activity (Forgac, M. (1999) *J. Biol. Chem.* 274:12951-12954).

The resting potential of the cell is utilized in many processes involving carrier proteins and gated ion channels. Carrier proteins utilize the resting potential to transport molecules into and out of the cell. Amino acid and glucose transport into many cells is linked to sodium ion co-transport (symport) so that the movement of Na⁺ down an electrochemical gradient drives transport of the other molecule up a concentration gradient. Similarly, cardiac muscle links transfer of Ca²⁺ out of the cell with transport of Na⁺ into the cell (antiport).

Gated Ion Channels

Gated ion channels control ion flow by regulating the opening and closing of pores. The ability to control ion flux through various gating mechanisms allows ion channels to mediate such diverse signaling and homeostatic functions as neuronal and endocrine signaling, muscle contraction, fertilization, and regulation of ion and pH balance. Gated ion channels are categorized according to the manner of regulating the gating function. Mechanically-gated channels open their pores in response to mechanical stress; voltage-gated channels (e.g., Na⁺, K⁺, Ca²⁺, and Cl⁻ channels) open their pores in response to changes in membrane potential; and ligand-gated channels (e.g.,

acetylcholine-, serotonin-, and glutamate-gated cation channels, and GABA- and glycine-gated chloride channels) open their pores in the presence of a specific ion, nucleotide, or neurotransmitter. The gating properties of a particular ion channel (i.e., its threshold for and duration of opening and closing) are sometimes modulated by association with auxiliary channel proteins and/or post translational modifications, such as phosphorylation.

Mechanically-gated or mechanosensitive ion channels act as transducers for the senses of touch, hearing, and balance, and also play important roles in cell volume regulation, smooth muscle contraction, and cardiac rhythm generation. A stretch-inactivated channel (SIC) was recently cloned from rat kidney. The SIC channel belongs to a group of channels which are activated by pressure or stress on the cell membrane and conduct both Ca²⁺ and Na⁺ (Suzuki, M. et al. (1999) *J. Biol. Chem.* 274:6330-6335).

The pore-forming subunits of the voltage-gated cation channels form a superfamily of ion channel proteins. The characteristic domain of these channel proteins comprises six transmembrane domains (S1-S6), a pore-forming region (P) located between S5 and S6, and intracellular amino and carboxy termini. In the Na⁺ and Ca²⁺ subfamilies, this domain is repeated four times, while in the K⁺ channel subfamily, each channel is formed from a tetramer of either identical or dissimilar subunits. The P region contains information specifying the ion selectivity for the channel. In the case of K⁺ channels, a GYG tripeptide is involved in this selectivity (Ishii, T.M. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:11651-11656).

Voltage-gated Na⁺ and K⁺ channels are necessary for the function of electrically excitable cells, such as nerve and muscle cells. Action potentials, which lead to neurotransmitter release and muscle contraction, arise from large, transient changes in the permeability of the membrane to Na⁺ and K⁺ ions. Depolarization of the membrane beyond the threshold level opens voltage-gated Na⁺ channels. Sodium ions flow into the cell, further depolarizing the membrane and opening more voltage-gated Na⁺ channels, which propagates the depolarization down the length of the cell. Depolarization also opens voltage-gated potassium channels. Consequently, potassium ions flow outward, which leads to repolarization of the membrane. Voltage-gated channels utilize charged residues in the fourth transmembrane segment (S4) to sense voltage change. The open state lasts only about 1 millisecond, at which time the channel spontaneously converts into an inactive state that cannot be opened irrespective of the membrane potential. Inactivation is mediated by the channel's N-terminus, which acts as a plug that closes the pore. The transition from an inactive to a closed state requires a return to resting potential.

Voltage-gated Na⁺ channels are heterotrimeric complexes composed of a 260 kDa pore-forming α subunit that associates with two smaller auxiliary subunits, β 1 and β 2. The β 2 subunit is a integral membrane glycoprotein that contains an extracellular Ig domain, and its association with α

and $\beta 1$ subunits correlates with increased functional expression of the channel, a change in its gating properties, as well as an increase in whole cell capacitance due to an increase in membrane surface area (Isom, L.L. et al. (1995) *Cell* 83:433-442).

Non voltage-gated Na^+ channels include the members of the amiloride-sensitive Na^+

channel/degenerin (NaCDEG) family. Channel subunits of this family are thought to consist of two transmembrane domains flanking a long extracellular loop, with the amino and carboxyl termini located within the cell. The NaCDEG family includes the epithelial Na^+ channel (ENaC) involved in Na^+ reabsorption in epithelia including the airway, distal colon, cortical collecting duct of the kidney, and exocrine duct glands. Mutations in ENaC result in pseudohypoaldosteronism type 1 and Liddle's syndrome (pseudohypoaldosteronism). The NaCDEG family also includes the recently characterized H^+ -gated cation channels or acid-sensing ion channels (ASIC). ASIC subunits are expressed in the brain and form heteromultimeric Na^+ -permeable channels. These channels require acid pH fluctuations for activation. ASIC subunits show homology to the degenerins, a family of mechanically-gated channels originally isolated from *C. elegans*. Mutations in the degenerins cause neurodegeneration. ASIC subunits may also have a role in neuronal function, or in pain perception, since tissue acidosis causes pain (Waldmann, R. and M. Lazdunski (1998) *Curr. Opin. Neurobiol.* 8:418-424; Egleon, R.M. et al. (1999) *Trends Pharmacol. Sci.* 20:337-342).

K^+ channels are located in all cell types, and may be regulated by voltage, ATP

concentration, or second messengers such as Ca^{2+} and cAMP. In non-excitable tissue, K^+ channels are involved in protein synthesis, control of endocrine secretions, and the maintenance of osmotic equilibrium across membranes. In neurons and other excitable cells, in addition to regulating action potentials and repolarizing membranes, K^+ channels are responsible for setting resting membrane potential. The cytosol contains non-diffusible anions and, to balance this net negative charge, the cell contains a Na^+ - K^+ pump and ion channels that provide the redistribution of Na^+ , K^+ , and Cl^- . The pump actively transports Na^+ out of the cell and K^+ into the cell in a 3:2 ratio. Ion channels in the plasma membrane allow K^+ and Cl^- to flow by passive diffusion. Because of the high negative charge within the cytosol, Cl^- flows out of the cell. The flow of K^+ is balanced by an electromotive force pulling K^+ into the cell, and a K^+ concentration gradient pushing K^+ out of the cell. Thus, the resting membrane potential is primarily regulated by K^+ flow (Salkoff, L., and T. Jegh (1995) *Neuron* 15:489-492).

Potassium channel subunits of the Shaker-like superfamily all have the characteristic six transmembrane/1 pore domain structure. Four subunits combine as homo- or heterotetramers to form functional K^+ channels. These pore-forming subunits also associate with various cytoplasmic β subunits that alter channel inactivation kinetics. The Shaker-like channel family includes the voltage-gated K^+ channels as well as the delayed rectifier type channels such as the human ether-a-go-go

related gene (HERG) associated with long QT, a cardiac dysrhythmia syndrome (Curran, M.E. (1998) *Curr. Opin. Biotechnol.* 9:565-572; Kaczorowski, G.J. and M.L. Garcia (1999) *Curr. Opin. Chem. Biol.* 3:448-458).

A second superfamily of K^+ channels is composed of the inward rectifying channels (Kir).

Kir channels have the property of preferentially conducting K^+ currents in the inward direction. These proteins consist of a single potassium selective pore domain and two transmembrane domains, which correspond to the fifth and sixth transmembrane domains of voltage-gated K^+ channels. Kir subunits also associate as tetramers. The Kir family includes ROMK1, mutations in which lead to Bartter syndrome, a renal tubular disorder. Kir channels are also involved in regulation of cardiac pacemaker activity, seizures and epilepsy, and insulin regulation (Doupnik, C.A. et al. (1995) *Curr. Opin. Neurobiol.* 5:268-277; Curran, *supra*).

The recently recognized TWIK K^+ channel family includes the mammalian TWIK-1, TREK-1 and TASK proteins. Members of this family possess an overall structure with four transmembrane domains and two P domains. These proteins are probably involved in controlling the resting potential in a large set of cell types (Duprat, F. et al. (1997) *EMBO J* 16:5464-5471).

The voltage-gated Ca^{2+} channels have been classified into several subtypes based upon their electrophysiological and pharmacological characteristics. L-type Ca^{2+} channels are predominantly expressed in heart and skeletal muscle where they play an essential role in excitation-contraction coupling. T-type channels are important for cardiac pacemaker activity, while N-type and P/Q-type channels are involved in the control of neurotransmitter release in the central and peripheral nervous system. The L-type and N-type voltage-gated Ca^{2+} channels have been purified and, though their functions differ dramatically, they have similar subunit compositions. The channels are composed of three subunits. The α_1 subunit forms the membrane pore and voltage sensor, while the α_2 and β subunits modulate the voltage-dependence, gating properties, and the current amplitude of the channel. These subunits are encoded by at least six α , one α_2 , and four β genes. A fourth subunit, γ , has been identified in skeletal muscle (Walker, D. et al. (1998) *J. Biol. Chem.* 273:2361-2367; McCleskey, E.W. (1994) *Curr. Opin. Neurobiol.* 4:304-312).

The transient receptor family (Trp) of calcium ion channels are thought to mediate capacitative calcium entry (CCE). CCE is the Ca^{2+} influx into cells to resupply Ca^{2+} stores depleted by the action of inositol triphosphate (IP_3) and other agents in response to numerous hormones and growth factors. Trp and Trp-like were first cloned from *Drosophila* and have similarity to voltage-gated Ca^{2+} channels in the S3 through S6 regions. This suggests that Trp and/or related proteins may form mammalian CCC entry channels (Zhu, X. et al. (1996) *Cell* 85:661-671; Boulay, G. et al. (1997) *J. Biol. Chem.* 272:29672-29680). Melastatin is a gene isolated in both the mouse and human, and whose expression in melanoma cells is inversely correlated with melanoma aggressiveness in vivo.

The human cDNA transcript corresponds to a 1533-amino acid protein having homology to members of the Trp family. It has been proposed that the combined use of malastatin mRNA expression status and tumor thickness might allow for the determination of subgroups of patients at both low and high risk for developing metastatic disease (Duncan, L.M. et al (2001) J. Clin. Oncol. 19:568-576).

Chloride channels are necessary in endocrine secretion and in regulation of cytosolic and organelle pH. In secretory epithelial cells, Cl⁻ enters the cell across a basolateral membrane through an Na⁺, K⁺/Cl⁻ cotransporter, accumulating in the cell above its electrochemical equilibrium concentration. Secretion of Cl⁻ from the apical surface, in response to hormonal stimulation, leads to flow of Na⁺ and water into the secretory lumen. The cystic fibrosis transmembrane conductance regulator (CFTR) is a chloride channel encoded by the gene for cystic fibrosis, a common fatal genetic disorder in humans. CFTR is a member of the ABC transporter family, and is composed of two domains each consisting of six transmembrane domains followed by a nucleotide-binding site. Loss of CFTR function decreases transepithelial water secretion and, as a result, the layers of mucus that coat the respiratory tree, pancreatic ducts, and intestine are dehydrated and difficult to clear. The resulting blockage of these sites leads to pancreatic insufficiency, "meconium ileus", and devastating "chronic obstructive pulmonary disease" (Al-Awqati, Q. et al. (1992) J. Exp. Biol. 172:245-266).

The voltage-gated chloride channels (CLC) are characterized by 10-12 transmembrane domains, as well as two small globular domains known as CBS domains. The CLC subunits probably function as homotetramers. CLC proteins are involved in regulation of cell volume, membrane potential stabilization, signal transduction, and transepithelial transport. Mutations in CLC-1, expressed predominantly in skeletal muscle, are responsible for autosomal recessive generalized myotonia and autosomal dominant myotonia congenita, while mutations in the kidney channel CLC-5 lead to kidney stones (Jentsch, T.J. (1996) Curr. Opin. Neurobiol. 6:303-310).

Ligand-gated channels open their pores when an extracellular or intracellular mediator binds to the channel. Neurotransmitter-gated channels are channels that open when a neurotransmitter binds to their extracellular domain. These channels exist in the postsynaptic membrane of nerve or muscle cells. There are two types of neurotransmitter-gated channels. Sodium channels open in response to excitatory neurotransmitters, such as acetylcholine, glutamate, and serotonin. This opening causes an influx of Na⁺ and produces the initial localized depolarization that activates the voltage-gated channels and starts the action potential. Chloride channels open in response to inhibitory neurotransmitters, such as γ -aminobutyric acid (GABA) and glycine, leading to hyperpolarization of the membrane and the subsequent generation of an action potential. Neurotransmitter-gated ion channels have four transmembrane domains and probably function as pentamers (Jentsch, *supra*). Amino acids in the second transmembrane domain appear to be important in determining channel permeation and selectivity (Sather, W.A. et al. (1994) Curr. Opin. Neurobiol.

4:313-323).

Ligand-gated channels can be regulated by intracellular second messengers. For example, calcium-activated K⁺ channels are gated by internal calcium ions. In nerve cells, an influx of calcium during depolarization opens K⁺ channels to modulate the magnitude of the action potential (Ishii et al., *supra*). The large conductance (BK) channel has been purified from brain and its subunit composition determined. The α subunit of the BK channel has seven rather than six transmembrane domains in contrast to voltage-gated K⁺ channels. The extra transmembrane domain is located at the subunit N-terminus. A 28-amino-acid stretch in the C-terminal region of the subunit (the "calcium bowl" region) contains many negatively charged residues and is thought to be the region responsible for calcium binding. The β subunit consists of two transmembrane domains connected by a glycosylated extracellular loop, with intracellular N- and C-termini (Kaczorowski, *supra*; Vergara, C. et al. (1998) Curr. Opin. Neurobiol. 8:321-329).

Cyclic nucleotide-gated (CNG) channels are gated by cytosolic cyclic nucleotides. The best examples of these are the cAMP-gated Na⁺ channels involved in olfaction and the cGMP-gated cation channels involved in vision. Both systems involve ligand-mediated activation of a G-protein coupled receptor which then alters the level of cyclic nucleotide within the cell. CNG channels also represent a major pathway for Cu²⁺ entry into neurons, and play roles in neuronal development and plasticity. CNG channels are tetramers containing at least two types of subunits, an α subunit which can form functional homomeric channels, and a β subunit, which modulates the channel properties. All CNG subunits have six transmembrane domains and a pore forming region between the fifth and sixth transmembrane domains, similar to voltage-gated K⁺ channels. A large C-terminal domain contains a cyclic nucleotide binding domain, while the N-terminal domain confers variation among channel subtypes (Zufall, F. et al. (1997) Curr. Opin. Neurobiol. 7:404-412).

The activity of other types of ion channel proteins may also be modulated by a variety of intracellular signalling proteins. Many channels have sites for phosphorylation by one or more protein kinases including protein kinase A, protein kinase C, tyrosine kinase, and casein kinase II, all of which regulate ion channel activity in cells. Kir channels are activated by the binding of the G $\beta\gamma$ subunits of heterotrimeric G-proteins (Reimann, F. and F.M. Ashcroft (1999) Curr. Opin. Cell Biol. 11:503-508). Other proteins are involved in the localization of ion channels to specific sites in the cell membrane. Such proteins include the PDZ domain proteins known as MAGUKs (membrane-associated guanylate kinases) which regulate the clustering of ion channels at neuronal synapses (Craven, S.E. and D.S. Bredt (1998) Cell 93:495-498).

Disease Correlation

The etiology of numerous human diseases and disorders can be attributed to defects in the

transport of molecules across membranes. Defects in the trafficking of membrane-bound transporters and ion channels are associated with several disorders, e.g., cystic fibrosis, glucose-galactose malabsorption syndrome, hypercholesterolemia, von Gierke disease, and certain forms of diabetes mellitus. Single-gene defect diseases resulting in an inability to transport small molecules across membranes include, e.g., cystinuria, iminoglycinuria, Hartup disease, and Fanconi disease (Van Hoff, W.G. (1996) *Exp. Nephrol.* 4:253-262; Talemie, G.M. et al. (1994) *Ann. Intern. Med.* 120:218-226; and Chillon, M. et al. (1995) *New Engl. J. Med.* 332:1475-1480).

Human diseases caused by mutations in ion channel genes include disorders of skeletal muscle, cardiac muscle, and the central nervous system. Mutations in the pore-forming subunits of sodium and chloride channels cause myotonia, a muscle disorder in which relaxation after voluntary contraction is delayed. Sodium channel myotonias have been treated with channel blockers.

Mutations in muscle sodium and calcium channels cause forms of periodic paralysis, while mutations in the sarcolemmal calcium release channel, T-tubule calcium channel, and muscle sodium channel cause malignant hyperthermia. Cardiac arrhythmia disorders such as the long QT syndromes and idiopathic ventricular fibrillation are caused by mutations in potassium and sodium channels (Cooper, E.C. and L.Y. Jan (1998) *Proc. Natl. Acad. Sci. USA* 96:4759-4766). All four known human idiopathic epilepsy genes code for ion channel proteins (Berkovic, S.F. and J.E. Scheffer (1999) *Curr. Opin. Neurology* 12:177-182). Other neurological disorders such as ataxia, hemiplegic migraine and hereditary deafness can also result from mutations in ion channel genes (Jen, J. (1999) *Curr. Opin. Neurobiol.* 9:274-280; Cooper, supra).

Ion channels have been the target for many drug therapies. Neurotransmitter-gated channels have been targeted in therapies for treatment of insomnia, anxiety, depression, and schizophrenia. Voltage-gated channels have been targeted in therapies for arrhythmia, ischemic stroke, head trauma, and neurodegenerative disease (Taylor, C.P. and L.S. Narasimhan (1997) *Adv. Pharmacol.* 39:47-98). Various classes of ion channels also play an important role in the perception of pain, and thus are potential targets for new analgesics. These include the vanilloid-gated ion channels, which are activated by the vanilloid capsaicin, as well as by noxious heat. Local anesthetics such as lidocaine and mexiletine which block voltage-gated Na⁺ channels have been useful in the treatment of neuropathic pain (Eglen, supra).

Ion channels in the immune system have recently been suggested as targets for immunomodulation. T-cell activation depends upon calcium signaling, and a diverse set of T-cell specific ion channels has been characterized that affect this signaling process. Channel blocking agents can inhibit secretion of lymphokines, cell proliferation, and killing of target cells. A peptide antagonist of the T-cell potassium channel Kv1.3 was found to suppress delayed-type hypersensitivity and allogenic responses in pigs, validating the idea of channel blockers as safe and efficacious

immunosuppressants (Cahalan, M.D. and K.G. Chanly (1997) *Curr. Opin. Biotechnol.* 8:749-756). The discovery of new transporters and ion channels, and the polynucleotides encoding them, satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of transport, neurological, muscle, immunological, and cell proliferative disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of transporters and ion channels.

SUMMARY OF THE INVENTION

The invention features purified polypeptides, transporters and ion channels, referred to collectively as "TRICH" and individually as "TRICH-1," "TRICH-2," "TRICH-3," "TRICH-4," "TRICH-5," "TRICH-6," "TRICH-7," "TRICH-8," "TRICH-9," "TRICH-10," "TRICH-11," "TRICH-12," "TRICH-13," "TRICH-14," "TRICH-15," "TRICH-16," "TRICH-17," "TRICH-18," "TRICH-19," "TRICH-20," "TRICH-21," "TRICH-22," "TRICH-23," "TRICH-24," "TRICH-25," "TRICH-26," "TRICH-27," "TRICH-28," "TRICH-29," and "TRICH-30." In one aspect, the invention provides an isolated polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-30.

The invention further provides an isolated polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-30. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:31-60.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting

of SEQ ID NO:1-30, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30.

The invention further provides an isolated polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:31-60, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:31-60, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group

consisting of SEQ ID NO:31-60, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:31-60, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:31-60, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:31-60, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, and a pharmaceutically acceptable excipient. In one embodiment, the composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-30. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional TRICH, comprising administering to a patient in need of such treatment the composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino

acid sequence selected from the group consisting of SEQ ID NO:1-30, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional TRIC^h, comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional TRIC^h, comprising administering to a patient in need of such treatment the composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a polynucleotide sequence selected from the group consisting of SEQ ID NO:31-60, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:31-60, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:31-60, iii) a polynucleotide having a sequence complementary to i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:31-60, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:31-60, iii) a polynucleotide complementary to the polynucleotide of ii), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Alternatively, the target

polynucleotide comprises a fragment of a polynucleotide sequence selected from the group consisting of i-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

BRIEF DESCRIPTION OF THE TABLES

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the present invention.

Table 2 shows the GenBank identification number and annotation of the nearest GenBank homolog for polypeptides of the invention. The probability score for the match between each polypeptide and its GenBank homolog is also shown.

Table 3 shows structural features of polypeptide sequences of the invention, including predicted motifs and domains, along with the methods, algorithms, and searchable databases used for analysis of the polypeptides.

Table 4 lists the cDNA and/or genomic DNA fragments which were used to assemble polynucleotide sequences of the invention, along with selected fragments of the polynucleotide sequences.

Table 5 shows the representative cDNA library for polynucleotides of the invention.

Table 6 provides an appendix which describes the tissues and vectors used for construction of the cDNA libraries shown in Table 5.

Table 7 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so

forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

"TRICH" refers to the amino acid sequences of substantially purified TRICH obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of TRICH. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of TRICH either by directly interacting with TRICH or by acting on components of the biological pathway in which TRICH participates.

An "allelic variant" is an alternative form of the gene encoding TRICH. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding TRICH include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as TRICH or a polypeptide with at least one functional characteristic of TRICH. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding TRICH, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding TRICH. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent TRICH. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of TRICH is retained. For example,

negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophobicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophobicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of TRICH. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of TRICH either by directly interfering with TRICH or by acting on components of the biological pathway in which TRICH participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab)₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind TRICH polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired.

Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA;

RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic TRICH, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding TRICH or fragments of TRICH may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows

amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
5	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
10	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
15	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
20	Thr	Ser, Val
	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide.

Chemical modifications of a polynucleotide can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

"Differential expression" refers to increased or upregulated; or decreased, downregulated, or absent gene or protein expression, determined by comparing at least two different samples. Such comparisons may be carried out between, for example, a treated and an untreated sample, or a diseased and a normal sample.

"Exon shuffling" refers to the recombination of different coding regions (exons). Since an exon may represent a structural or functional domain of the encoded protein, new proteins may be assembled through the novel reassortment of stable substructures, thus allowing acceleration of the evolution of new protein functions.

A "fragment" is a unique portion of TRICH or the polynucleotide encoding TRICH which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50%) of a polypeptide as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:31-60 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:31-60, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:31-60 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:31-60 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:31-60 and the region of SEQ ID NO:31-60 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purposes for the fragment.

A fragment of SEQ ID NO:1-30 is encoded by a fragment of SEQ ID NO:31-60. A fragment of SEQ ID NO:1-30 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-30. For example, a fragment of SEQ ID NO:1-30 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-30. The precise length of a fragment of SEQ ID NO:1-30 and the region of SEQ ID NO:1-30 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A "full length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full length" polynucleotide sequence encodes a "full length" polypeptide sequence.

"Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

5 The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

10 Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequences.

15 Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/ncbi/blast2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Reward for match: 1

30 *Penalty for mismatch: -2*

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 11

35 *Filter: on*

23

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

10 Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

15 The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

25 Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

30 *Open Gap: 11 and Extension Gap: 1 penalties*

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

35 Percent identity may be measured over the length of an entire defined polypeptide sequence,

24

for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC

concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%.

Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C₄t or R₄t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of TRICH which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of TRICH which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of TRICH. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of TRICH.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the

antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Openly linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Openly linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition.

PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an TRICH may involve lipidation, glycosylation, phosphorylation, acetylation, trypsinization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications may vary by cell type depending on the enzymatic milieu of TRICH.

"Probe" refers to nucleic acid sequences encoding TRICH, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

"Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) *Current Protocols in Molecular Biology*, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) *PCR Protocols, A Guide to Methods and Applications*, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that

purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, *supra*. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is

expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing TRICH, nucleic acids encoding TRICH, or fragments thereof may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" refers to the collective pattern of gene expression by a particular cell

type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed cells" includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or *in vitro* fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook et al. (1989), *supra*.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to

another. The resulting polypeptides will generally have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length of one of the polypeptides.

15 THE INVENTION

The invention is based on the discovery of new human transporters and ion channels (TRICCH), the polynucleotides encoding TRICCH, and the use of these compositions for the diagnosis, treatment, or prevention of transport, neurological, muscle, immunological, and cell proliferative disorders.

20 Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the invention. Each polynucleotide and its corresponding polypeptide are correlated to a single Incyte project identification number (Incyte Project ID). Each polypeptide sequence is denoted by both a polypeptide sequence identification number (Polypeptide SEQ ID NO.) and an Incyte polypeptide sequence number (Incyte Polypeptide ID) as shown. Each polynucleotide sequence is denoted by both a polynucleotide sequence identification number (Polynucleotide SEQ ID NO.) and an Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) as shown.

25 Table 2 shows sequences with homology to the polypeptides of the invention as identified by BLAST analysis against the GenBank protein (genpept) database. Columns 1 and 2 show the polypeptide sequence identification number (Polypeptide SEQ ID NO.) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for polypeptides of the invention. Column 3 shows the GenBank identification number (Genbank ID NO.) of the nearest GenBank homolog. Column 4 shows the probability score for the match between each polypeptide and its GenBank homolog. Column 5 shows the annotation of the GenBank homolog along with relevant citations where applicable, all of which are expressly incorporated by reference herein.

35 Table 3 shows various structural features of the polypeptides of the invention. Columns 1

and 2 show the polypeptide sequence identification number (SEQ ID NO.) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for each polypeptide of the invention. Column 3 shows the number of amino acid residues in each polypeptide. Column 4 shows potential phosphorylation sites, and column 5 shows potential glycosylation sites, as determined by the MOTIFS program of the GCG sequence analysis software package (Genetics Computer Group, Madison WI). Column 6 shows amino acid residues comprising signature sequences, domains, and motifs. Column 7 shows analytical methods for protein structure/function analysis and in some cases, searchable databases to which the analytical methods were applied.

Together, Tables 2 and 3 summarize the properties of polypeptides of the invention, and these properties establish that the claimed polypeptides are transporters and ion channels. For example, SEQ ID NO:6 is 89% identical to rat neuronal nicotinic acetylcholine receptor subunit (GenBank ID g6746563) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 1.7e-188, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:6 also contains a neurotransmitter-gated ion channel domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESSCAN analyses provide further corroborative evidence that SEQ ID NO:6 is a neurotransmitter-gated ion channel. In an alternative example, SEQ ID NO:14 is 93% identical to rat TAP-like ABC transporter (GenBank ID g6045150) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 0.0, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:14 also contains an ABC transporter domain and an ABC transporter transmembrane region as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESSCAN analyses provide further corroborative evidence that SEQ ID NO:14 is an ABC transporter. In an alternative example, SEQ ID NO:16 is 98% identical to human voltage-dependent anion channel (GenBank ID g340199) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 1.2e-130, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:16 also contains a eukaryotic porin active site domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESSCAN analyses provide further corroborative evidence that SEQ ID NO:16 is a mitochondrial porin. In an alternative example, SEQ ID NO:20 is 28% identical to a rat voltage-gated calcium channel (GenBank ID g4386963) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The

BLAST probability score is 2.4e-27, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. Data from BLIMPS and BLAST analyses provide further corroborative evidence that SEQ ID NO:20 is a voltage-gated calcium channel. In an alternative example, SEQ ID NO:22 is 82% identical to human inhibitory glycine receptor (GenBank ID g31849) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 1.1e-175, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:22 also contains a neurotransmitter-gated ion channel domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESAN analyses provide further corroborative evidence that SEQ ID NO:30 is 36% identical to human ATP binding cassette (ABC)-C transporter (GenBank ID g1514530) as determined by the Basic Local Alignment Search Tool (BLAST, see Table 2). The BLAST probability score is 2.3e-127, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:30 also contains ABC transporter domains as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains (see Table 3). Data from BLIMPS, MOTIFS, and PROFILESAN analyses provide further corroborative evidence that SEQ ID NO:30 is an ABC transporter. SEQ ID NO:1-5, SEQ ID NO:7-13, SEQ ID NO:15, SEQ ID NO:17-19, SEQ ID NO:21, and SEQ ID NO:23-29 were analyzed and annotated in a similar manner. The algorithms and parameters for the analysis of SEQ ID NO:1-30 are described in Table 7.

As shown in Table 4, the full length polynucleotide sequences of the present invention were assembled using cDNA sequences or coding (exon) sequences derived from genomic DNA, or any combination of these two types of sequences. Columns 1 and 2 list the polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and the corresponding Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) for each polynucleotide of the invention. Column 3 shows the length of each polynucleotide sequence in basepairs. Column 4 lists fragments of the polynucleotide sequences which are useful, for example, in hybridization or amplification technologies that identify SEQ ID NO:31-60 or that distinguish between SEQ ID NO:31-60 and related polynucleotide sequences. Column 5 shows identification numbers corresponding to cDNA sequences, coding sequences (exons) predicted from genomic DNA, and/or sequence assemblages comprised of both cDNA and genomic DNA. These sequences were used to assemble the full length polynucleotide sequences of the invention. Columns 6 and 7 of Table 4 show the nucleotide start (5') and stop (3') positions of the cDNA and/or genomic sequences in column 5 relative to their respective full length sequences.

The identification numbers in Column 5 of Table 4 may refer specifically, for example, to Incyte cDNAs along with their corresponding cDNA libraries. For example, 6340750H1 is the identification number of an Incyte cDNA sequence, and BRANDIN01 is the cDNA library from which it is derived. Incyte cDNAs for which cDNA libraries are not indicated were derived from pooled cDNA libraries (e.g., 7191330V1). Alternatively, the identification numbers in column 5 may refer to GenBank cDNAs or ESTs (e.g., g5110579) which contributed to the assembly of the full length polynucleotide sequences. In addition, the identification numbers in column 5 may identify sequences derived from the ENSEMBL (The Sanger Centre, Cambridge, UK) database (i.e., those sequences including the designation "ENST"). Alternatively, the identification numbers in column 5 may be derived from the NCBI RefSeq Nucleotide Sequence Records Database (i.e., those sequences including the designation "NM" or "NT") or the NCBI RefSeq Protein Sequence Records (i.e., those sequences including the designation "NP"). Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon stitching" algorithm. For example, FL_XXXXXX_N_N_YYYYY_N_N represents a "stitched" sequence in which XXXXX is the identification number of the cluster of sequences to which the algorithm was applied, and YYYYY is the number of the prediction generated by the algorithm, and N_{1,2,3...} if present, represent specific exons that may have been manually edited during analysis (See Example V). Alternatively, the identification numbers in column 5 may refer to assemblages of exons brought together by an "exon-stretching" algorithm. For example,

FL_XXXXXX_gAAAAA_gBBBBB_1_N is the identification number of a "stretched" sequence, with XXXXXX being the Incyte project identification number, gAAAAA being the GenBank identification number of the human genomic sequence to which the "exon-stretching" algorithm was applied, gBBBBB being the GenBank identification number or NCBI RefSeq identification number of the nearest GenBank protein homolog, and N referring to specific exons (See Example V). In instances where a RefSeq sequence was used as a protein homolog for the "exon-stretching" algorithm, a RefSeq identifier (denoted by "NM," "NP," or "NT") may be used in place of the GenBank identifier (i.e., gBBBBB).

Alternatively, a prefix identifies component sequences that were hand-edited, predicted from genomic DNA sequences, or derived from a combination of sequence analysis methods. The following Table lists examples of component sequence prefixes and corresponding sequence analysis methods associated with the prefixes (see Example IV and Example V).

Prefix	Type of analysis and/or examples of programs
--------	--

GNN, GFG, ENST	Exon prediction from genomic sequences using, for example, GENSCAN (Stanford University, CA, USA) or FGENSES (Computer Genomics Group, The Sanger Centre, Cambridge, UK).
GBI	Hand-edited analysis of genomic sequences.
FL	Stitched or stretched genomic sequences (see Example V).
INCY	Full length transcript and exon prediction from mapping of EST sequences to the genome. Genomic location and EST composition data are combined to predict the exons and resulting transcript.

In some cases, Incyte cDNA coverage redundant with the sequence coverage shown in column 5 was obtained to confirm the final consensus polynucleotide sequence, but the relevant Incyte cDNA identification numbers are not shown.

Table 5 shows the representative cDNA libraries for those full length polynucleotide sequences which were assembled using Incyte cDNA sequences. The representative cDNA library is the Incyte cDNA library which is most frequently represented by the Incyte cDNA sequences which were used to assemble and confirm the above polynucleotide sequences. The tissues and vectors which were used to construct the cDNA libraries shown in Table 5 are described in Table 6.

The invention also encompasses TRICH variants. A preferred TRICH variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the TRICH amino acid sequence, and which contains at least one functional or structural characteristic of TRICH.

The invention also encompasses polynucleotides which encode TRICH. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:31-60, which encodes TRICH. The polynucleotide sequences of SEQ ID NO:31-60, as presented in the Sequence Listing, enhance the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding TRICH. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding TRICH. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:31-60 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting

of SEQ ID NO:31-60. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of TRICH.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding TRICH, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring TRICH, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode TRICH and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring TRICH under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding TRICH or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding TRICH and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode TRICH and TRICH derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding TRICH or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:31-60 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied Biosystems), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or

combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding TRICH may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.)

Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060.) Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 primer analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence

into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode TRICH may be cloned in recombinant DNA molecules that direct expression of TRICH, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express TRICH.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter TRICH-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULAR BREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Cramer, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of TRICH, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of

homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding TRICH may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser.* 7:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser.* 7:225-232.)

Alternatively, TRICH itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) *Proteins, Structures and Molecular Properties*, W.H. Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the amino acid sequence of TRICH, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chize, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, *supra*, pp. 28-53.)

In order to express a biologically active TRICH, the nucleotide sequences encoding TRICH or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding TRICH. Such elements may vary in their strength and specificity.

Specific initiation signals may also be used to achieve more efficient translation of sequences encoding TRICH. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding TRICH and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Schauf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression

vectors containing sequences encoding TRICH and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. (See, e.g., Sambrook, J. et al. (1989) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) *Current Protocols in Molecular Biology*, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding TRICH. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, *supra*; Ausubel, *supra*; Van Hecke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509; Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-1945; Takamatsu, N. (1987) *EMBO J.* 6:307-311; *The McGraw Hill Yearbook of Science and Technology* (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659; and Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) *Cancer Gen. Ther.* 5(6):350-356; Yu, M. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90(13):6340-6344; Butler, R.M. et al. (1985) *Nature* 317(6040):813-815; McGregor, D.P. et al. (1994) *Mol. Immunol.* 31(2):219-226; and Verma, I.M. and N. Sonnia (1997) *Nature* 389:239-242.) The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding TRICH. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding TRICH can be achieved using a multifunctional *E. coli* vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or pSPORT1 plasmid (Life Technologies). Ligation of sequences encoding TRICH into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for *in vitro* transcription, diideoxy sequencing, single strand rescue with helper phages, and creation of nested deletions in the cloned sequence. (See, e.g., Van Hecke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509.) When large quantities of TRICH are needed, e.g. for the production of antibodies, vectors which direct high level expression of TRICH may be used. For example, vectors

containing the strong, inducible SP6 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of TRICH. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, *supra*; Bitter, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; and Sorer, C.A. et al. (1994) *BioTechnology* 12:181-184.)

Plant systems may also be used for expression of TRICH. Transcription of sequences encoding TRICH may be driven by viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., *The McGraw Hill Yearbook of Science and Technology* (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding TRICH may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses TRICH in host cells. (See, e.g., Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of TRICH in cell lines is preferred. For example, sequences encoding TRICH can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in

enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *ap^r* cells, respectively. (See, e.g., Wigler, M. et al. (1977) *Cell* 11:223-232; Lowy, I. et al. (1980) *Cell* 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *ds* and *par* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) *Proc. Natl. Acad. Sci. USA* 77:3567-3570; Colbere-Garapin, F. et al. (1981) *J. Mol. Biol.* 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) *Proc. Natl. Acad. Sci. USA* 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) *Methods Mol. Biol.* 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding TRICH is inserted within a marker gene sequence, transformed cells containing sequences encoding TRICH can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding TRICH under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding TRICH and that express TRICH may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR

amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of TRICH using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing

monoclonal antibodies reactive to two non-interfering epitopes on TRICH is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) *Serological Methods, a Laboratory Manual*, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) *Current Protocols in Immunology*, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) *Immunochemical Protocols*, Humana Press, Totowa NJ)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding TRICH include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding TRICH, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding TRICH may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode TRICH may be designed to contain signal sequences which direct secretion of TRICH through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "pro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WID8) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding TRICH may be ligated to a heterologous sequence resulting in translation of a

fusion protein in any of the aforementioned host systems. For example, a chimeric TRICH protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of TRICH activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, c-myc, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, c-myc, and hemagglutinin (HA) enable immunaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the TRICH encoding sequence and the heterologous protein sequence, so that TRICH may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, *supra*, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled TRICH may be achieved *in vitro* using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

TRICH of the present invention or fragments thereof may be used to screen for compounds that specifically bind to TRICH. At least one and up to a plurality of test compounds may be screened for specific binding to TRICH. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of TRICH, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, e.g., Coligan, J.E. et al. (1991) *Current Protocols in Immunology* 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which TRICH binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express TRICH, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing TRICH or cell membrane fractions which contain TRICH are then contacted with a test compound and binding, stimulation, or inhibition of activity of either TRICH or the

compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with TRICH, either in solution or affixed to a solid support, and detecting the binding of TRICH to the compound.

Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

TRICH of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of TRICH. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for TRICH activity, wherein TRICH is combined with at least one test compound, and the activity of TRICH in the presence of a test compound is compared with the activity of TRICH in the absence of the test compound. A change in the activity of TRICH in the presence of the test compound is indicative of a compound that modulates the activity of TRICH. Alternatively, a test compound is combined with an *in vitro* or cell-free system comprising TRICH under conditions suitable for TRICH activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of TRICH may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding TRICH or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent Number 5,175,383 and U.S. Patent Number 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) *Science* 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knock-out a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) *Clin. Invest.* 97:1999-2002; Wagner, K.U. et al. (1997) *Nucleic Acids Res.* 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential

therapeutic or toxic agents.

Polynucleotides encoding TRICH may also be manipulated *in vitro* in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) *Science* 282:1145-1147).

Polynucleotides encoding TRICH can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding TRICH is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress TRICH, e.g., by secreting TRICH in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) *Biotechnol. Annu. Rev.* 4:55-74).

15 THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of TRICH and transporters and ion channels. In addition, the expression of TRICH is closely associated with brain, liver, tumor, colon, thymus, small intestine, myometrium, testicular, bone marrow neuroblastoma tumor, parotid gland, lung, pituitary gland, and placental tissues, and Pompe's disease. Therefore, TRICH appears to play a role in transport, neurological, muscle, immunological, and cell proliferative disorders. In the treatment of disorders associated with increased TRICH expression or activity, it is desirable to decrease the expression or activity of TRICH. In the treatment of disorders associated with decreased TRICH expression or activity, it is desirable to increase the expression or activity of TRICH.

Therefore, in one embodiment, TRICH or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TRICH. Examples of such disorders include, but are not limited to, a transport disorder such as akinesia, amyotrophic lateral sclerosis, ataxia telangiectasia, cystic fibrosis, Becker's muscular dystrophy, Bell's palsy, Charcot-Marie-Tooth disease, diabetes mellitus, diabetes insipidus, diabetic neuropathy, Duchenne muscular dystrophy, hyperkalemic periodic paralysis, normokalemic periodic paralysis, Parkinson's disease, malignant hyperthermia, multidrug resistance, myasthenia gravis, myotonic dystrophy, cataplexy, tardive dyskinesia, dystonias, peripheral neuropathy, cerebral neoplasms, prostate cancer, cardiac disorders associated with transport, e.g., angina, bradyarrhythmia, tachyarrhythmia, hypertension, Long QT syndrome, myocarditis, cardiomyopathy, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, thyrotoxic myopathy,

ethanol myopathy, dermatomyositis, inclusion body myositis, infectious myositis, polymyositis, neurological disorders associated with transport, e.g., Alzheimer's disease, amnesia, bipolar disorder, dementia, depression, epilepsy, Tourette's disorder, paranoid psychoses, and schizophrenia, and other disorders associated with transport, e.g., neurofibromatosis, postherpetic neuralgia, trigeminal neuropathy, sarcoidosis, sickle cell anemia, Wilson's disease, cataracts, infertility, pulmonary artery stenosis, sensorineural autosomal deafness, hyperglycemia, hypoglycemia, Grave's disease, goiter, Cushing's disease, Addison's disease, glucose-galactose malabsorption syndrome, hypercholesterolemia, adrenoleukodystrophy, Zellweger syndrome, Menkes disease, occipital horn syndrome, von Gierke disease, cystinuria, iminoglycinuria, Hartup disease, and Fanconi disease; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Strausler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurothromatosis, tubercous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, cataplexy, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; a muscle disorder such as cardiomyopathy, myocarditis, Duchenne's muscular dystrophy, Becker's muscular dystrophy, myotonic dystrophy, central core disease, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, infectious myositis, polymyositis, dermatomyositis, inclusion body myositis, thyrotoxic myopathy, ethanol myopathy, angina, anaphylactic shock, arrhythmias, asthma, cardiovascular shock, Cushing's syndrome, hypertension, hypoglycemia, myocardial infarction, migraine, pheochromocytoma, and myopathies including encephalopathy, epilepsy, Kearns-Sayre syndrome, lactic acidosis, myoclonic disorder, ophthalmoplegia, and acid malasse deficiency (AMD), also known

as Pompe's disease); an immunological disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, antilysozing spoudyrlitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, alopic dermatitis, dermatomyositis, diabetes mellitus, empty sella, episodic lymphopenia with lymphocytotoxicins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypersensitivity, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polyomyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, trunna, and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus.

In another embodiment, a vector capable of expressing TRICH or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TRICH including, but not limited to, those described above.

In a further embodiment, a composition comprising a substantially purified TRICH in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TRICH including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of TRICH may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TRICH including, but not limited to, those listed above.

In a further embodiment, an antagonist of TRICH may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of TRICH. Examples of such disorders include, but are not limited to, those transport, neurological, muscle, immunological, and cell proliferative disorders described above. In one aspect, an antibody which specifically binds

TRICH may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express TRICH.

In an additional embodiment, a vector expressing the complement of the polynucleotide

encoding TRICH may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of TRICH including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of TRICH may be produced using methods which are generally known in the art. In particular, purified TRICH may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind TRICH. Antibodies to TRICH may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with TRICH or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (*Bacilli Calmette-Guerin*) and *Corynebacterium parvum* are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to TRICH have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of TRICH amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to TRICH may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma

technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce TRICH-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

Antibodies may also be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

Antibody fragments which contain specific binding sites for TRICH may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between TRICH and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering TRICH epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for TRICH. Affinity is expressed as an association constant, K_a, which is defined as the molar concentration of TRICH-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple TRICH epitopes, represents the average affinity, or avidity, of the antibodies

for TRICH. The K_d determined for a preparation of monoclonal antibodies, which are nonspecific for a particular TRICH epitope, represents a true measure of affinity. High-affinity antibody preparations with K_d ranging from about 10^8 to 10^{12} L/mole are preferred for use in immunoassays in which the TRICH-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_d ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of TRICH, preferably in active form, from the antibody (Catty, D. (1988) *Antibodies, Volume I: A Practical Approach*, IRL Press, Washington DC; Liddell, J.E. and A. Cyster (1991) *A Practical Guide to Monoclonal Antibodies*, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of TRICH-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (Sec, e.g., Catty, *supra*, and Coligan et al., *supra*.)

In another embodiment of the invention, the polynucleotides encoding TRICH, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding TRICH. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding TRICH. (Sec, e.g., Agrawal, S., ed. (1996) *Antisense Therapeutics*, Humana Press Inc., Totowa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (Sec, e.g., Shier, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (Sec, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, *supra*; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (Sec, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.*

25(4):2730-2736.)

In another embodiment of the invention, polynucleotides encoding TRICH may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaise, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Sommer (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:11395-11399), hepatitis B or C virus (HBV, HCV), fungal parasites, such as *Candida albicans* and *Paracoccidioides brasiliensis*, and protozoan parasites such as *Plasmodium falciparum* and *Trypanosoma cruzi*). In the case where a genetic deficiency in TRICH expression or regulation causes disease, the expression of TRICH from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in TRICH are treated by constructing mammalian expression vectors encoding TRICH and introducing these vectors by mechanical means into TRICH-deficient cells. Mechanical transfer technologies for use with cells *in vivo* or *ex vivo* include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J.-L. and H. Rétignon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

Expression vectors that may be effective for the expression of TRICH include, but are not limited to, the pCDNA 3.1, EPTAG, pRCMV2, PREP, PVAX, PCR2-TOPOTA vectors (Invitrogen, Carlsbad CA), pCMV-Script, pCMV-TAG, pEGSHIPERV (Stratagene, La Jolla CA), and pTET-ORF, pTET-ON, pTRE2, pTRE2-LUC, pTK-HYG (Clontech, Palo Alto CA). TRICH may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or β -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl.*

Acad. Sci. USA 89:5547-5551; Gossen, M. et al. (1995) Science 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) Curr. Opin. Biotechnol. 9:451-456, commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVORXR and PND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and Blau, H.M. *supra*)), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding TRICH from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID

TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental

parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) Virology 52:456-467), or by electroporation (Neumann, E. et al. (1982) EMBO J. 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to TRICH expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding TRICH under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are

commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) Proc. Natl. Acad. Sci. USA 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) J. Virol. 61:1647-1650; Bender, M.A. et al. (1987) J. Virol. 61:1639-1646; Adam, M.A. and A.D. Miller (1988) J. Virol. 62:3802-3806; Dull, T. et al. (1998) J. Virol. 72:8463-8471; Zufferey, R. et al. (1998) J. Virol. 72:9873-9880). U.S. Patent Number 5,910,434 to Riggs ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant")

discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4⁺ T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) J. Virol. 71:7020-7029; Bauer, G. et al. (1997) Blood 89:2259-2267; Bonyhadi, M.L. (1997) J. Virol. 71:4707-4716; Ranga, U. et al. (1998) Proc. Natl. Acad. Sci. USA 95:1201-1206; Su, L. (1997) Blood 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver

polynucleotides encoding TRICH to cells which have one or more genetic abnormalities with respect to the expression of TRICH. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) Transplantation 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) Annu. Rev. Nutr. 19:511-544 and Verma, I.M. and N. Somia (1997) Nature 389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding TRICH to target cells which have one or more genetic abnormalities with respect to the expression of TRICH. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing TRICH to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) Exp. Eye Res. 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S.

Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant

HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goinis, W.F. et al. (1999) J. Virol. 73:519-532 and Xu, H. et al. (1994) Dev. Biol. 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding TRICH to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) Curr. Opin. Biotechnol. 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity

(e.g., protease and polymerase). Similarly, inserting the coding sequence for TRICH into the alphavirus genome in place of the capsid-coding region results in the production of a large number of TRICH-coding RNAs and the synthesis of high levels of TRICH in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) *Virology* 228:74-83). The wide host range of alphaviruses will allow the introduction of TRICH into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, *Molecular and Immunologic Approaches*, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding TRICH.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques

for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding TRICH. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2'-O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding TRICH. Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased TRICH expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding TRICH may be therapeutically useful, and in the treatment of disorders associated with decreased TRICH expression or activity, a compound which specifically promotes expression of the polynucleotide encoding TRICH may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding TRICH is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an *in vitro* cell-free or reconstituted

biochemical system. Alterations in the expression of a polynucleotide encoding TRICH are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding TRICH. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a *Schizosaccharomyces pombe* gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of TRICH, antibodies to TRICH, and mimetics, agonists, antagonists, or inhibitors of TRICH.

The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising TRICH or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, TRICH or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example TRICH or fragments thereof, antibodies of TRICH, and agonists, antagonists or inhibitors of TRICH, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED₅₀ (the dose therapeutically effective in 50% of the population) or LD₅₀ (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD₅₀/ED₅₀ ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED₅₀ with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

5 The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

10 Normal dosage amounts may vary from about 0.1 μ g to 100,000 μ g, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

15 In another embodiment, antibodies which specifically bind TRICH may be used for the diagnosis of disorders characterized by expression of TRICH, or in assays to monitor patients being treated with TRICH or agonists, antagonists, or inhibitors of TRICH. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics.

20 Diagnostic assays for TRICH include methods which utilize the antibody and a label to detect TRICH in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

25 A variety of protocols for measuring TRICH, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of TRICH expression. Normal or standard values for TRICH expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibodies to TRICH under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of TRICH expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values.

30 Deviation between standard and subject values establishes the parameters for diagnosing disease. In another embodiment of the invention, the polynucleotides encoding TRICH may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of TRICH may be correlated

with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of TRICH, and to monitor regulation of TRICH levels during therapeutic intervention.

5 In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding TRICH or closely related molecules may be used to identify nucleic acid sequences which encode TRICH. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding TRICH, allelic variants, or related sequences.

10 Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the TRICH encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:31-60 or from genomic sequences including promoters, enhancers, and introns of the TRICH gene.

15 Means for producing specific hybridization probes for DNAs encoding TRICH include the cloning of polynucleotide sequences encoding TRICH or TRICH derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as 32 P or 35 S, or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

20 Polynucleotide sequences encoding TRICH may be used for the diagnosis of disorders associated with expression of TRICH. Examples of such disorders include, but are not limited to, a transport disorder such as akinesia, amyotrophic lateral sclerosis, ataxia telangiectasia, cystic fibrosis, Becker's muscular dystrophy, Bell's palsy, Charcot-Marie Tooth disease, diabetes mellitus, diabetes insipidus, diabetic neuropathy, Duchenne muscular dystrophy, hyperkalemic periodic paralysis, normokalemic periodic paralysis, Parkinson's disease, malignant hyperthermia, mulling resistance, myasthenia gravis, myotonic dystrophy, caudonia, tardive dyskinesia, dystonias, peripheral neuropathy, cerebral neoplasms, prostate cancer, cardiac disorders associated with transport, e.g., angina, bradyarrhythmia, tachyarrhythmia, hypertension, Long QT syndrome, myocarditis, cardiomyopathy, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, thyrotoxic myopathy, ethanol myopathy, dermatomyositis, inclusion body myositis, infectious myositis, polymyositis, neurological disorders associated with transport, e.g., Alzheimer's disease, amnesia, bipolar disorder, dementia, depression, epilepsy, Tourette's disorder, paranoid psychoses, and schizophrenia, and other disorders associated with transport, e.g., neurofibromatosis, postherpetic neuralgia, trigeminal neuropathy, sarcoidosis, sickle cell anemia, Wilson's disease,

cataracts, infertility, pulmonary artery stenosis, sensorineural autosomal deafness, hyperglycemia, hypoglycemia, Grave's disease, goiter, Cushing's disease, Addison's disease, glucose-galactose malabsorption syndrome, hypercholesterolemia, adrenoleukodystrophy, Zellweger syndrome, Menkes disease, occipital horn syndrome, von Gierke disease, cystinuria, iminoglycinuria, Hartup disease, and Fanconi disease; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Strausler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalovirginal syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, cataplexy, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; a muscle disorder such as cardiomyopathy, myocardiitis, Duchenne's muscular dystrophy, Becker's muscular dystrophy, myotonic dystrophy, central core disease, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, infectious myositis, polymyositis, dermatomyositis, inclusion body myositis, thyrotoxic myopathy, ethanol myopathy, angina, anaphylactic shock, arrhythmias, asthma, cardiovascular shock, Cushing's syndrome, hypertension, hypoglycemia, myocardial infarction, migraine, pheochromocytoma, and myopathies including encephalopathy, epilepsy, Kearns-Sayre syndrome, lactic acidosis, myoclonic disorder, ophthalmoplegia, and acid maltase deficiency (AMD, also known as Pompe's disease); an immunological disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact

dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hyperesinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. The polynucleotide sequences encoding TRICH may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered TRICH expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding TRICH may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding TRICH may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding TRICH in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of TRICH, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding TRICH, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from

normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding TRICH may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced *in vitro*. Oligomers will preferably contain a fragment of a polynucleotide encoding TRICH, or a fragment of a polynucleotide complementary to the polynucleotide encoding TRICH, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding TRICH may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (FSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding TRICH are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In FSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplifiers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (iSNP), are capable of identifying polymorphisms by comparing the

sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of TRICH include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:225-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described below. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, TRICH fragments of TRICH, or antibodies specific for TRICH may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis" U.S. Patent Number 5,640,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the

hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) *Mol. Carcinog.* 24:153-159; Steiner, S. and N.L. Anderson (2000) *Toxicol. Lett.* 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at

<http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global

pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, 1999a). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for TRICH to quantify the levels of TRICH expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; Mendoza, L.G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schema, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO99/251116; Shalton, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in *DNA Microarrays: A Practical Approach*, M. Schema, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding TRICH may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence.

Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial

chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism

(RFLP). (See, for example, Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

Fluorescent *in situ* hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Urich, et al. (1995) in Meyers, *supra*, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding TRICH on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23,

any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, TRICH, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between TRICH and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with TRICH, or fragments thereof, and washed. Bound TRICH is then detected by methods well known in the art. Purified TRICH can also be coated directly onto plates for use in the aforementioned drug screening techniques.

Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding TRICH specifically compete with a test compound for binding TRICH. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more

antigenic determinants with TRICH.

In additional embodiments, the nucleotide sequences which encode TRICH may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, including U.S. Ser. No. 60/223,269, U.S. Ser. No. 60/224,456, U.S. Ser. No. 60/226,410, U.S. Ser. No. 60/228,140, U.S. Ser. No. 60/230,067, and U.S. Ser. No. 60/231,434, are hereby expressly incorporated by reference.

EXAMPLES

I. Construction of cDNA Libraries

Incyte cDNAs were derived from cDNA libraries described in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA) and shown in Table 4, column 5. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A)⁺ RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)⁺PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, *supra*, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the

appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSFORT1 plasmid (Life Technologies), PCDNA2.1 plasmid (Invitrogen, Carlsbad CA), PBK-CMV plasmid (Stratagene), or pINCY (Incyte Genomics, Palo Alto CA), or derivatives thereof. Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5α, DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by *in vivo* excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows.

Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Applied Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIODYE Terminator cycle sequencing ready reaction kit (Applied Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading

frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, *supra*, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VIII.

The polynucleotide sequences derived from Incyte cDNAs were validated by removing vector, linker, and poly(A) sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The Incyte cDNA sequences or translations thereof were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and hidden Markov model (HMM)-based protein family databases such as PFAM. (HMM is a probabilistic approach which analyzes consensus primary structures of gene families. See, for example, Eddy, S.R. (1996) *Curr. Struct. Biol.* 6:361-365.) The queries were performed using programs based on BLAST, FASTA, BLMPFS, and HMMER. The Incyte cDNA sequences were assembled to produce full length polynucleotide sequences. Alternatively, GenBank cDNAs, GenBank ESTs, stitched sequences, stretched sequences, or Genscan-predicted coding sequences (see Examples IV and V) were used to extend Incyte cDNA assemblies to full length. Assembly was performed using programs based on Phred, Phrap, and Consed, and cDNA assemblies were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length polypeptide sequences. Alternatively, a polypeptide of the invention may begin at any of the methionine residues of the full length translated polypeptide. Full length polypeptide sequences were subsequently analyzed by querying against databases such as the GenBank protein databases (genpept), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and hidden Markov model (HMM)-based protein family databases such as PFAM. Full length polynucleotide sequences are also analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments are generated using default parameters specified by the CLUSTAL algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

Table 7 summarizes the tools, programs, and algorithms used for the analysis and assembly of Incyte cDNA and full length sequences and provides applicable descriptions, references, and threshold parameters. The first column of Table 7 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score or the lower the probability value,

the greater the identity between two sequences).

The programs described above for the assembly and analysis of full length polynucleotide and polypeptide sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:31-60. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies are described in Table 4, column 4.

IV. Identification and Editing of Coding Sequences from Genomic DNA

Putative transporters and ion channels were initially identified by running the Genscan gene identification program against public genomic sequence databases (e.g., *gdbpi* and *gdbhg*). Genscan is a general-purpose gene identification program which analyzes genomic DNA sequences from a variety of organisms (See Bugge, C. and S. Karlin (1997) *J. Mol. Biol.* 268:78-94, and Bugge, C. and S. Karlin (1998) *Curr. Opin. Struct. Biol.* 8:346-354). The program concatenates predicted exons to form an assembled cDNA sequence extending from a methionine to a stop codon. The output of Genscan is a FASTA database of polynucleotide and polypeptide sequences. The maximum range of sequence for Genscan to analyze at once was set to 30 kb. To determine which of these Genscan predicted cDNA sequences encode transporters and ion channels, the encoded polypeptides were analyzed by querying against PFAM models for transporters and ion channels. Potential transporters and ion channels were also identified by homology to Incyte cDNA sequences that had been annotated as transporters and ion channels. These selected Genscan-predicted sequences were then compared by BLAST analysis to the *genpept* and *gbpi* public databases. Where necessary, the Genscan-predicted sequences were then edited by comparison to the top BLAST hit from *genpept* to correct errors in the sequence predicted by Genscan, such as extra or omitted exons. BLAST analysis was also used to find any Incyte cDNA or public cDNA coverage of the Genscan-predicted sequences, thus providing evidence for transcription. When Incyte cDNA coverage was available, this information was used to correct or confirm the Genscan predicted sequence. Full length polynucleotide sequences were obtained by assembling Genscan-predicted coding sequences with Incyte cDNA sequences and/or public cDNA sequences using the assembly process described in Example III. Alternatively, full length polynucleotide sequences were derived entirely from edited or unedited Genscan-predicted coding sequences.

V. Assembly of Genomic Sequence Data with cDNA Sequence Data

"Stitched" Sequences

Partial cDNA sequences were extended with exons predicted by the Genscan gene identification program described in Example IV. Partial cDNAs assembled as described in Example III were mapped to genomic DNA and parsed into clusters containing related cDNAs and Genscan exon predictions from one or more genomic sequences. Each cluster was analyzed using an algorithm based on graph theory and dynamic programming to integrate cDNA and genomic information,

generating possible splice variants that were subsequently confirmed, edited, or extended to create a full length sequence. Sequence intervals in which the entire length of the interval was present on more than one sequence in the cluster were identified, and intervals thus identified were considered to be equivalent by transitivity. For example, if an interval was present on a cDNA and two genomic sequences, then all three intervals were considered to be equivalent. This process allows unrelated but consecutive genomic sequences to be brought together, bridged by cDNA sequence. Intervals thus identified were then "stitched" together by the stitching algorithm in the order that they appear along their parent sequences to generate the longest possible sequence, as well as sequence variants. Linkages between intervals which proceed along one type of parent sequence (cDNA to cDNA or genomic sequence to genomic sequence) were given preference over linkages which change parent type (cDNA to genomic sequence). The resultant stitched sequences were translated and compared by BLAST analysis to the genpept and gbpi public databases. Incorrect exons predicted by Genscan were corrected by comparison to the top BLAST hit from genpept. Sequences were further extended with additional cDNA sequences, or by inspection of genomic DNA, when necessary.

15 "Stretched" Sequences

Partial DNA sequences were extended to full length with an algorithm based on BLAST analysis. First, partial cDNAs assembled as described in Example III were queried against public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases using the BLAST program. The nearest GenBank protein homolog was then compared by BLAST analysis to either Incyte cDNA sequences or GenScan exon predicted sequences described in Example IV. A chimeric protein was generated by using the resultant high-scoring segment pairs (HSPs) to map the translated sequences onto the GenBank protein homolog. Insertions or deletions may occur in the chimeric protein with respect to the original GenBank protein homolog. The GenBank protein homolog, the chimeric protein, or both were used as probes to search for homologous genomic sequences from the public human genome databases. Partial DNA sequences were therefore "stretched" or extended by the addition of homologous genomic sequences. The resultant stretched sequences were examined to determine whether it contained a complete gene.

25 VI. Chromosomal Mapping of TRICH Encoding Polynucleotides

The sequences which were used to assemble SEQ ID NO:31-60 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:31-60 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Piprap (Table 7). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Génethon were used to determine if any of the clustered sequences

had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO., to that map location.

Map locations are represented by ranges, or intervals, of human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Génethon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Human genome maps and other resources available to the public, such as the NCBI "GeneMap 99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

10 VII. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, *supra*, ch. 7; Ausubel (1995) *supra*, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$25 \quad \frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum (length(Seq. 1), length(Seq. 2))}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the

entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

- 5 Alternatively, polynucleotide sequences encoding TRICH are analyzed with respect to the tissue sources from which they were derived. For example, some full length sequences are assembled, at least in part, with overlapping lincyte cDNA sequences (see Example III). Each cDNA sequence is derived from a cDNA library constructed from a human tissue. Each human tissue is classified into one of the following organ/tissue categories: cardiovascular system; connective tissue; digestive system; embryonic structures; endocrine system; exocrine glands; genitalia, female; genitalia, male; germ cells; hemic and immune system; liver; musculoskeletal system; nervous system; pancreas; respiratory system; sense organs; skin; stomatognathic system; unclassified/mixed; or urinary tract. The number of libraries in each category is counted and divided by the total number of libraries across all categories. Similarly, each human tissue is classified into one of the following disease/condition categories: cancer, cell line, developmental, inflammation, neurological, trauma, cardiovascular, pooled, and other, and the number of libraries in each category is counted and divided by the total number of libraries across all categories. The resulting percentages reflect the tissue- and disease-specific expression of cDNA encoding TRICH. cDNA sequences and cDNA library/tissue information are found in the LIFESEQ GOLD database (Lincyte Genomics, Palo Alto CA).

20 VIII. Extension of TRICH Encoding Polynucleotides

Full length polynucleotide sequences were also produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer was synthesized to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and 2-mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer

pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

5 The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (LabSystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose gel to determine which reactions were successful in extending the sequence.

15 The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviII cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

25 The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIODYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

35 In like manner, full length polynucleotide sequences are verified using the above procedure or are used to obtain 5' regulatory sequences using the above procedure along with oligonucleotides designed for such extension, and an appropriate genomic library.

IX. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:31-60 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Cl of [γ - 32 P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech).

10 An aliquot containing 10⁷ counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases:

Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

X. Microarrays

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing. See, e.g., Baldeschweiler, *supra*), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the aforementioned technologies should be uniform and solid with a non-porous surface (Schna (1999), *supra*). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Schna, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645; Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection.

After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

Tissue or Cell Sample Preparation

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)⁺ RNA is purified using the oligo-(dT) cellulose method. Each poly(A)⁺ RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/ μ l oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/ μ l RNase inhibitor, 500 μ M dATP, 500 μ M dGTP, 500 μ M dTTP, 40 μ M dCTP, 40 μ M dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)⁺ RNA with GEMBRIGHT[®] kits (Incyte). Specific control poly(A)⁺ RNAs are synthesized by *in vitro* transcription from non-coding yeast genomic DNA. After incubation at 37°C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85°C to the stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μ l 5X SSC/0.2% SDS.

Microarray Preparation

Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 μ g. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a

110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522, incorporated herein by reference. 1 μ l of the array element DNA, at an average concentration of 100 ng/ μ l, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60°C followed by washes in 0.2% SDS and distilled water as before.

Hybridization

Hybridization reactions contain 9 μ l of sample mixture consisting of 0.2 μ g each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65°C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm² coverslip. The arrays are transferred to a watertight chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 μ l of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60°C. The arrays are washed for 10 min at 45°C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45°C in a second wash buffer (0.1X SSC), and dried.

Detection

Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source,

although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:1,000,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Uncyey).

XI. Complementary Polynucleotides

Sequences complementary to the TRICH-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring TRICH. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of TRICH. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the TRICH-encoding transcript.

XII. Expression of TRICH

Expression and purification of TRICH is achieved using bacterial or virus-based expression systems. For expression of TRICH in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*lac*) hybrid

promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express TRICH upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of TRICH in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding TRICH by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (SF9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, TRICH is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from TRICH at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunofluorescence purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified TRICH obtained by these methods can be used directly in the assays shown in Examples XVI, XVII, and XVIII, where applicable.

XIII. Functional Assays

TRICH function is assessed by expressing the sequences encoding TRICH at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include PCMV SPORT (Life Technologies) and PCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the

recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of TRICH on gene expression can be assessed using highly purified

populations of cells transfected with sequences encoding TRICH and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding TRICH and other genes of interest can be analyzed by northern analysis or microarray techniques.

XIV. Production of TRICH Specific Antibodies

TRICH substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) Methods Enzymol. 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the TRICH amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using FMOC chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (NHS) to increase immunogenicity. (See, e.g., Ausubel, 1995, supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-TRICH activity by, for example, binding the peptide or TRICH to a substrate,

blocking with 1% BSA, reacting with rabbit antiserum, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XV. Purification of Naturally Occurring TRICH Using Specific Antibodies

Naturally occurring or recombinant TRICH is substantially purified by immunoaffinity chromatography using antibodies specific for TRICH. An immunoaffinity column is constructed by covalently coupling anti-TRICH antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing TRICH are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of TRICH (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/TRICH binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and TRICH is collected.

XVI. Identification of Molecules Which Interact with TRICH

Molecules which interact with TRICH may include transporter substrates, agonists or antagonists, modulatory proteins such as G $\beta\gamma$ proteins (Reimann, *supra*) or proteins involved in TRICH localization or clustering such as MAGUKs (Craven, *supra*). TRICH, or biologically active fragments thereof, are labeled with 125 I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) Biochem. J. 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled TRICH, washed, and any wells with labeled TRICH complex are assayed. Data obtained using different concentrations of TRICH are used to calculate values for the number, affinity, and association of TRICH with the candidate molecules.

Alternatively, proteins that interact with TRICH are isolated using the yeast 2-hybrid system (Fields, S. and O. Song (1989) Nature 340:245-246). TRICH, or fragments thereof, are expressed as fusion proteins with the DNA binding domain of Gal4 or LexA, and potential interacting proteins are expressed as fusion proteins with an activation domain. Interactions between the TRICH fusion protein and the TRICH interacting proteins (fusion proteins with an activation domain) reconstitute a transactivation function that is observed by expression of a reporter gene. Yeast 2-hybrid systems are commercially available, and methods for use of the yeast 2-hybrid system with ion channel proteins are discussed in Niebhammer, M. and M. Sheng (1998, *Meth. Enzymol.* 293:104-122).

TRICH may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

Potential TRICH agonists or antagonists may be tested for activation or inhibition of TRICH

ion channel activity using the assays described in section XV.III.

XVII. Demonstration of TRICH Activity

Ion channel activity of TRICH is demonstrated using an electrophysiological assay for ion conductance. TRICH can be expressed by transforming a mammalian cell line such as COS \pm , HeLa or CHO with a eukaryotic expression vector encoding TRICH. Eukaryotic expression vectors are commercially available, and the techniques to introduce them into cells are well known to those skilled in the art. A second plasmid which expresses any one of a number of marker genes, such as β -galactosidase, is co-transformed into the cells to allow rapid identification of those cells which have taken up and expressed the foreign DNA. The cells are incubated for 48-72 hours after transformation under conditions appropriate for the cell line to allow expression and accumulation of TRICH and β -galactosidase.

Transformed cells expressing β -galactosidase are stained blue when a suitable colorimetric substrate is added to the culture media under conditions that are well known in the art. Stained cells are tested for differences in membrane conductance by electrophysiological techniques that are well known in the art. Untransformed cells, and/or cells transformed with either vector sequences alone or β -galactosidase sequences alone, are used as controls and tested in parallel. Cells expressing TRICH will have higher anion or cation conductance relative to control cells. The contribution of TRICH to conductance can be confirmed by incubating the cells using antibodies specific for TRICH. The antibodies will bind to the extracellular side of TRICH, thereby blocking the pore in the ion channel, and the associated conductance.

Alternatively, ion channel activity of TRICH is measured as current flow across a TRICH-containing *Xenopus laevis* oocyte membrane using the two-electrode voltage-clamp technique (Ishii et al., *supra*; Jegla, T. and L. Salter (1997) J. Neurosci. 17:32-44). TRICH is subcloned into an appropriate *Xenopus* oocyte expression vector, such as pBFP, and 0.5-5 ng of mRNA is injected into mature stage IV oocytes. Injected oocytes are incubated at 18°C for 1-5 days. Inside-out macropatches are excised into an intracellular solution containing 116 mM K-gluconate, 4 mM KCl, and 10 mM HEPES (pH 7.2). The intracellular solution is supplemented with varying concentrations of the TRICH mediator, such as cAMP, cGMP, or Ca $^{2+}$ (in the form of CaCl $_2$), where appropriate. Electrode resistance is set at 2-5 M Ω and electrodes are filled with the intracellular solution lacking mediator. Experiments are performed at room temperature from a holding potential of 0 mV. Voltage ramps (2.5 s) from -100 to 100 mV are acquired at a sampling frequency of 500 Hz. Current measured is proportional to the activity of TRICH in the assay.

In particular, the activity of TRICH-20 is measured as Ca $^{2+}$ conductance, the activity of TRICH-22 is measured as Cl $^{-}$ conductance in the presence of glycine, the activity of TRICH-23 is

measured as Ca^{2+} conductance, and the activity of TRICH-24 is measured as K^{+} conductance in the presence of Ca^{2+} , and the activity of TRICH-26 is measured as cation conductance in the presence of heat.

5 Transport activity of TRICH is assayed by measuring uptake of labeled substrates substrates (including but not limited to, maltoose, glucose, or glycogen) into *Xenopus laevis* oocytes. Oocytes at stages V and VI are injected with TRICH mRNA (10 ng per oocyte) and incubated for 3 days at 18°C in OR2 medium (82.5mM NaCl, 2.5 mM KCl, 1mM CaCl_2 , 1mM MgCl_2 , 1mM Na_2HPO_4 , 5 mM Hepes, 3.8 mM NaOH, .50µg/ml gentamycin, pH 7.8) to allow expression of TRICH. Oocytes are then transferred to standard uptake medium (100mM NaCl, 2 mM KCl, 1mM CaCl_2 , 1mM MgCl_2 , 10 mM Hepes/Tris pH 7.5). Uptake of various substrates (e.g., amino acids, sugars, drugs, ions, and neurotransmitters) is initiated by adding labeled substrate (e.g. radiolabeled with ^3H , fluorescently labeled with rhodamine, etc.) to the oocytes. After incubating for 30 minutes, uptake is terminated by washing the oocytes three times in Na^{+} -free medium, measuring the incorporated label, and comparing with controls. TRICH activity is proportional to the level of internalized labeled substrate.

15 In particular, test substrates include sulfate for TRICH-13, tricarboxylates for TRICH-21, dicarboxylates and Na^{+} for TRICH-25, ornithine for TRICH-27, and monocarboxylates for TRICH-28.

ATPase activity associated with TRICH can be measured by hydrolysis of radiolabeled ATP- $[\gamma\text{-}^{32}\text{P}]$, separation of the hydrolysis products by chromatographic methods, and quantitation of the recovered ^{32}P using a scintillation counter. The reaction mixture contains ATP- $[\gamma\text{-}^{32}\text{P}]$ and varying amounts of TRICH in a suitable buffer incubated at 37°C for a suitable period of time. The reaction is terminated by acid precipitation with trichloroacetic acid and then neutralized with base, and an aliquot of the reaction mixture is subjected to membrane or filter paper-based chromatography to separate the reaction products. The amount of ^{32}P liberated is counted in a scintillation counter. The amount of radioactivity recovered is proportional to the ATPase activity of TRICH in the assay.

XVIII. Identification of TRICH Agonists and Antagonists

TRICH is expressed in a eukaryotic cell line such as CHO (Chinese Hamster Ovary) or HEK (Human Embryonic Kidney) 293. Ion channel activity of the transformed cells is measured in the presence and absence of candidate agonists or antagonists. Ion channel activity is assayed using patch clamp methods well known in the art or as described in Example XVII. Alternatively, ion channel activity is assayed using fluorescent techniques that measure ion flux across the cell membrane (Veleceblei, G. et al. (1999) Meth. Enzymol. 294:20-47; West, M.R. and C.R. Molloy (1996) Anal. Biochem. 241:51-58). These assays may be adapted for high-throughput screening using microplates. Changes in internal ion concentration are measured using fluorescent dyes such as the Ca^{2+} indicator Fluo-4 AM, sodium-sensitive dyes such as SBFI and sodium green, or the Cl^{-}

indicator MQAE (all available from Molecular Probes) in combination with the FLIPR fluorimetric plate reading system (Molecular Devices). In a more generic version of this assay, changes in membrane potential caused by ionic flux across the plasma membrane are measured using oxonyl dyes such as DiBAC₄ (Molecular Probes). DiBAC₄ equilibrates between the extracellular solution and cellular sites according to the cellular membrane potential. The dye's fluorescence intensity is 20-fold greater when bound to hydrophobic intracellular sites, allowing detection of DiBAC₄ entry into the cell (Gonzalez, J.E. and P.A. Negulescu (1998) Curr. Opin. Biotechnol. 9:624-631).

Candidate agonists or antagonists may be selected from known ion channel agonists or antagonists, peptide libraries, or combinatorial chemical libraries.

10 Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID
2194064	1	2194064CD1	31	2194064CB1
2744094	2	2744094CD1	32	2744094CB1
2798241	3	2798241CD1	33	2798241CB1
3105257	4	3105257CD1	34	3105257CB1
3200979	5	3200979CD1	35	3200979CB1
6754139	6	6754139CD1	36	6754139CB1
6996659	7	6996659CD1	37	6996659CB1
7472747	8	7472747CD1	38	7472747CB1
7474121	9	7474121CD1	39	7474121CB1
7475615	10	7475615CD1	40	7475615CB1
7475656	11	7475656CD1	41	7475656CB1
7480632	12	7480632CD1	42	7480632CB1
6952742	13	6952742CD1	43	6952742CB1
7478795	14	7478795CD1	44	7478795CB1
656293	15	656293CD1	45	656293CB1
7473957	16	7473957CD1	46	7473957CB1
7474111	17	7474111CD1	47	7474111CB1
7480826	18	7480826CD1	48	7480826CB1
6025572	19	6025572CD1	49	6025572CB1
5686561	20	5686561CD1	50	5686561CB1
1553725	21	1553725CD1	51	1553725CB1
1695770	22	1695770CD1	52	1695770CB1
4672222	23	4672222CD1	53	4672222CB1
6176128	24	6176128CD1	54	6176128CB1
7473418	25	7473418CD1	55	7473418CB1
7474129	26	7474129CD1	56	7474129CB1
7481414	27	7481414CD1	57	7481414CB1
7481461	28	7481461CD1	58	7481461CB1
7472541	29	7472541CD1	59	7472541CB1
6999183	30	6999183CD1	60	6999183CB1

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
1	2194064CD1	g2463634	1.60E-41	Monocarboxylate transporter [Homo sapiens] (Price, N. T. et al. (1998) Biochem. J. 329:321-328)
2	2744094CD1	g13346481	0	ATP-binding cassette transporter MRP8 [Homo sapiens]
3	2798241CD1	g1699038	2.90E-142	ABC3 [Homo sapiens] (Connors, T. D. et al. (1997) Genomics 39:231-234)
4	3105257CD1	g8650412	0	M-ABC2 protein [Homo sapiens] (Zhang, F. et al. (2000) Characterization of ABCB9, an ATP binding cassette protein associated with lysosomes J. Biol. Chem. 275:23287-23294)
5	3200979CD1	g1514530	3.10E-119	ABC-C transporter [Homo sapiens] (Klugbauer, N. and P. Hofmann (1996) FEBS Lett. 391:61-65)
6	6754139CD1	g6746563	1.70E-188	neuronal nicotinic acetylcholine receptor subunit [Rattus norvegicus] (Elgoyhen, A. B. et al. (2001) alpha 10: A determinant of nicotinic cholinergic receptor function in mammalian vestibular and cochlear mechanosensory hair cells Proc. Natl. Acad. Sci. U.S.A. 98:3501-3506)
7	6996659CD1	g1050330	0	Ionotropic glutamate receptor [Rattus norvegicus] (Ciabarra, A.M. et al. (1995) J. Neurosci. 15:6498- 6508)
8	7472747CD1	g13926108	1.00E-157	2P domain potassium channel Talk-1 [Homo sapiens] (Girard, C. et al. (2001) Genomic and functional characteristics of novel human pancreatic 2P domain K(+) channels. Biochem Biophys Res Commun. 282:249-256)
9	7474121CD1	g2465542	7.00E-20	TWIK-related acid-sensitive K+ channel [Homo sapiens] (Duprat, F. et al. (1997) EMBO J. 16:5464-5471)
10	7475615CD1	g2654005	5.70E-114	Pendrin [Homo sapiens] (Everett, L.A. et al. (1997) Nature Genet. 17:411-422)

Table 2 (cont.)

Polypeptide Incyte	Genbank ID	Genbank ID	Probability	Genbank Homolog
SEQ ID NO: Polypeptide ID	NO: ID	score		
20	5686561CD1	g4586963	2.40E-27	voltage-gated Ca channel [Rattus norvegicus] (Tashbashi, K. et al. (2000) Molecular cloning of a novel form (two-repeat) protein related to voltage-gated sodium and calcium channels. Biochem. Biophys. Res. Commun. 270:370-376)
21	1553725CD1	g545998	1.60E-89	tricarboxylate carrier [Rattus sp.] (Azzi, A. et al. (1993) The mitochondrial tricarboxylate carrier. J. Bioenerg. Biomembr. 25:515-524)
22	1695770CD1	g31849	1.10E-175	Inhibitory glycine receptor [Homo sapiens] (Greeningloh, G. et al. (1990) Alpha subunit variants of the human glycine receptor: primary structures, functional expression and chromosomal localization of the corresponding genes. EMBO J. 9:771-776)
23	4672222CD1	g13562153	0	channel-kinase 1 [Homo sapiens] (Ryazanov, A. G. et al. (1999) Alpha-kinases: a new class of protein kinases with a novel catalytic domain. Curr. Biol. 9:R43-R45)
24	6176128CD1	g3978472	0	potassium channel subunit [Rattus norvegicus] (Jolivet, W.J. et al. (1998) Formation of intermediate-conductance calcium-activated potassium channels by interaction of Slack and Slo subunits. Nat. Neurosci. 1:462-469)
25	7473418CD1	g2811122	2.90E-177	NADC-2 [Xenopus laevis] (Nat. Neurosci. 1:462-469)
26	7474129CD1	g2570933	1.20E-134	vanilloid receptor subtype 1 [Rattus norvegicus] (Caterina, M.J. et al. (1997) The capsaicin receptor: a heat-activated ion channel in the pain pathway. Nature 389:816-824)

Table 2 (cont.)

11	7475656CD1	g3168874	0	Ion channel BCNG-1 [Homo sapiens] (Santero, B. et al. (1997) Proc. Natl. Acad. Sci. USA 94:14815-14820)
12	7480632CD1	g1514530	9.80E-123	ABC-C transporter [Homo sapiens] (Kugbauer, N. and F. Hofmann (1996) FEBS Lett. 391:61-65)
13	6952742CD1	g10719650	0	sulfate/anion transporter SAT-1 protein [Homo sapiens] (Loht, H. et al. (2000) Mapping of Five New Putative Anion Transporter Genes in Human and Characterization of SLCT26A6, A Candidate Gene for Pancreatic Anion Exchanger. Genomics 70:102-112)
14	7478795CD1	g6045150	0	TAP-like ABC transporter [Rattus norvegicus] (Yamaguchi, Y. et al. (1999) An ABC transporter homologous to TAP proteins. FEBS Lett. 457:231-236)
15	6562933CD1	g6746563	1.30E-220	neuronal nicotinic acetylcholine receptor [Rattus norvegicus] (Blachly-Dyson, B. et al. (1993) J. Biol. Chem. 268:1835-1841)
16	7473957CD1	g340199	1.20E-130	voltage-dependent anion channel [Homo sapiens] (Cardiac potassium channel subunit (Kv6.2) [Homo sapiens] (Zhu, X. et al. (1999) Receptors Channels 6:337-350)
17	7474111CD1	g6006493	1.50E-75	Cardiac potassium channel subunit (Kv6.2) [Homo sapiens] (Zhu, X. et al. (1999) Receptors Channels 6:337-350)
18	7480826CD1	g8248427	1.50E-235	anion acid transporter system A [Rattus norvegicus] (Sugawara, M. et al. (2000) J. Biol. Chem. 275:16473-16477)
19	6025572CD1	g402628	4.20E-114	adenine nucleotide carrier [Mus musculus] (Sugawara, M. et al. (2000) J. Biol. Chem. 275:16473-16477)

Table 2 (cont.)

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
27	7481414CD1	g13445630	1.00E-151	mutant ornithine transporter 2 [Mus musculus] (Wu, Q. and Maniatis, T. (1999) A striking organization of a large family of human neural cadherin-like cell adhesion genes. Cell 97:779-790)
28	7481461CD1	g458247	1.40E-136	X-linked PEST-containing transporter [Homo sapiens] (Lafreniere, R.G. et al. (1994) A novel transmembrane transporter encoded by the XPCT gene in Xq13.2. Mol. Genet. 3:1133-1139)
29	7472541CD1	g6457270	0	Putative E1-E2 ATPase [Mus musculus] (Halleck, M.S. et al. (1999) Differential expression of putative transbilayer amphipath transporters. Physiol. Genomics (Online) 1:139-150)
30	6999183CD1	g1514530	2.30E-127	ABC-C transporter [Homo sapiens] (Klugbauer N. and Hofmann F. (1996) Primary structure of a novel ABC transporter with a chromosomal localization on the band encoding the multidrug resistance- associated protein, FEBS Lett. 391:61-65)

91

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
1	2194064CD1	308	S287 S51 T132		Signal peptide: M1-A17 Transmembrane domains: W197-V224, Y248-G270 PEST transporter: DM05037 P53988 1-465:M1-L109, L126-K289 DM05037 Q03064 1-475:M1-L109, V110-K289 DM05037 P36021 155-612:G3-G288	SPScan HMMER BLAST-DOMO
2	2744094CD1	606	S116 S133 S266 S299 S403 S503 S604 S63 T112 T253 T318 T330 T388 T455 T543 T70	N216 N386 N62 N68	Transmembrane domains: P25-W49, Q82-I107, L166-L187, P184-M203 ABC transporter: H392-G575 ABC transporter transmembrane region: S30-A319 ABC transporters family signature: A483-D533 ABC transporter: F502-V516 ATP/GTP binding site: G399-S406 ATP-binding transporter: PD00131:G141-D150, S403-I456, G550-R587 ABC transporters family: DM00008 P33527 1293-1502: F367-G575 DM00008 Q10185 1239-1448: I365-G575 DM00008 P39109 1272-1482: I365-G575 DM00008 S64757 1302-1528: I365-K486 ATP-binding transport protein: PD000130: T61-G292 PD002040: G434-P488	HMMER HMMER-PFAM HMMER-PFAM ProfileScan MOTIFS MOTIFS BLIMPS-PRODOM BLAST-DOMO BLAST-PRODOM

92

Table 3 (cont.)

Seq ID	Incyte Polypeptide Acid	Amino Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Potential Signature Sequences, Domains and Motifs	Analytical Methods and Databases
4	3105257CD1	659	S206 S26 S300 S452 S504 S583 S62 T261 T284 T293 T348 T520 T615 Y121 Y298	N131 N210	ABC transporter: G441-G628 ABC transporter transmembrane region: L92-L366 ABC transporters family signature: A535-D586 ABC transporter: L555-L569 ATP/GTP binding site: G448-S455 ABC transporters family: BL00211: L446-V457, L555-D586 ATP-binding transporter: PD00131: G190-D199, S452-L505, G603-L640 ABC transporters family: DM00008 P34712 1076-L290: F415-G628 ATP-binding transporter protein: PD000130: L135-Y358 Multidrug resistance ATP-binding transport protein: PD167072: W486-G552	BLAST-PRODOM BLAST-PRODOM BLAST-PRODOM BLIMPS-PRODOM BLIMPS-BLOCKS MOTIFS MOTIFS ProfileScan HMMEP-PFAM HMMEP-PFAM Databases

Table 3 (cont.)

Seq ID	Incyte Polypeptide Acid	Amino Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Potential Signature Sequences, Domains and Motifs	Analytical Methods and Databases
3	2798241CD1	1642	S199 S32 T1366 N190 N388 N458 N499 N458 N499 N576 N86 N943 N973 N943 N973 S624 S761 T1454 S815 S859 T1461 S861 S885 S1558 S962 T28 T1635 T486 T518 T1099 T572 T606 T1126 T779 T780 S1190 T854 Y168 S1236 S1247 S1308 S1372 T1429 Y1552	N1627	Transmembrane domains: Q34-M52, S272-P292, S295-P313, V327- I346, I401-L427, V865-H883, P1075-V1098, L1095-P1114, W1137-I1162, I1165-I1184 ABC transporter: G507-G689, G1326-G1509 ABC transporters family signature: V595D646, I1413-D1464 ABC transporter: L615-V629 ATP/GTP binding sites: G514-S521, G1333-S1340 ABC transporters family: DM00008 P41233 839-1045: I478-S687, K1313-M1506 ABC transporters family: DM00008 P34358 611-816: I478-S687, I1319-M1506 DM00008 P26050 8-212: K1313-S1508, I478-I686 DM00008 P41233 1851-2058: R1309-S1508, I478-I686	HMMEP HMMEP-PFAM ProfileScan MOTIFS MOTIFS BLAST-PRODOM BLAST-PRODOM BLAST-PRODOM Databases

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
5	3200979CD1	1592	S125 S187 T1117 S207 S386 T1135 S453 Y906 T1214 S714 S733 T1346 S745 S770 T1388 S778 S874 T1417 S882 S994 S1454 T368 T439 T1494 T484 T542 T1580 T565 T673 S1116 T691 T706 S1206 T766 T1257 T782 T801 T1264 Y927 T98 T1265 Y1192 S7 S1297 S1320 T77 S1328 T1434 T1466	N185 N62 N75 N870 N871 N899 N949 N1164 N1273	Transmembrane domains: I265-V285, L296-I315, M319-L340, I390-F410, L815-M834, L1063-M1082, W1099-T1117, L1126-L1145 ABC transporter: G500-G642, G1281-G1465 ABC transporters family signature: L1372-D1420 ATP/GTP binding sites: G507-S514, G1288-S1295 ABC transporters family: BL00211: I505-L516, L1389-D1420 ABC transporters family: DM00008 P41233 839-1045:K1268-M1462, I471-P600, E587-N641 DM00008 P34358 611-816:F1262-M1462, I471-D592, E585-N641 DM00008 P41233 1851-2058:K1266-S1464, I471-V584, V588-N641 DM00008 P23703 41-246:K1268-G1465, V476-L609, E585-G642	HMMER HMMER-PFAM ProfileScan MOTIFS BLIMPS-BLOCKS BLAST-DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
6	6754139CD1	382	S124 S260 S340 S85 T337		Transmembrane domains: A168-H191, V200-L217, Y233-N253, F361-L378 Neurotransmitter-gated ion channel: D2-L378 Neurotransmitter-gated ion-channels signature: V66-G120 Neurotransmitter-gated ion channel: C86-C100 Neurotransmitter-gated ion channel: BL00236:M1-D26, Y155-S196, V43-N52, D71-H109 Neurotransmitter-gated ion channel: PR00252:T9-W25, L42-K53, C86-C100, L162-N174 Nicotinic acetylcholine channel: PR00254:M1-L12, Y30-W44, I48-G60, V66- S84 Neurotransmitter-gated ion channel: DM00195 P43144 5-478:M1-E296, R323-A381 DM00195 JH0173 14-503:M1-P314, L327-A381 DM00195 P09478 5-538:R4-L297, E296-A381 DM00195 P54131 3-491:M1-A312, L327-A381 Postsynaptic ion channel: PD000153: M1-R262, S298-V377	HMMER HMMER-PFAM ProfileScan MOTIFS BLIMPS-BLOCKS BLIMPS-PRINTS BLIMPS-PRINTS BLAST-DOMO BLAST-PRODOM

Table 3 (cont.)

SEQ ID NO	Incyste Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Potential Ion Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases	HMME	HMME-PRAM	BLIMPS-BLOCKS	BLAST-DOMO	BLAST-PRODOM
10	7475615CD1	769	S200 S3 S407	N195 N198	N195 N198	Transmembrane domains: P245-I265, N294-V311, P491-V510	HMME		Sulfate transporter family: L229-T513	Sulfate transporters profile: BL01130: G119-V172, T217-L268	BLAST-DOMO	BLAST-PRODOM
									Sulfate transporter protein: PD001755: H641-R720, L521-D579			

Table 3 (cont.)

SEQ ID NO	Incyste Polypeptide	Amino Acid	Potential Phosphorylation Sites	Potential Glycosylation Sites	Potential Ion Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases	HMME	HMME-PRAM	HMME	HMME-PRAM	HMME	HMME-PRAM	HMME
7	6996659CD1	Residues	S110 S202 S1030	N145 N264	N145 N264	Signal peptide: M1-V24	HMME		Signal peptide: M1-S33	Transmembrane domains: M677-T693, P931-I946	HMME		HMME	
8	7472747CD1	Residues	S193 S199 S91	N57 N86	N57 N86	Glutamate receptor: DM00247 P35436 615-886: T731-Q993	BLAST-DOMO		Ionotropic glutamate receptor: PD156309: S170-Y577	BLAST-PRODOM				
9	7474121CD1	Residues	S205 S252 S267	N70 N96	N70 N96	NMDA receptor signature: PR00177: M677-G702, P744-E771, P931-V955, F593-T621	BLAST-DOMO		Transmembrane domains: F95-L114, V167-P187	HMME		HMME		
						Transmembrane domains: G23-A43, P103-I122, L132-D150, P337-Q359	HMME							

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
11	7475656CD1	882	S102 S108 S13 S324 S360 S394 S395 S518 S544 S591 T190 T242 T649 T754 T799 T869 Y240 Y529	N330 N640 N770 N8	Transmembrane domains: L139-F159, T242-L258, I366-L392 Transmembrane region cyclic nucleotide domain: Y209-I453 Cyclic nucleotide-binding domain: K482-M570 Cyclic nucleotide-binding domain: I494-I515 Cyclic nucleotide-binding site: BL00888: G491-V514, G527-L536 Cyclic nucleotide-binding domain: DM01165 A55251 333-706: H302-E576 DM01165 P29973 311-684: H302-E576 DM01165 Q03041 286-658: H302-E576 DM01165 S52072 262-635: H302-R572 Cyclic nucleotide gated hyperpolarization activated cation channel: PD079330: P747-L882 PD089437: A627-M722 PD108745: M1-D62 PD151315: T577-Q626	HMMER HMMER-PFAM HMMER-PFAM MOTIFS BLIMPS-BLOCKS BLAST-DOMO BLAST-PRODOM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
12	7480632CD1	1547	S134 S196 S1102 S216 S395 T1301 S7 T1343 T1389 S723 S742 T1372 S754 S779 S1405 S787 S883 T1449 S891 T107 T1535 T377 T448 S1158 T493 T551 T1212 T574 T682 S1218 T700 T715 T1219 T775 T791 S1252 T810 T86 S1275 T936 T975 S1283 Y915 S462 T1421 Y1144	N194 N71 N84 N879 N880 N908 N958 N1100 N1228	Transmembrane domains: I274-V294, L305-I324, M328-L349, I399-F419, L824-M843, M946-I963, L1021-F1040, L1046-L1064, D1105-F1123 ABC transporter: G509-G651, G1236-G1420 ABC transporters family signature: L1327-D1375 ATP/GTP binding sites: G516-S523, G1243-S1250 ABC transporters family: BL00211: I514-L525, L1344-D1375 ABC transporters family: DM00008 P41233 839-1045: K1223-M1417, I480-P609, E596-N650 DM00008 P41233 1851-2058: R1220-S1419, I480-V593, V597-N650 DM00008 P34358 611-816: F1217-M1417, I480-D601, E594-N650 DM00008 P23703 41-246: K1223-G1420, V485-L618, E594-G651	HMMER HMMER-PFAM ProfileScan MOTIFS BLIMPS-BLOCKS BLAST-DOMO

Table 3 (cont.)

Protein ID	14	7478795CD1	Residues	766	S161 S275 S28	N280 N508	761	ATP-BINDING TRANSPORT PROTEIN	BLAST-PRODOM	ATP-BINDING TRANSPORT TRANSMEMBRANE	BLIMPS-PRODOM	REGION	PD00131:G283-D292, S543-I596, K691-L728	Transmembrane domain	V85-F104, V185-F204, L328-G347, Y411-G431	ABC transporter transmembrane region.	ABC membrane: L188-M459	ABC transporter	ABC trans: G532-G716	ABC transporter: L643-L657	ATP/GTP-binding site motif A (P-loop)	ATP_GTP_A: G539-S546	ABC transporters family signature	atp_bind_transport.prf: I625-D674	PROFILESSCAN
------------	----	------------	----------	-----	---------------	-----------	-----	-------------------------------	--------------	-------------------------------------	---------------	--------	---	----------------------	---	---------------------------------------	-------------------------	-----------------	----------------------	----------------------------	---------------------------------------	----------------------	-----------------------------------	-----------------------------------	--------------

Table 3 (cont.)

[illegible]

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
15	656293CD1	450	S153 S192 S328 S408 T405	N40 N56	NEUROTRANSMITTER-GATED ION-CHANNELS DM00195 P43144 5-478:A25-E364, R391-A449 CHANNEL IONIC TRANSMEMBRANE GLYCOPROTEIN POSTSYNAPTIC MEMBRANE RECEPTOR PRECURSOR SIGNAL PROTEIN PD000153:S131-R361 Neurotransmitter-gated ion channel BL00236:D139-H177, Y223-S264, V57-D94, V111-N120 NEUROTRANSMITTER-GATED Ion Channel PR00252:T77-W93, L110-K121, C154-C168, L230-N242 NICOTINIC ACETYLCHOLINE RECEPTOR SIGNATURE PR00254:V134-S152, S64-L80, Y98-W112, I116-G128 signal peptide: M1-G24 transmembrane domain: A236-H259, V268-L285, Y301-N321, F429-L446 Neurotransmitter-gated ion-channel neur_chan:A30-L446 Neurotr_Ion_Channel C154-C168 Neurotransmitter-gated ion-channels signature neurotr_ion_channel.prf:V134-G188 signal_cleavage: M1-G24	BLAST_DOMO BLAST_PRODOM BLIMPS_BLOCKS BLIMPS_PRINTS BLIMPS_PRINTS HMMER HMMER HMMER_PFAM MOTIFS PROFILESCAN SPSCAN

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
16	7473957CD1	260	S114 S12 S211 T136 T227 T28 T47 T49 T63 T84	N215 N216	EUKARYOTIC MITOCHONDRIAL PORIN DM01893 P45879 1-282:S12-A260 PORIN CHANNEL VOLTAGEDEPENDENT OUTER MEMBRANE PROTEIN MITOCHONDRION ANIONSELECTIVE MITOCHONDRIAL VDAC PD003211:A15-Q259 Eukaryotic mitochondrial porin BL00558:G33-L46, T57-S81 EUKARYOTIC PORIN SIGNATURE PR00185:G45-T60, E124-E135, Y224-D241 Eukaryotic porin Euk_porin:A5-A260 Eukaryotic_Porin Y202-Y224 Eukaryotic mitochondrial porin signature eukaryotic_porin.prf:M16-S81	BLAST_DOMO BLAST_PRODOM BLIMPS_BLOCKS BLIMPS_PRINTS HMMER_PFAM MOTIFS PROFILESCAN
17	7474111CD1	506	S187 S194 S2 S231 S286 S423 S493 S57 T241 T273 T357 T385	N284	do CHANNEL; POTASSIUM; CDRK; FORM; DM00436 JH0595 144-307:P230-I366 CHANNEL IONIC PROTEIN POTASSIUM SUBUNIT VOLTAGEGATED TRANSMEMBRANE CALCIUM TRANSPORT ION PD000141:F319-Y486 POTASSIUM CHANNEL SIGNATURE PR00169:F319-V339, M363-C389, E392-E415, F427-M449, G456-F482, E211-P230, P245-T273, I293-K316 transmembrane domain: I253-C270, V356-A373, V394-L413 Ion transport protein ion_trans:I263-I478	BLAST_DOMO BLAST_PRODOM BLIMPS_PRINTS HMMER HMMER_PFAM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
20	5686561CD1	540	S162 S180 S24 S29 S327 S349 S454 T527	N399 N406	Transmembrane domains: A77-Y100, Y220-L243, I259-L285, V291-Y311, A369-F389 Sodium channel signature: PR00170:G362-F389, Y76-G105, L361-F389, K109-G134 Calcium channel: DM00043 A55645 1137-1259: A250-V298 (P-value = 2.7e-5) Voltage gated calcium channel PD000032:Y221-G391, I460-F486, N423-W443 (P-value = 1.1e-6)	HMMER BLIMPS-PRINTS BLAST-DOMO BLAST-PRODOM
21	1553725CD1	322	S142 S217 S295 S39 T133 T168 T304 T62 Y315	N123 N131 N29	PROTEIN TRANSMEMBRANE CHROMOSOME PUTATIVE TRANSPORTER C17G6.15C TRANSPORT XV READING FRAME PD006986:F8-L253	BLAST_PRODOM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
22	1695770CD1	417	S108 S122 S163 S43 S56 T196 T239 T243 T410 T411 T88	N72	Signal peptide: M1-A28 Transmembrane domains: M255-I279, I320-I339 Neurotransmitter-gated ion-channel domain: P44-F341 Neurotransmitter-gated ion channels signature BL00236: V73-R110, I127-N136, N157-Y195, F242-A283 Neurotransmitter-gated ion-channels signature: L152-E206 Neurotransmitter-gated ion-channel family signature PR00252:R93-Y109, S126-E137, C172-C186, F249-Q261 Gamma-aminobutyric acid A (GABAA) receptor signature PR00253:Y258-W278, A284-S305, I318-I339 CHANNEL IONIC TRANSMEMBRANE GLYCOPROTEIN POSTSYNAPTIC MEMBRANE RECEPTOR PRECURSOR SIGNAL PROTEIN PD000153: R99-K347 NEUROTRANSMITTER-GATED ION-CHANNELS DM00560 S18836 18-453: R24-D417 Neurotransmitter-gated ion channel motif: C172-C186	HMMER HMMER HMMER_PFAM BLIMPS_BLOCKS PROFILES SCAN BLIMPS_PRINTS BLIMPS_PRINTS BLAST_PRODOM BLAST_DOMO MOTIFS

Seq ID	Incyte	Amino	Potential	Potential	Signature Sequences,	Analytical
ID NO:	Polypeptide	Residues	Phosphorylation	Glycosylation	Domains and Motifs	Methods and Databases
24	6176128CD1	1237			Transmembrane domains: M155-Y177, M248-P264, L310-L330 CHANNEL POTASSIUM IONIC CALCIUMACTIVATED ALPHA CALCIUM SUBUNIT ACTIVATED PROTEIN P983, PD003090:R337-P629, I784-M889, L926- Y1003-E1033, Q1176-S1215 DO CHANNEL, POTASSIUM, MSLO, ACTIVATED; P944-P983, Q1176-S1226	BLAST_PRODOM HMMER
25	7473418CD1	539	S299 S321 T535 S1221 S1125 S1215 T1190 T1231 T1146 T1155 S1013 S1030 S1098 S1219 T777 T994 S1090 T517 T609 T718 T351 T391 T446 S954 T156 T302 S9 S924 S93 S770 S808 S871 S713 S720 S726 S538 S686 S690 S407 S438 S439 S168 S179 S361 S102 S135 S339	N533	Transmembrane domains: V15-C38, C50-P67, F264-F282, A323-R341 Sodium:sulfate symporter signature: BL01271:S451-I505, T132-I151, M216-V240, P378-G399 PROTEIN TRANSMEMBRANE TRANSPORT MEMBRANE INNER TRANSPORTER SODIUM SYMPORT OF COTRANSPORTER PD000549:V15-V173, M216-W518 DO KENAL, BOUND, PRO-SER-ALA, NA; P199-W349, L367-T517	BLAST_PRODOM HMMER
25	7473418CD1	539	S299 S321 T535 S1221 S1125 S1215 T1190 T1231 T1146 T1155 S1013 S1030 S1098 S1219 T777 T994 S1090 T517 T609 T718 T351 T391 T446 S954 T156 T302 S9 S924 S93 S770 S808 S871 S713 S720 S726 S538 S686 S690 S407 S438 S439 S168 S179 S361 S102 S135 S339	N533	Transmembrane domains: V15-C38, C50-P67, F264-F282, A323-R341 Sodium:sulfate symporter signature: BL01271:S451-I505, T132-I151, M216-V240, P378-G399 PROTEIN TRANSMEMBRANE TRANSPORT MEMBRANE INNER TRANSPORTER SODIUM SYMPORT OF COTRANSPORTER PD000549:V15-V173, M216-W518 DO KENAL, BOUND, PRO-SER-ALA, NA; P199-W349, L367-T517	BLAST_PRODOM HMMER
24	6176128CD1	1237			Transmembrane domains: M155-Y177, M248-P264, L310-L330 CHANNEL POTASSIUM IONIC CALCIUMACTIVATED ALPHA CALCIUM SUBUNIT ACTIVATED PROTEIN LARGE PD003090:R337-P629, I784-M889, L926- Y1003-E1033, Q1176-S1215 DO CHANNEL, POTASSIUM, MSLO, ACTIVATED; P944-P983, Q1176-S1226	BLAST_PRODOM HMMER

Table 3 (cont.)

Seq	Incye	Amino	Potential	Potential	Signature Sequences,	Analytical	
ID	Polypeptide	Residues	Phosphorylation	Glycosylation	Domains and Motifs	Methods and	
ID	ID	1864	S103 S195 S196 S2 S22 S406 S5 S547 S697 S727 S757 S836 S87 S883 T115 T12 T299 T318 T349 T367 T508 T523 T529 T593 T603 T615 T675 T778 T795 T842 Y327 S1476 S1503 S1163 S1191 S1361 S1413 S1430 S1493 S1526 S1555 S1614 T1631 S1613 T1742 T1758 S1850 T1245 S1410 S1456 T1471 S1499 S1698 S1859 Y1220 Y1552	N404 N550 N715 N718 N805 N925 N1058 N1465 N1466 N1595 N1595 N1773 N1849	PROTEIN MEMBRANE CHROMOSOME TRANSMEMBRANE COSC12.3 TOH8.5 I P54DL.5 TV PD018035:X108-L439 PD039592:E597-N801 PD151509:V974-P1063, W1030-K1253 PD022180:W434-R545	BLAST_PRODOM	HMMER Databases
23	4672222CD1				Transmembrane domains: F858-M878, N999-L1022, V1079-Q1102		

Table 3 (cont.)

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
26	7474129CD1	755	S339 S353 S367 S463 S53 S572 S589 S653 S732 T128 T132 T255 T270 T277 T300 T343 T358 T362 T37 T376 T441 T664 Y225 Y347 Y587	N417 N648 N735	Transmembrane domains: V490-F507, L556-L573, P616-M642 Ank repeat: E179-K211, F226-S259, D305-K333 VANILLOID RECEPTOR SUBTYPE 1 PD101189: Q52-L291 PROTEIN OLFACTORY CHANNEL B0212.5 T09A12.3 T10B10.7 VANILLOID RECEPTOR SUBTYPE F28H7.10 PD011151:N303-E430	HMMER HMMER_PFAM BLAST_PRODOM BLAST_PRODOM
27	7481414CD1	301	S143 S203 S290 T136 T32		Transmembrane domain: L212-V230 Mitochondrial carrier proteins domain: Q8-M294 Mitochondrial energy transfer proteins signature: BL00215:L214-Q238, V256-G268 Mitochondrial energy transfer proteins signature: A10-G59, L107-I160, K204-A276, K213-N259 PROTEIN TRANSPORT TRANSMEMBRANE REPEAT MITOCHONDRIUM CARRIER MEMBRANE INNER MITOCHONDRIAL ADP/ATP PD000117: Y44-S241 Mitochondrial carrier protein motifs: P126-L134 P229-I237	HMMER HMMER_PFAM BLIMPS_BLOCKS PROFILES CAN BLAST_PRODOM MOTIFS

111

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
28	7481461CD1	515	S10 S104 S163 S257 S272 S277 S4 S474 S511 S97 T233 T250 T484	N81	Transmembrane domains: V117-F135, Y169-L191, I190-I215, G229-P245, I376-F395 Monocarboxylate transporter domain: A77-A455 XLINKED PESTCONTAINING TRANSPORTER SOLUTE CARRIER FAMILY MONOCARBOXYLIC ACID TRANSPORTERS MEMBER PD030892:P33-V111 do PEST; TRANSPORTER; LINKED; DM05037 P36021 155-612:E63-M489	HMMER HMMER_PFAM BLAST_PRODOM BLAST_DOMO

112

Seq ID	Incyte	Polypeptide	Amino Acid	Phosphorylation	Potential	Signature Sequences, Domains and Motifs	Methods and Databases
30	6999183CD1	1585	Residues	Phosphorylation	Potential	ABC TRANSPORTERS FAMILY: DM000008 P41233 839-1045: I1268-M1455, I482-P611, E598-N652 ABC transporters family: BL00211: L516-L527, L1382-D1413 Transmembrane domain (transmem_domain): L1167-M1193, T306-I326, P329-L346, W271-I289, T306-I326, P329-L346, L1167-M1193, T306-I326, P329-L346, P358-M375, Y398-Y420, V1034-F1053 ABC transporter (ABC_tran): G511-G653, G1280-G1458 ATP/GTP-binding site motif A (P-loop) (Acp_gcp_A): G518-T525, G1287-S1294 ABC transporters family signature (acp_bind_transport.prf): I1362-D1413 PROFILESSCAN	ANALYTICAL METHODS AND DATABASES

Table 3 (cont.)

Seq ID	Incyte	Polypeptide	Amino Acid	Phosphorylation	Potential	Signature Sequences, Domains and Motifs	Methods and Databases
29	7472541CD1	1519	Residues	Phosphorylation	Potential	Transmembrane domains: M13-G331, L358-L383, L1317-C1337 HMMEP-PFAM E1-E2 ATPase domains: E422-V444, L935-H985 E1-E2 ATPases phosphorylation site BL00154:G173-L190, I427-P445, D949-L989 PROFILESSCAN E1-E2 ATPases phosphorylation site: I413-A461 P-type cation-transporting ATPase PR00119:P431-P445, A965-D975, I1111-I1130 ATPASE HYDROLASE TRANSMEMBRANE PHOSPHORYLATION ATP-BINDING PROTEIN PROBABLE CALCIUMTRANSPORTING CALCIUM TRANSPORT PD004657:A1145-P1374 PD006317:Y162-E255 PD149930:C1085-P1144 PD004932:R65-P121 do ATPASE; CALCIUM; TRANSPORTING; DM02405 P32660 318-1225:R157-E475, E776-N1209 BLAST_DOMO E1-E2 ATPase motif: D433-T439 ATP/GTP binding site (P-loop): G1053-T1060 MOTIFS MOTIFS	ANALYTICAL METHODS AND DATABASES

Table 3 (cont.)

Table 4

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
31	2194064CB1	1129	1071-1129, 833-898	g5110579 FL2194064_g7770598_000019_g7 670446 6542780F9 (LNODNOK02)	1 203 32	485 1129 481
32	2744094CB1	2699	1-2196, 2541-2587	FL097646_00001 55058921H1 70317743D1 70317681D1	431 1 2347 2209	2542 793 2699 2639
33	2798241CB1	6369	1-1210, 1759-5012	71911330V1 70300809D1 6340750H1 (BRANDIN01) 7601441J1 (ESOGTME01) 6314138H1 (NERDTN03) 7690596H1 (PROSTME06) 7753104J1 (HEAONOE01) 4013186F9 (MUSCNOT10) 7606344H1 (COLRTUE01) 6913644H1 (PITUDIR01) 55052455J1 7400061H1 (SINIDME01) 2798241T6 (NPOLNOT01) 55058989J1 7100413F7 (BRAWTDRO2) 6744456H1 (BRAFNOT02) 55053647J1 6586921H1 (TLYMUNT03)	5832 5128 5650 4623 5235 4145 5764 3758 1764 4608 1981 1 1325 2548 483 568 3011 1157	6369 5690 6322 5186 5750 4636 6357 4391 2219 5181 2827 502 1955 3298 1185 1274 3823 1724
34	3105257CB1	2558	1-587, 2435-2558	70864718V1 70549000V1 FL3105257CT1_00001 6451207H1 (BRAINOC01)	1864 1608 1 1868	2353 2310 1843 2558

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
35	3200979CB1	5065	5030-5065, 1-3313	FL3200979_g3810670_g4240130 71698878V1	1 4463	4779 5065
36	6754139CB1	1677	1-686	656293H1 (BOSINOT03) 55062573H1 GBI:edit GNN:g8017750_000028_004 g5678193 6754139J1 (SINTFER02)	532 789 1 386 684 684	800 875 531 1677 883 874
37	6996659CB1	3714	1-1916, 3071-3091, 2092-2619	6996659F8 (BRAXTDR17) GBI.g9211864_01_04_05_12.edi t 55098348H2 1596150T6 (BRAINF014) 7124651F6 (COLNDIY01) g4622477 1596150F6 (BRAINF014) 55063531J1 7291716R6 (BRAIFER06) 7291716F6 (BRAIFER06) 55063924J1	1180 1303 2752 3116 2605 3322 2967 1 510 219 1768	1915 3006 2942 3707 2776 3714 3466 309 1209 1174 1994
38	7472747CB1	1009	1-388, 571- 704, 778- 1009	FL7472747_g6983242_000026_g3 925427 7616162H1 (COLNTUN03)	122 1 1	1009 450 1155
39	7474121CB1	1155	1-1155	GNN.g7259672_000014_002	1	1155
40	7475615CB1	2733	1852-2185, 1484-1579, 665-1340, 1-249, 2334-2733, 454-495	FL7475615_g8980204_000002_g2 654005_1_11-12 FL7475615_g8980204_000002_g2 654005_1_6-7 FL7475615_g8980204_000002_g2 654005_1_12-13	1580 986 1687	1756 1221 1849

Polynucleotide	Incyte	Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	Position
43		6952742CB1	2600	2329-2600	588402278 (LIVNON08) 55063579H1 GBI:97232144_000013.edlt.3 6816048H1 (ADRETTUR01) 6952742H1 (BRAITDR02) GNN:96970605_000013_002 GBI:97232144_000013.fasta.ed	5' 3'
44		7478795CB1	2917	2698-2917	72016954V1 71989431V1 72017820V1 72017055V1 72017371V1 72017076V1 72017430V1 55076285H1 GBI:98017750.edlt	5' 3'
45		6562933CB1	1474	1-362	FL656293_g8017750_000028_g67 46563_2-2-3 GBI:98017750_000028_g67	5' 3'
46		7473957CB1	1742	1-367	7675576H1 (NOSETVE01) 4648731F9 (PROSTUT20) 71166638V1 71165785V1 6830443J1 (SINTNOR01) 7761487H1 (THYMNOR02) 6770140H1 (BRAUNOR01) 7761487J1 (THYMNOR02) GNN:97243948_CDS_1	5' 3'
47		7474111CB1	2312	1-639	1860-1908 2004-2312 1686-1712	5' 3'

Table 4 (cont.)

Polynucleotide	Incyte	Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	Position
41		7475656CB1	3457	3284-3346	550733909H2 654005_1-10-11 FL7475615_g8980204_000002_g2 6806177J1 (SKIRNOR01) GNN:97342135_000012_002 654005_1-8-9 FL7475615_g8980204_000002_g2 1509180F6 (LUNGNOT14) 55083049H1 55092029J1 654005_1-7-8	5' 3'
42		7480632CB1	5622	1-3676	2428507R6 (SCORNON02) 5373417F8 (BRAINOT22) 5893974H1 (BRAYDIN03) 4787380H1 (BRATNOT03) 1450339F1 (PENTITUT01) 7270152H1 (OVARIDJ01) 71697049V1 3488927H1 (EPIGNOT01) 6774619J1 (OVARIDR01) 5063703F6 (ARFTTUT01) 55072886J1 GBI:93810670_000001.edlt 6488228F9 (MIXDUNB01) 7670233H2 (BONRNOCO1)	5' 3'

Table 4 (cont.)

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
48	7480826CB1	2320	161-224, 2044-2320	7752763J1 (HEAONE01) 60143671D1 6052064J1 (BRABDIR03) 6484950H1 (MIXDUNB01) 2944045H1 (BRAITUT23) 7469461H1 (LUNGNOE02)	1668 467 1080 1276 827 1	2320 917 1658 1723 1118 498
49	6025572CB1	1781	1-170	FL6025572_g7382154_000015_g1 197164 4923834H1 (TESTNOT11) g3838735 g3734777 71970611V1 6025572F6 (TESTNOT11)	347 1 1313 252 1285 883	1063 291 1781 472 1780 1627
50	5686561CB1	2433	1-1078, 1197-1275	71412362V1 6060785H1 (BRAENOT04) 7695065J1 (LNODTUE01) 7633409H1 (SINTDIE01) 3776733H1 (BRSTNOT27) 2802364F6 (PENCNOT01) 5564984F6 (TLYMNOT08) 70730430V1	1088 551 387 1 2148 1765 860 1525	1702 1100 1052 483 2433 2304 1528 2108
51	1553725CB1	1772	1571-1772	60211064U1 72050509V1 70300327D1 70300706D1 1553725X15C1 (BLADTUT04) 70300332D1	344 1176 984 1 54 729	823 1772 1428 262 694 1286
52	1695770CB1	1874	1-479, 1298-1874, 1131-1216, 886-984	55117454H1 55110123H1 55072985J1	1155 286 1	1874 1179 542

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
53	4672222CB1	6211	3238-3683, 4625-4798, 2313-2462, 1-1636	55047368J1 71007436V1 71998604V1 71995592V1 3462433F7 (293TF2T01) 71997753V1 71995863V1 55073038H1 55141177J1 71998657V1 6141577F6 (BMARTXT03) 55140386J1 GBI.g8189326.edit 5092011F6 (UTRSTMR01) 7743692H1 (ADRETUE04) 2505959F6 (CONUTUT01)	1925 5663 4613 3913 2738 4522 3346 818 2942 3811 1 1086 2957 1797 5374 5325	2815 6211 5344 4598 3162 5239 3886 1499 3318 4537 878 1915 3903 2436 5927 5866
54	6176128CB1	3714	1-197, 329- 2513, 3301- 3336	GBI.g979669_000005_000004.ed it 6859776H1 (BRAIFEN08) GBI.g979669_000002.edit GBI.g7739135_000005.edit 6772216J1 (BRAUNOR01) 6887873J1 (BRAITDR03) 8039114H1 (SPLNNOE01) 6907605J1 (PITUDIR01) 6445788H2 (BRAINOC01) 6891702F6 (BRAITDR03) 7065904R6 (BRATNOR01)	1 2265 3612 3115 2991 899 1741 2586 1383 543 383	1143 2953 3714 3711 3324 1503 2374 3088 2006 1053 645
55	7473418CB1	3115	1-1411	FL7473418_g3176728_g5531902_ 1_4-5 7056016H1 (BRALNON02)	369 2658	740 3115

58	Polynucleotide	Incye	Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
			7481461CBI	1840	1-91	70481006V1 70465445V1 17487222F6 (STOMTUT02) 60266587D1 7637372J1 (SINTDIE01) 65933739	551 673 1250 1840 1801 1225 221 642 407	2204 4634 3550 2001 526 4862 3426 1260 5348 4428 792 4131 3918 2994 3167 4903 4087 384 909 1573 4767 3836 2616 3434 2939 3947 3263 1126 1592 5149
59			7472541CBI	5348	1384-1560, 1-1188, 4239-4906, 2145-2970, 4944-5348	6772907J1 (BRAVOR01) 2182261F6 (SININOT01) 5459667H1 (SINITUT03) GNN.97710567_000006_002.ed1c 7362215H1 (BRAIFEE05) GNN.97708823_000019_002 7032970H1 (BRAXTDR12) 7313608H1 (BRABDIE02) 71462931V1 5767060H1 (STOMFE02) 8069315J1 (BRAIFEE05) 7582660H1 (BRAIFEC01) GNN.97454125_000004_002.ed1c 6772907H1 (BRAVOR01) GBI.93873182_000001.ed1c5p	1641 4301 3276 1057 1 4306 2692 731 4732 3887 91 3510 1382 2476 1 4087 384 909 1573 4767 3836 2616 3434 2939 3947 3263 1126 1592 5149	
60			6999183CBI	5149	1-1797, 4753-4852, 3028-3711, 2471-2667	72017610V1 55076606J1 55144834J1 55144835H1 72299322V1 72017349V1 55051672H1 6999183R8 (HEALDIR01) 72017145V1 GBI.93873182_000001.ed1c5p	4219 1126 3263 2154 2616 4767 3836 909 1573 1128 4087 4903 3167 2994 1 4087 384 909 1573 4767 3836 2616 3434 2939 3947 3263 1126 1592 5149	

Table 4 (cont.)

Seq ID NO:	Polynucleotide	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	Position	3'
56	7474129CBI	2846	1-1696, 2073-2846, 1777-2012		FL7473418_g3176728_g5531902_1-7-8 1324158F6 (LPRN0702) 2114 6899347H1 (LIVRTR01) 1074 FL7473418_g3176728_g5531902_1-5-6 548 70075691U1 2291 FL7473418_g3176728_g5531902_1-6-7 739 FL7473418_g3176728_g5531902_1-1-2 1 FL7473418_g3176728_g5531902_1-10-11 1351 FL7473418_g3176728_g5531902_1-2-3 103 7114876H1 (BRAENOK01) 1549 FL7473418_g3176728_g5531902_1-3-4 232 4895008P6 (LIVRUT12) 1752 55109928H1 2480 55109306J1 2660 55124533H1 832 55124525H1 1073 55073088J1 796 GBI.94544493_000005_000056.e 1	2240 2846 2660 832 1893 1208 1 1073 796 1	
57	7481414CBI	906	441-541, 262-348		GBI.94544493_000005_000056.e 55073088J1 55124525H1 55124533H1 55109306J1 55109928H1 4895008P6 (LIVRUT12) 1-3-4 FL7473418_g3176728_g5531902_1-3-4 232 547 1954	906 1208 1893 832 2660 2846 2240	

Table 4 (cont.)

Table 5

Polynucleotide SEQ ID NO:	Incyte Project ID	Representative Library
31	2194064CB1	THYRTUT03
32	2744094CB1	BRSTTUT15
33	2798241CB1	PROSTME06
34	3105257CB1	BLADNOT01
35	3200979CB1	PENITUT01
36	6754139CB1	BRSTNOR01
37	6996659CB1	BRAIFER06
38	7472747CB1	COLNTUN03
40	7475615CB1	LUNGNON07
41	7475656CB1	BRAINOT22
42	7480632CB1	PENITUT01
43	6952742CB1	LIVRNON08
44	7478795CB1	BRAENOT02
45	656293CB1	COLNNOT22
46	7473957CB1	BRAHTDR03
47	7474111CB1	THYMNOE02
48	7480826CB1	MIXDUNB01
49	6025572CB1	TESTNOT11
50	5686561CB1	BRAENOT04
51	1553725CB1	THYMNON04
52	1695770CB1	COLNNOT23
53	4672222CB1	PITUDIR01
54	6176128CB1	BRAITDR03
55	7473418CB1	LPARNOT02
56	7474129CB1	PLACNOT05
58	7481461CB1	OVARTUT05
59	7472541CB1	BRAIFEE05
60	6999183CB1	HEALDIR01

WO 02/12340

PCT/USN/24217

Table 6

Library	Vector	Library Description
BLADNOT01	PBLUESCRIPT	Library was constructed using RNA isolated from the bladder tissue of a 78-year-old Caucasian female, who died from an intracranial bleed. Patient history included basal cell carcinoma, arthritis, and chronic hypertension.
BRAENOT02	pINCY	Library was constructed using RNA isolated from posterior parietal cortex tissue removed from the brain of a 35-year-old Caucasian male who died from cardiac failure.
BRAENOT04	pINCY	Library was constructed using RNA isolated from inferior parietal cortex tissue removed from the brain of a 35-year-old Caucasian male who died from cardiac failure. Pathology indicated moderate leptomeningeal fibrosis and multiple microinfarctions of the cerebral neocortex. Patient history included dilated cardiomyopathy, congestive heart failure, cardiomegaly and an enlarged spleen and liver.
BRAHTDR03	PCDNA2.1	This random primed library was constructed using RNA isolated from archaecortex, anterior hippocampus tissue removed from a 55-year-old Caucasian female who died from cholangiocarcinoma. Pathology indicated mild meningeal fibrosis predominately over the convexities, scattered axonal spheroids in the white matter of the cingulate cortex and the thalamus, and a few scattered neurofibrillary tangles in the entorhinal cortex and the periaqueductal gray region. Pathology for the associated tumor tissue indicated well-differentiated cholangiocarcinoma of the liver with residual or relapsed tumor. Patient history included cholangiocarcinoma, post-operative Budd-Chiari syndrome, biliary ascites, hydrothorax, dehydration, malnutrition, oliguria and acute renal failure. Previous surgeries included cholecystectomy and resection of 85% of the liver.
BRAIFEE05	PCDNA2.1	This 5' biased random primed library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus who was stillborn with a hypoplastic left heart at 23 weeks' gestation.
BRAIFER06	PCDNA2.1	This random primed library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus who was stillborn with a hypoplastic left heart at 23 weeks' gestation. Serologies were negative.

WO 02/12340

PCT/USN/24217

Table 6 (cont.)

Library	Vector	Library Description
COLN0722	PINCY	Library was constructed using RNA isolated from a 56-year-old Caucasian female with Crohn's disease during a partial resection of the small intestine. Pathology indicated Crohn's disease of the ileum and ileal-colonic anastomosis, causing a fistula at the anastomotic site that extended into pericolic fat. The ileal mucosa showed linear and punctate ulcers with intervening normal tissue. Previous surgeries included a partial ileal resection and permanent ileostomy. Family history included irritable bowel syndrome in the mother and the siblings.
COLN0723	PINCY	Library was constructed using RNA isolated from a 16-year-old Caucasian male during a total colectomy with abdominal/pelvic resection. Pathology indicated gastritis and pancolitis consistent with the acute phase of ulcerative colitis. Inflammation was more severe in the transverse colon, with inflammation confined to the mucosa. There was only mild involvement of the ascending and sigmoid colon, and no significant involvement of the cecum, rectum, or terminal ileum. Family history included irritable bowel syndrome.
COLN0703	PINCY	This normalized pooled colon tumor tissue library was constructed from 1.16 million independent clones from a pooled colon tumor library. Starting library was constructed using pooled cDNA from 6 donors. cDNA was generated using mRNA isolated from colon tumor tissue removed from a 55-year-old Caucasian male (A) during hemicolectomy; from a 60-year-old Caucasian male (B) during hemicolectomy; from a 62-year-old Caucasian male (C) during sigmoidectomy; from a 30-year-old Caucasian female (D) during hemicolectomy; from a 64-year-old Caucasian female (E) during hemicolectomy; and from a 70-year-old Caucasian female (F) during hemicolectomy. Pathology indicated invasive grade 3 adenocarcinoma (A); invasive grade 2 adenocarcinoma (B); invasive grade 2 adenocarcinoma (C); carcinoma tumor grade 2 adenocarcinoma (E); and invasive grade 2 adenocarcinoma (F). Donors B, C, D, E, and F had positive lymph nodes. Patient medications included Activan (A); Seldane (B); Tri-Leylen (D); Synthroid (E); Tamoxifen, prednisone, Synthroid, and glibizide (F). The library was normalized in two rounds using conditions adapted from Soares et al., PNAS (1994) 91:9228-9232 and Bonaldo et al., Genome Research 6 (1996):791, except that a significantly longer (48 hours/round) reannealing hybridization was used.

Table 6 (cont.)

Library	Vector	Library Description
BRAIN0722	PINCY	Library was constructed using RNA isolated from right temporal lobe tissue removed from a 45-year-old Black male during a brain lobectomy. Pathology for the associated tumor tissue indicated dysembryoplastic neuroepithelial tumor of the right temporal lobe. The right temporal region dura was consistently with calcifying pseudotumor of the neurexus. Family history included obesity, benign hypertension, cirrhosis of the liver, obesity, hyperlipidemia, cerebrovascular disease, and type II diabetes.
BRAITDR03	PCDNA2.1	This random primed library was constructed using RNA isolated from a 55-year-old Caucasian female who died from cholangiocarcinoma. Pathology indicated mild meningioma fibrosis predominantly over the convolutes, scattered axonal spheroids in the white matter of the cingulate cortex and the thalamus, and a few scattered neurofibrillary tangles in the entorhinal cortex and the periaqueductal gray region. Pathology for the associated tumor tissue indicated well-differentiated cholangiocarcinoma of the liver with residual or relapsed tumor. Patient history included cholangiocarcinoma, post-operative Budd-Chiari syndrome, biliary ascites, hydrothorax, dehydration, malnutrition, oliguria and acute renal failure. Previous surgeries included cholecystectomy and resection of 85% of the liver.
BRS10R01	PINCY	Library was constructed using RNA isolated from diseased breast tissue removed from a 59-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology for the associated tumor tissue indicated an invasive lobular carcinoma with extension into ducts. Patient history included cirrhosis, esophageal ulcer, hyperlipidemia, and neuropathy.
BRS10T015	PINCY	Library was constructed using RNA isolated from breast tissue removed from a 46-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated invasive grade 3, nuclear grade 2 adenocarcinoma, ductal type. An intraductal carcinoma component, non-comedo, comprised approximately 50% of the neoplasm, including the lactiferous ducts. Angiolymphatic involvement was present. Metastatic adenocarcinoma was present in 7 of 10 axillary lymph nodes. The largest nodal metastasis measured 3 cm, and focal extracapsular extension was identified. Family history included atherosclerotic coronary artery disease, type II diabetes, cerebrovascular disease, and depressive disorder.

Table 6 (cont.)

Library	Vector	Library Description
HEALDIR01	PCDNA2.1	This random primed library was constructed using RNA isolated from diseased left ventricle tissue removed from a 7-month-old Caucasian male who died from cardiopulmonary arrest due to Pompe's disease. Patient history included Pompe's disease, left ventricular hypertrophy, pyrexia, right complete cleft lip, cleft palate, chronic serous otitis media, hypertrophic cardiomyopathy, congestive heart failure, and developmental delays. Family history included acute myocardial infarction, diabetes, cystic fibrosis and Down's syndrome.
LIVRNON08	pINCY	This normalized library was constructed from 5.7 million independent clones from a pooled liver tissue library. Starting RNA was made from pooled liver tissue removed from a 4-year-old Hispanic male who died from anoxia and a 16 week female fetus who died after 16-weeks gestation from anencephaly. Serologies were positive for cytomegalovirus in the 4-year-old. Patient history included asthma in the 4-year-old. Family history included taking daily prenatal vitamins and mitral valve prolapse in the mother of the fetus. The library was normalized in 2 rounds using conditions adapted from Soares et al., PNAS (1994) 91:9228 and Bonaldo et al., Genome Research 6 (1996):791, except that a significantly longer (48 hours/round) reannealing hybridization was used.
LPARNOT02	pINCY	Library was constructed using RNA isolated from tissue obtained from the left parotid (salivary) gland of a 70-year-old male with parotid cancer.
LUNGNON07	pINCY	This normalized lung tissue library was constructed from 5.1 million independent clones from a lung tissue library. Starting RNA was made from RNA isolated from lung tissue. The library was normalized in two rounds using conditions adapted from Soares et al., PNAS (1994) 91:9228-9232 and Bonaldo et al., Genome Research (1996) 6:791, except that a significantly longer (48 hours/round) reannealing hybridization was used.

Table 6 (cont.)

Library	Vector	Library Description
MIXDUNB01	pINCY	Library was constructed using RNA isolated from myometrium removed from a 41-year-old Caucasian female (A) during vaginal hysterectomy with a dilatation and curettage and untreated smooth muscle cells removed from the renal vein of a 57-year-old Caucasian male. Pathology for donor A indicated the myometrium and cervix were unremarkable. The endometrium was secretory and contained fragments of endometrial polyps. Benign endo- and ectocervical mucosa were identified in the endocervix. Pathology for the associated tumor tissue indicated uterine leiomyoma. Medical history included an unspecified menstrual disorder, ventral hernia, normal delivery, a benign ovarian neoplasm, and tobacco abuse in donor A. Previous surgeries included a bilateral destruction of fallopian tubes, removal of a solitary ovary, and an exploratory laparotomy in donor A. Medications included ferrous sulfate in donor A.
OVARTUT05	pINCY	Library was constructed using RNA isolated from ovarian tumor tissue removed from a 62-year-old Caucasian female during a total abdominal hysterectomy, removal of the fallopian tubes and ovaries, exploratory laparotomy, regional lymph node excision, and dilation and curettage. Pathology indicated a grade 4 endometrioid carcinoma with extensive squamous differentiation, forming a solid mass in the right ovary. The uterine endometrium was inactive, the cervix showed mild chronic cervicitis, and focal endometriosis was observed in the posterior uterine serosa. Curettings indicated weakly proliferative endometrium with excessive stromal breakdown in the uterus, and a prior cervical biopsy indicated mild chronic cervicitis with a prominent nabothian cyst in the cervix. Patient history included longitudinal deficiency of the radioulna, osteoarthritis, thrombophlebitis, and abnormal blood chemistries. Family history included atherosclerotic coronary artery disease, pulmonary embolism, and cerebrovascular disease.
PENITUT01	pINCY	Library was constructed using RNA isolated from tumor tissue removed from the penis of a 64-year-old Caucasian male during penile amputation. Pathology indicated a fungating invasive grade 4 squamous cell carcinoma involving the inner wall of the foreskin and extending onto the glans penis. Patient history included benign neoplasm of the large bowel, atherosclerotic coronary artery disease, angina pectoris, gout, and obesity. Family history included malignant pharyngeal neoplasm, chronic lymphocytic leukemia, and chronic liver disease.

Table 6 (cont.)

Library	Vector	Library Description
THYMOB02	PCDNA2.1	This 5' biased random primed library was constructed using RNA isolated from thymus tissue removed from a 3-year-old Hispanic male during a thymectomy and closure of a patent ductus arteriosus. The patient presented with severe pulmonary stenosis and cyanosis. Patient history included a cardiac catheterization and echocardiogram. Previous surgeries included Blalock-Taussig shunt and pulmonary valvotomy. The patient was not taking any medications. Family history included benign hypertension, osteoarthritis, depressive disorder, and extrinsic asthma in the grandparent(s).
THYMON04	PSORT1	This normalized library was constructed from a thymus tissue library. Starting RNA was made from thymus tissue removed from a 3-year-old Caucasian male, who died from anoxia. Serologies were negative. The patient was not taking any medications. The library was normalized in two rounds using conditions adapted from Soares et al., PNAS (1994) 91:9228 and Bonaldo et al., Genome Research (1996) 6:791, except that a significantly longer (48-hours/round) reannealing hybridization was used.
THYRT03	PINCY	Library was constructed using RNA isolated from benign thyroid tumor tissue removed from a 17-year-old Caucasian male during a thyroidectomy. Pathology indicated encapsulated follicular adenoma forming a circumscribed mass.

Table 6 (cont.)

Library	Vector	Library Description
PITUDIR01	PCDNA2.1	This random primed library was constructed using RNA isolated from pituitary gland tissue removed from a 70-year-old female who died from metastatic adenocarcinoma.
PLACNO705	PINCY	Library was constructed using RNA isolated from placental tissue removed from a Caucasian male fetus, who died after 18 weeks' gestation from fetal demise.
PROSTM06	PCDNA2.1	This 5' biased random primed library was constructed using RNA isolated from diseased prostate tissue removed from a 57-year-old Caucasian male during closed prostate biopsy, radical prostatectomy, and regional lymph node excision. Pathology indicated adenocarcinoma, Gleason grade 3+3, forming a predominant mass involving the right side centrally. The patient presented with elevated prostate specific antigen and prostate cancer. Patient history included cholecystectomy, repair of diaphragm hernia, and repair of vertebral fracture. Patient medications included Pepcid, Omeprazole, and Dulcetin. Family history included benign hypertension, cerebrovascular accident, atherosclerotic coronary artery disease, uterine cancer and type II diabetes in the mother; prostate cancer in the father; drug abuse, prostate cancer, and breast cancer in the sibling(s).
TESTNO711	PINCY	Library was constructed using RNA isolated from testicular tissue removed from a 16-year-old Caucasian male who died from hanging. Patient history included drug use (cocaine, marijuana, and cocaine use), and medications included Lithium, Ritalin, and Paxil.

Table 7

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) <i>J. Mol. Biol.</i> 215:403-410; Altschul, S.F. et al. (1997) <i>Nucleic Acids Res.</i> 25:3389-3402.	<i>ESTs</i> : Probability value= 1.0E-8 or less <i>Full Length sequences</i> : Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and search.	Pearson, W.R. and D.J. Lipman (1988) <i>Proc. Natl. Acad. Sci. USA</i> 85:2444-2448; Pearson, W.R. (1990) <i>Methods Enzymol.</i> 183:63-98; and Smith, T.F. and M.S. Waterman (1981) <i>Adv. Appl. Math.</i> 2:482-489.	<i>ESTs</i> : fasta E value=1.06E-6 <i>Assembled ESTs</i> : fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less <i>Full Length sequences</i> : fastx score=100 or greater
BLIMPS	A BLOCKS IMPROVED Searcher that matches a sequence against those in BLOCKS, PRINTS, DMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) <i>Nucleic Acids Res.</i> 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) <i>Methods Enzymol.</i> 266:88-105; and Attwood, T.K. et al. (1997) <i>J. Chem. Inf. Comput. Sci.</i> 37:417-424.	Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) <i>J. Mol. Biol.</i> 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) <i>Nucleic Acids Res.</i> 26:320-322; Durbin, R. et al. (1998) <i>Our World View</i> , in a Nutshell, Cambridge Univ. Press, pp. 1-350.	<i>PFAM hits</i> : Probability value= 1.0E-3 or less <i>Signal peptide hits</i> : Score= 0 or greater

131

Table 7 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) <i>CABIOS</i> 4:61-66; Gribskov, M. et al. (1989) <i>Methods Enzymol.</i> 183:146-159; Bairoch, A. et al. (1997) <i>Nucleic Acids Res.</i> 25:217-221.	Normalized quality score \geq GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) <i>Genome Res.</i> 8:175-185; Ewing, B. and P. Green (1998) <i>Genome Res.</i> 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) <i>Adv. Appl. Math.</i> 2:482-489; Smith, T.F. and M.S. Waterman (1981) <i>J. Mol. Biol.</i> 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) <i>Genome Res.</i> 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) <i>Protein Engineering</i> 10:1-6; Claverie, J.M. and S. Audic (1997) <i>CABIOS</i> 12:431-439.	Score=3.5 or greater
TMAP	A program that uses weight matrices to delineate transmembrane segments on protein sequences and determine orientation.	Persson, B. and P. Argos (1994) <i>J. Mol. Biol.</i> 237:182-192; Persson, B. and P. Argos (1996) <i>Protein Sci.</i> 5:363-371.	
TMHMMER	A program that uses a hidden Markov model (HMM) to delineate transmembrane segments on protein sequences and determine orientation.	Sonnhammer, E.L. et al. (1998) <i>Proc. Sixth Intl. Conf. on Intelligent Systems for Mol. Biol.</i> , Glasgow et al., eds., The Am. Assoc. for Artificial Intelligence Press, Menlo Park, CA, pp. 175-182.	
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) <i>Nucleic Acids Res.</i> 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

132

What is claimed is:

1. An isolated polypeptide selected from the group consisting of:
 - a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-30,
 - b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-30,
 - c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, and
 - d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30.

2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-30.

3. An isolated polynucleotide encoding a polypeptide of claim 1.

4. An isolated polynucleotide encoding a polypeptide of claim 2.

5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:31-60.

6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

7. A cell transformed with a recombinant polynucleotide of claim 6.

8. A transgenic organism comprising a recombinant polynucleotide of claim 6.

9. A method of producing a polypeptide of claim 1, the method comprising:

- a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and

- b) recovering the polypeptide so expressed.

10. An isolated antibody which specifically binds to a polypeptide of claim 1.

11. An isolated polynucleotide selected from the group consisting of:

- a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:31-60,
- b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:31-60,
- c) a polynucleotide complementary to a polynucleotide of a),
- d) a polynucleotide complementary to a polynucleotide of b), and
- e) an RNA equivalent of a)-d).

12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.

13. A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

15. A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and
- b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

16. A composition comprising a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

17. A composition of claim 16, wherein the polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO:1-30.

18. A method for treating a disease or condition associated with decreased expression of functional TRIC^h, comprising administering to a patient in need of such treatment the composition of claim 16.

19. A method of screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

21. A method for treating a disease or condition associated with decreased expression of functional TRIC^h, comprising administering to a patient in need of such treatment a composition of claim 20.

22. A method of screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

24. A method for treating a disease or condition associated with overexpression of functional TRIC^h, comprising administering to a patient in need of such treatment a composition of claim 23.

25. A method of screening for a compound that specifically binds to the polypeptide of claim

1, the method comprising:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and

b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, the method comprising:

- a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,

b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and

c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

27. A method of screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,

b) detecting altered expression of the target polynucleotide, and

c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

28. A method of assessing toxicity of a test compound, the method comprising:

- a) treating a biological sample containing nucleic acids with the test compound,

b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof,

c) quantifying the amount of hybridization complex, and

d) comparing the amount of hybridization complex in the treated biological sample with the

amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

29. A diagnostic test for a condition or disease associated with the expression of TRICH in a biological sample, the method comprising:
- combining the biological sample with an antibody of claim 10, under conditions suitable for the antibody to bind the polypeptide and form an antibody:polypeptide complex, and
 - detecting the complex, wherein the presence of the complex correlates with the presence of the polypeptide in the biological sample.

30. The antibody of claim 10, wherein the antibody is:

- a chimeric antibody,
- a single chain antibody,
- a Fab fragment,
- a F(ab')₂ fragment, or
- a humanized antibody.

31. A composition comprising an antibody of claim 10 and an acceptable excipient.

32. A method of diagnosing a condition or disease associated with the expression of TRICH in a subject, comprising administering to said subject an effective amount of the composition of claim 31.

33. A composition of claim 31, wherein the antibody is labeled.

34. A method of diagnosing a condition or disease associated with the expression of TRICH in a subject, comprising administering to said subject an effective amount of the composition of claim 33.

35. A method of preparing a polyclonal antibody with the specificity of the antibody of claim 10, the method comprising:

- immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, or an immunogenic fragment thereof, under conditions to

elicit an antibody response,

- isolating antibodies from said animal, and
- screening the isolated antibodies with the polypeptide, thereby identifying a polyclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30.

36. An antibody produced by a method of claim 35.

37. A composition comprising the antibody of claim 36 and a suitable carrier.

38. A method of making a monoclonal antibody with the specificity of the antibody of claim 10, the method comprising:

- immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30, or an immunogenic fragment thereof, under conditions to elicit an antibody response,
- isolating antibody producing cells from the animal,
- fusing the antibody producing cells with immortalized cells to form monoclonal antibody-producing hybridoma cells,
- culturing the hybridoma cells, and
- isolating from the culture monoclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30.

39. A monoclonal antibody produced by a method of claim 38.

40. A composition comprising the antibody of claim 39 and a suitable carrier.

41. The antibody of claim 10, wherein the antibody is produced by screening a Fab expression library.

42. The antibody of claim 10, wherein the antibody is produced by screening a recombinant immunoglobulin library.

43. A method of detecting a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30 in a sample, the method comprising:

- a) incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide, and
- b) detecting specific binding, wherein specific binding indicates the presence of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30 in the sample.

44. A method of purifying a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30 from a sample, the method comprising:

- a) incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide, and
- b) separating the antibody from the sample and obtaining the purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-30.

45. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:1.

46. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:2.

47. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:3.

48. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:4.

49. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:5.

50. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:6.

51. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:7.

52. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:8.

53. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:9.

54. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:10.

55. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:11.

56. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:12.

57. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:13.

58. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:14.

59. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:15.

60. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:16.

61. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:17.

62. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:18.

63. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:19.

64. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:20.

65. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:21.

66. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:22.

67. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:23.

68. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:24.

69. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:25.

70. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:26.

71. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:27.

72. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:28.

73. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:29.

74. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:30.

5 75. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:31.

76. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:32.

10 77. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:33.

15 78. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:34.

79. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:35.

20 80. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:36.

81. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:37.

25 82. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:38.

30 83. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:39.

84. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:40.

85. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:41.

5 86. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:42.

87. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:43.

10 88. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:44.

89. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:45.

15 90. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:46.

20 91. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:47.

92. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:48.

25 93. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:49.

94. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:50.

30 95. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:51.

96. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID

NO:52.

97. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:53.

98. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:54.

99. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:55.

100. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:56.

101. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:57.

102. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:58.

103. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:59.

104. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:60.

<110> INCYTE GENOMICS, INC.

YUE, Henry

THORNTON, Michael

RAKUCMAR, Jajalaxmi

TANG, Y. Tom

AZIMZAI, Yalda

BAUGHN, Mariah R.

YANG, Junming

YAO, Monique G.

LAL, Preeti

WALIA, Narinder K.

GANDHI, Ameena R.

HAFALIA, April J.A.

NGUYEN, Daniel B.

PATERSON, Chandra

ELLIOTT, Vicki S.

TRIBOUTLEY, Catherine M.

LU, Dzung Alma M.

XU, Yuning

REDDY, Roopa

HERNANDEZ, Roberto

BOROMSKY, Mark L.

LO, Terence P.

LU, Yan

POLICKY, Jennifer L.

GREENE, Barrie D.

SANTANWALA, Madhu S.

RAUDMAN, Brigitte E.

BURFORD, Neil

ISON, Craig H.

LEE, Ernestine A.

DING, Li

DAS, Debopriya

KALLICK, Deborah A.

KHAN, Farrah A.

SEILHNER, Jeffrey J.

<120> TRANSPORTERS AND ION CHANNELS

<130> PT-0183 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/223,269; 60/224,456; 60/226,410; 60/228,140; 60/230,067;

60/231,434

<151> 2000-08-03; 2000-08-10; 2000-08-18; 2000-08-25; 2000-08-31;

2000-09-08

<160> 60

<170> PERL Program

<210> 1

<211> 308

<212> PRG

<213> Homo sapiens

<220>

<221> misc feature

<223> Incyte ID No: 2194064CD1

<400> 1

Met Val Gly Gly Val Leu Ala Ser Leu Gly Phe Val Phe Ser Ala

1 Phe Ala Ser Asp Leu Leu His Leu Tyr Leu Gly Leu Gly Leu Leu 15
20 Ala Gly Phe Gly Trp Ala Leu Val Phe Ala Pro Ala Leu Gly Thr 30
35 Leu Ser Arg Tyr Phe Ser Arg Arg Val Leu Ala Val Gly Leu 45
50 Ala Leu Thr Gly Asn Gly Ala Ser Ser Leu Leu Leu Ala Pro Ala 60
65 Leu Gln Leu Leu Asp Thr Phe Gly Trp Arg Gly Ala Leu Leu 75
80 Leu Leu Gly Ala Ile Thr Leu His Leu Thr Pro Cys Gly Ala Leu 90
95 Leu Leu Pro Leu Val Leu Pro Gly Asp Pro Pro Ala Pro Arg 105
110 Ser Pro Leu Ala Ala Leu Gly Thr Ala Leu Val Gly Gly Tyr 120
125 Phe Ser Ile Phe Ala Leu Gly Thr Ala Leu Val Gly Gly Thr 135
140 Phe Val Pro Tyr Val His Leu Ala Pro His Ala Leu Asp Arg 145
155 Leu Gly Gly Tyr Gly Ala Leu Val Val Ala Val Ala Ala Met 165
170 Gly Asp Ala Gly Ala Arg Leu Val Cys Gly Trp Leu Ala Asp 180
185 Gly Trp Val Pro Leu Pro Arg Leu Leu Ala Val Phe Gly Ala 195
200 Thr Gly Leu Gly Leu Trp Val Val Gly Leu Val Pro Val Val 210
215 Gly Gln Glu Ser Trp Gly Gly Pro Leu Leu Ala Ala Val Ala 225
230 Tyr Gly Leu Ser Ala Gly Ser Tyr Ala Pro Leu Val Phe Gly 240
245 Leu Pro Gly Leu Val Gly Val Gly Val Val Gln Ala Thr Gly 255
260 Leu Val Met Met Leu Met Ser Leu Gly Gly Leu Leu Gly Pro 270
275 Leu Ser Gly Lys Asp Leu Ser Ser Gln Ile Cys Leu Gln Leu Ser 285
290 Ser Ala Pro Gly Val Arg Gly Phe 300
305
<210> 2
<211> 606
<212> PRT
<213> Homo sapiens
<220>
<221> misc_feature
<223> Incyte ID No: 2744094CD1
<400> 2
Met Ala Glu Gln Leu Ser Gln Gln Leu Pro Arg Thr Cys Leu Trp 15
1 His Leu Tyr Ile Thr Thr Val Ser Leu Pro Gly Tyr Met Val Ser 20
25 Cys Ile Ile Phe Phe Val Val Pro Ile Val Phe Leu Thr Ile 30
35 Phe Ser Phe Trp Trp Leu Ser Tyr Trp Trp Leu Gln Gly Ser Gly 40
45 Thr Asn Ser Ser Arg Glu Ser Asn Gly Thr Met Ala Asp Leu Gly 50
55 Asn Ile Ala Asp Asn Pro Gln Leu Ser Phe Tyr Gln Leu Val Tyr 60
65

80 Gly Leu Asn Ala Leu Leu Leu Ile Cys Val Gly Val Cys Ser Ser 85
95 Gly Ile Phe Thr Lys Val Thr Arg Lys Ala Ser Thr Ala Leu His 100
110 Asn Lys Leu Phe Asn Lys Val Phe Arg Cys Pro Met Ser Phe Phe 115
125 Asp Thr Ile Pro Ile Gly Arg Leu Leu Asn Cys Phe Ala Gly Asp 130
140 Leu Gln Gln Leu Asp Gln Leu Leu Pro Ile Phe Ser Glu Gln Phe 145
155 Leu Val Leu Ser Leu Met Val Ile Ala Val Leu Leu Ile Val Ser 160
170 Val Leu Ser Pro Tyr Ile Leu Leu Met Gly Ala Ile Ile Met Val 175
185 Ile Cys Phe Ile Tyr Tyr Met Met Phe Lys Lys Ala Ile Gly Val 190
200 Phe Lys Arg Leu Glu Asn Tyr Ser Arg Ser Pro Leu Phe Ser His 205
215 Ile Leu Asn Ser Leu Gln Gly Leu Ser Ser Ile His Val Tyr Gly 220
230 Lys Thr Glu Asp Phe Ile Ser Gln Phe Lys Arg Leu Thr Asp Ala 235
245 Gln Asn Asn Tyr Leu Leu Leu Phe Leu Ser Ser Thr Arg Trp Met 250
260 Ala Leu Arg Leu Glu Ile Met Thr Asn Leu Val Thr Leu Ala Val 265
275 Ala Leu Phe Val Ala Phe Gly Ile Ser Ser Thr Pro Tyr Ser Phe 280
290 Lys Val Met Ala Val Asn Ile Val Leu Gln Leu Ala Ser Ser Phe 295
305 Gln Ala Thr Ala Arg Ile Gly Leu Glu Thr Glu Ala Gln Phe Thr 310
320 Ala Val Glu Arg Ile Leu Gln Tyr Met Lys Met Cys Val Ser Glu 325
335 Ala Pro Leu His Met Glu Gly Thr Ser Cys Pro Gln Gly Trp Pro 340
350 Gln His Gly Glu Ile Ile Phe Gln Asp Tyr His Met Lys Tyr Arg 355
365 Asp Asn Thr Pro Thr Val Leu His Gly Ile Asn Leu Thr Ile Arg 370
380 Gly His Glu Val Val Gly Ile Val Gly Arg Thr Gly Ser Gly Lys 385
395 Ser Ser Leu Gly Met Ala Leu Phe Arg Leu Val Glu Pro Met Ala 400
410 Gly Arg Ile Leu Ile Asp Gly Val Asp Ile Cys Ser Ile Gly Leu 415
425 Glu Asp Leu Arg Ser Lys Leu Ser Val Ile Pro Gln Asp Pro Val 430
440 Leu Leu Ser Gly Thr Ile Arg Phe Asn Leu Asp Pro Phe Asp Arg 445
455 His Thr Asp Gln Gln Ile Trp Asp Ala Leu Glu Arg Thr Phe Leu 460
470 Thr Lys Ala Ile Ser Lys Phe Pro Lys Lys Leu His Thr Asp Val 475
485 Val Glu Asn Gly Tyr Phe Ser Val Gly Glu Arg Gln Leu Leu 490
500 Cys Ile Ala Arg Ala Val Leu Arg Asn Ser Lys Ile Ile Leu Ile 505
515 Asp Glu Ala Thr Ala Ser Ile Asp Met Glu Thr Asp Thr Leu Ile 520
530 Gln Arg Thr Ile Arg Glu Ala Phe Gln Gly Cys Thr Val Leu Val 535
545 550

Ile Ala His Arg Val Thr Thr Val Leu Asn Cys Asp Arg Ile Leu
560 565 570
Val Met Gly Asn Gly Lys Val Val Glu Phe Asp Arg Pro Glu Val
575 580 585
Leu Arg Lys Lys Pro Gly Ser Leu Phe Ala Ala Leu Met Ala Thr
590 595 600
Ala Thr Ser Ser Leu Arg 605

<210> 3
<211> 1642
<212> PRT
<213> Homo sapiens

<220>
<221> misc.feature
<223> Incyte ID No: 2798241CD1

<400> 3
Met Ser Thr Ala Ile Arg Glu Val Gly Val Trp Arg Gln Thr Arg
1 5 10 15
Thr Leu Leu Leu Lys Asn Tyr Leu Ile Lys Cys Arg Thr Lys Lys
20 25 30
Ser Ser Val Gln Glu Ile Leu Phe Pro Leu Phe Phe Leu Trp
35 40 45
Leu Ile Leu Ile Ser Met Met His Pro Asn Lys Lys Tyr Glu Glu
50 55 60
Val Pro Asn Ile Glu Leu Asn Pro Met Asp Lys Phe Thr Leu Ser
65 70 75
Asn Leu Ile Leu Gly Tyr Thr Pro Val Thr Asn Ile Thr Ser Ser
80 85 90
Ile Met Gln Lys Val Ser Thr Asp His Leu Pro Asp Val Ile Ile
95 100 105
Thr Glu Glu Tyr Thr Asn Glu Lys Glu Met Leu Thr Ser Ser Lys
110 115 120
Ser Lys Pro Ser Asn Phe Val Gly Val Val Phe Lys Asp Ser Met
125 130 135
Ser Tyr Glu Leu Arg Phe Pro Asp Met Ile Pro Val Ser Ser
140 145 150
Ile Tyr Met Asp Ser Arg Ala Gly Cys Ser Lys Ser Cys Glu Ala
155 160 165
Ala Gln Tyr Trp Ser Ser Gly Phe Thr Val Leu Gln Ala Ser Ile
170 175 180
Asp Ala Ala Ile Ile Gln Leu Lys Thr Asn Val Ser Leu Trp Lys
185 190 195
Glu Leu Glu Ser Thr Lys Ala Val Ile Met Gly Glu Thr Ala Val
200 205 210
Val Glu Ile Asp Thr Phe Pro Arg Gly Val Ile Leu Ile Tyr Leu
215 220 225
Val Ile Ala Phe Ser Pro Phe Gly Tyr Phe Leu Ala Ile His Ile
230 235 240
Val Ala Glu Lys Glu Lys Lys Ile Lys Glu Phe Leu Lys Ile Met
245 250 255
Gly Leu His Asp Thr Ala Phe Trp Leu Ser Trp Val Leu Leu Tyr
260 265 270
Thr Ser Leu Ile Phe Leu Met Ser Leu Leu Met Ala Val Ile Ala
275 280 285
Thr Ala Ser Leu Leu Phe Pro Gln Ser Ser Ile Val Ile Phe
290 295 300
Leu Leu Phe Phe Leu Tyr Gly Leu Ser Ser Val Phe Phe Ala Leu
305 310 315
Met Leu Thr Pro Leu Phe Lys Lys Ser Lys His Val Gly Ile Val
320 325 330

Glu Phe Phe Val Thr Val Ala Phe Gly Phe Ile Gly Leu Met Ile
335 340 345
Ile Leu Ile Glu Ser Phe Pro Lys Ser Leu Val Trp Leu Phe Ser
350 355 360
Pro Phe Cys His Cys Thr Phe Val Ile Gly Ile Ala Gln Val Met
365 370 375
His Leu Glu Asp Phe Asn Glu Gly Ala Ser Phe Ser Asn Leu Thr
380 385 390
Ala Gly Pro Tyr Pro Leu Ile Ile Thr Ile Ile Met Leu Thr Leu
395 400 405
Asn Ser Ile Phe Tyr Val Leu Leu Ala Val Tyr Leu Asp Gln Val
410 415 420
Ile Pro Gly Glu Phe Gly Leu Arg Arg Ser Ser Leu Tyr Phe Leu
425 430 435
Lys Pro Ser Tyr Trp Ser Lys Ser Lys Arg Asn Tyr Glu Glu Leu
440 445 450
Ser Glu Gly Asn Val Asn Gly Asn Ile Ser Phe Ser Glu Ile Ile
455 460 465
Glu Pro Val Ser Ser Glu Phe Val Gly Lys Glu Ala Ile Arg Ile
470 475 480
Ser Gly Ile Gln Lys Thr Tyr Arg Lys Lys Gly Glu Asn Val Glu
485 490 495
Ala Leu Arg Asn Leu Ser Phe Asp Ile Tyr Glu Gly Gln Ile Thr
500 505 510
Ala Leu Leu Gly His Ser Gly Thr Gly Lys Ser Thr Leu Met Asn
515 520 525
Ile Leu Cys Gly Leu Cys Pro Pro Ser Asp Gly Phe Ala Ser Ile
530 535 540
Tyr Gly His Arg Val Ser Glu Ile Asp Glu Met Phe Glu Ala Arg
545 550 555
Lys Met Ile Gly Ile Cys Pro Gln Leu Asp Ile His Phe Asp Val
560 565 570
Leu Thr Val Glu Glu Asn Leu Ser Ile Leu Ala Ser Ile Lys Gly
575 580 585
Ile Pro Ala Asn Asn Ile Ile Gln Glu Val Gln Lys Val Leu Leu
590 595 600
Asp Leu Asp Met Gln Thr Ile Lys Asp Asn Gln Ala Lys Lys Leu
605 610 615
Ser Gly Gly Gln Lys Arg Lys Leu Ser Leu Gly Ile Ala Val Leu
620 625 630
Gly Asn Pro Lys Ile Leu Leu Leu Asp Glu Pro Thr Ala Gly Met
635 640 645
Asp Pro Cys Ser Arg His Ile Val Trp Asn Leu Lys Tyr Arg
650 655 660
Lys Ala Asn Arg Val Thr Val Phe Ser Thr His Phe Met Asp Glu
665 670 675
Ala Asp Ile Leu Ala Asp Arg Lys Ala Val Ile Ser Gln Gly Met
680 685 690
Leu Lys Cys Val Gly Ser Ser Met Phe Leu Lys Ser Lys Trp Gly
695 700 705
Ile Gly Tyr Arg Leu Ser Met Tyr Ile Asp Lys Tyr Cys Ala Thr
710 715 720
Glu Ser Leu Ser Ser Leu Val Lys Gln His Ile Pro Gly Ala Thr
725 730 735
Leu Leu Gln Gln Asn Asp Gln Gln Leu Val Tyr Ser Leu Pro Phe
740 745 750
Lys Asp Met Asp Lys Phe Ser Gly Leu Phe Ser Ala Leu Asp Ser
755 760 765
His Ser Asn Leu Gly Val Ile Ser Tyr Gly Val Ser Met Thr Thr
770 775 780
Leu Glu Asp Val Phe Leu Lys Leu Glu Val Glu Ala Glu Ile Asp
785 790 795
Gln Ala Asp Tyr Ser Val Phe Thr Gln Gln Pro Leu Glu Glu Glu

Met Asp Ser Lys Ser Phe Asp Glu Met Glu Gln Ser Leu Leu Ile 800 805 810
 Leu Ser Glu Thr Lys Ala Ser Leu Val Ser Thr Met Ser Leu Trp 815 820 825
 Lys Gln Gln Met Tyr Thr Ile Ala Lys Phe His Phe Thr Leu 830 835 840
 Lys Arg Glu Ser Lys Ser Val Arg Ser Val Leu Leu Leu Leu 845 850 855
 Ile Phe Phe Thr Val Gln Ile Phe Met Phe Leu Val His His Ser 860 865 870
 Phe Lys Asn Ala Val Val Pro Ile Lys Leu Val Pro Asp Leu Tyr 875 880 885
 Phe Leu Lys Pro Gly Asp Lys Pro His Lys Tyr Lys Thr Ser Leu 890 895 900
 Leu Leu Gln Asn Ser Ala Asp Ser Asp Ile Ser Asp Leu Ile Ser 905 910 915
 Phe Phe Thr Ser Gln Asn Ile Met Val Thr Met Ile Asn Asp Ser 920 925 930
 Asp Tyr Val Ser Val Ala Pro His Ser Ala Ala Leu Asn Val Val 935 940 945
 His Ser Glu Lys Asp Tyr Val Phe Ala Ala Val Phe Asn Ser Thr 950 955 960
 Met Val Tyr Ser Leu Pro Ile Leu Val Asn Ile Ile Ser Asn Tyr 965 970 975
 Tyr Leu Tyr His Leu Asn Val Thr Glu Thr Ile Gln Ile Trp Ser 980 985 990
 Thr Pro Phe Phe Gln Glu Ile Thr Asp Ile Val Phe Lys Ile Glu 995 1000 1005
 Leu Tyr Phe Gln Ala Ala Leu Leu Gly Ile Ile Val Thr Ala Met 1010 1015 1020
 Pro Pro Tyr Phe Ala Met Glu Asn Ala Glu Asn His Lys Ile Lys 1025 1030 1035
 Ala Tyr Thr Gln Leu Lys Leu Ser Gly Leu Leu Pro Ser Ala Tyr 1040 1045 1050
 Trp Ile Gly Gln Ala Val Val Asp Ile Pro Leu Phe Phe Ile Ile 1055 1060 1065
 Leu Ile Leu Met Leu Gly Ser Leu Leu Ala Phe His Tyr Gly Leu 1070 1075 1080
 Tyr Phe Tyr Thr Val Lys Phe Leu Ala Val Val Phe Cys Leu Ile 1085 1090 1095
 Gly Tyr Val Pro Ser Val Ile Leu Phe Thr Tyr Ile Ala Ser Phe 1100 1105 1110
 Thr Phe Lys Lys Ile Leu Asn Thr Lys Glu Phe Thr Ser Phe Ile 1115 1120 1125
 Tyr Ser Val Ala Ala Leu Ala Cys Ile Ala Ile Thr Glu Ile Thr 1130 1135 1140
 Phe Phe Met Gly Tyr Thr Ile Ala Thr Ile Leu His Tyr Ala Phe 1145 1150 1155
 Cys Ile Ile Pro Ile Tyr Pro Leu Leu Gly Cys Leu Ile Ser 1160 1165 1170
 Phe Ile Lys Ile Ser Trp Lys Asn Val Arg Lys Asn Val Asp Thr 1175 1180 1185
 Tyr Asn Pro Trp Asp Arg Leu Ser Val Ala Val Ile Ser Pro Tyr 1190 1195 1200
 Leu Gln Cys Val Leu Trp Ile Phe Leu Leu Gln Tyr Tyr Glu Lys 1205 1210 1215
 Lys Tyr Gly Gly Arg Ser Ile Arg Lys Asp Pro Phe Phe Arg Asn 1220 1225 1230
 Leu Ser Thr Lys Ser Lys Asn Arg Lys Leu Pro Glu Pro Pro Asp 1235 1240 1245
 Asn Glu Asp Glu Asp Glu Asp Val Lys Ala Glu Arg Leu Lys Val 1250 1255 1260
 1265 1270 1275

Lys Glu Leu Met Gly Cys Gln Cys Cys Glu Glu Lys Pro Ser Ile 1280 1285 1290
 Met Val Ser Asn Leu His Lys Glu Tyr Asp Asp Lys Lys Asp Phe 1295 1300 1305
 Leu Leu Ser Arg Lys Val Lys Lys Val Ala Thr Lys Tyr Ile Ser 1310 1315 1320
 Phe Cys Val Lys Lys Gly Glu Ile Leu Gly Leu Leu Gly Pro Asn 1325 1330 1335
 Gly Ala Gly Lys Ser Thr Ile Ile Asn Ile Leu Val Gly Asp Ile 1340 1345 1350
 Glu Pro Thr Ser Gly Gln Val Phe Leu Gly Asp Tyr Ser Ser Glu 1355 1360 1365
 Thr Ser Glu Asp Asp Ser Leu Lys Cys Met Gly Tyr Cys Pro 1370 1375 1380
 Gln Ile Asn Pro Leu Trp Pro Asp Thr Thr Leu Gln Glu His Phe 1385 1390 1395
 Glu Ile Tyr Gly Ala Val Lys Gly Met Ser Ala Ser Asp Met Lys 1400 1405 1410
 Glu Val Ile Ser Arg Ile Thr His Ala Leu Asp Leu Lys Glu His 1415 1420 1425
 Leu Gln Lys Thr Val Lys Lys Leu Pro Ala Gly Ile Lys Arg Lys 1430 1435 1440
 Leu Cys Phe Ala Leu Ser Met Leu Gly Asn Pro Gln Ile Thr Leu 1445 1450 1455
 Leu Asp Glu Pro Ser Thr Gly Met Asp Pro Lys Ala Lys Gln His 1460 1465 1470
 Met Trp Arg Ala Ile Arg Thr Ala Phe Lys Asn Arg Lys Arg Ala 1475 1480 1485
 Ala Ile Leu Thr Thr His Tyr Met Glu Glu Ala Glu Ala Val Cys 1490 1495 1500
 Asp Arg Val Ala Ile Met Val Ser Gly Gln Leu Arg Cys Ile Gly 1505 1510 1515
 Thr Val Gln His Leu Lys Ser Lys Phe Gly Lys Gly Tyr Phe Leu 1520 1525 1530
 Glu Ile Lys Leu Lys Asp Trp Ile Glu Asn Leu Glu Val Asp Arg 1535 1540 1545
 Leu Gln Arg Glu Ile Gln Tyr Ile Phe Pro Asn Ala Ser Arg Gln 1550 1555 1560
 Glu Ser Phe Ser Ser Ile Leu Ala Tyr Lys Ile Pro Lys Glu Asp 1565 1570 1575
 Val Gln Ser Leu Ser Gln Ser Phe Phe Lys Leu Glu Glu Ala Lys 1580 1585 1590
 His Ala Phe Ala Ile Glu Glu Tyr Ser Phe Ser Gln Ala Thr Leu 1595 1600 1605
 Glu Gln Val Phe Val Glu Leu Thr Lys Glu Gln Glu Glu Asp 1610 1615 1620
 Asn Ser Cys Gly Thr Leu Asn Ser Thr Leu Trp Trp Glu Arg Thr 1625 1630 1635
 Gln Glu Asp Arg Val Val Phe 1640 1645

<210> 4
 <211> 659
 <212> PRT
 <213> Homo sapiens
 <220>
 <221> misc_feature
 <223> Incyte ID No: 3105257CD1
 <400> 4
 Met Gly Arg Gly Ala Gly Ala Ala Leu Glu Arg Trp Ser Arg Ala 1
 5 10 15

Pro Leu Glu Glu Leu Leu Pro Gly Arg Gly Ser Gly Arg Leu Gly
 20 25 30
 Gly Pro Arg Gly Pro Arg Thr Ala Pro Gly Ala Val Gly Leu Gly
 35 40 45
 Pro Ala Ala Ala Gly Glu Glu Ala Trp Arg Arg Gly Arg Ala Ala
 50 55 60
 Pro Ser Arg Asp Asp Glu Arg Leu Arg Pro Met Ala Pro Gly Leu
 65 70 75
 Ser Glu Ala Gly Lys Leu Leu Gly Leu Glu Tyr Pro Glu Arg Glu
 80 85 90
 Arg Leu Ala Ala Val Gly Phe Leu Thr Met Ser Gly Val Ile
 95 100 105
 Ser Met Ser Ala Pro Phe Leu Gly Lys Ile Ile Asp Ala Ile
 110 115 120
 Tyr Thr Asn Pro Thr Val Asp Tyr Ser Asp Asn Leu Thr Arg Leu
 125 130 135
 Cys Leu Gly Leu Ser Ala Val Phe Leu Cys Gly Ala Ala Asn
 140 145 150
 Ala Ile Arg Val Tyr Leu Met Glu Thr Ser Gly Glu Arg Ile Val
 155 160 165
 Asn Arg Leu Arg Thr Ser Leu Phe Ser Ile Leu Arg Glu Glu
 170 175 180
 Val Ala Phe Phe Asp Lys Thr Arg Thr Gly Glu Leu Ile Asn Arg
 185 190 195
 Leu Ser Ser Asp Thr Ala Leu Leu Gly Arg Ser Val Thr Glu Asn
 200 205 210
 Leu Ser Asp Gly Leu Arg Ala Gly Ala Glu Ala Ser Val Gly Ile
 215 220 225
 Ser Met Met Phe Val Ser Pro Asn Leu Ala Thr Phe Val Leu
 230 235 240
 Ser Val Val Pro Pro Val Ser Ile Ile Ala Val Ile Tyr Gly Arg
 245 250 255
 Tyr Leu Arg Lys Leu Thr Lys Val Thr Glu Asp Ser Leu Ala Glu
 260 265 270
 Ala Thr Glu Leu Ala Glu Glu Arg Ile Gly Asn Val Arg Thr Val
 275 280 285
 Arg Ala Phe Gly Lys Glu Met Thr Glu Ile Glu Lys Tyr Ala Ser
 290 295 300
 Lys Val Asp His Val Met Glu Leu Ala Arg Lys Glu Ala Phe Ala
 305 310 315
 Arg Ala Gly Phe Phe Gly Ala Thr Gly Leu Ser Gly Asn Leu Ile
 320 325 330
 Val Leu Ser Val Leu Tyr Lys Gly Gly Leu Leu Met Gly Ser Ala
 335 340 345
 His Met Thr Val Gly Glu Leu Ser Ser Phe Leu Met Tyr Ala Phe
 350 355 360
 Trp Val Gly Ile Ser Ile Gly Gly Leu Ser Ser Phe Tyr Ser Glu
 365 370 375
 Leu Met Lys Gly Leu Gly Ala Gly Arg Leu Trp Glu Leu Leu
 380 385 390
 Glu Arg Glu Pro Lys Leu Pro Phe Asn Glu Gly Val Ile Leu Asn
 395 400 405
 Glu Lys Ser Phe Glu Gly Ala Leu Glu Phe Lys Asn Val His Phe
 410 415 420
 Ala Tyr Pro Ala Arg Pro Glu Val Pro Ile Phe Glu Asp Phe Ser
 425 430 435
 Leu Ser Ile Pro Ser Gly Ser Val Thr Ala Leu Val Gly Pro Ser
 440 445 450
 Gly Ser Gly Lys Ser Thr Val Leu Ser Leu Leu Arg Leu Tyr
 455 460 465
 Asp Pro Ala Ser Gly Thr Ile Ser Leu Asp Gly His Asp Ile Arg
 470 475 480
 Glu Leu Asn Pro Val Trp Leu Arg Ser Lys Ile Gly Thr Val Ser

8/85

Glu Glu Pro Ile Leu Phe Ser Cys Ser Ile Ala Glu Asn Ile Ala
 485 490 495
 Tyr Gly Ala Asp Asp Pro Ser Ser Val Thr Ala Glu Glu Ile Glu
 500 505 510
 Arg Val Ala Glu Val Ala Asn Thr Val Ala Phe Ile Arg Asn Phe
 515 520 525
 Pro Glu Gly Phe Asn Thr Val Val Gly Glu Lys Gly Val Leu Leu
 530 535 540
 Ser Gly Gly Glu Lys Glu Arg Ile Ala Ile Ala Arg Ala Leu Leu
 545 550 555
 Lys Asn Pro Lys Ile Leu Leu Leu Asp Glu Ala Thr Ser Ala Leu
 560 565 570
 Asp Ala Glu Asn Glu Tyr Leu Val Glu Glu Ala Leu Asp Arg Leu
 575 580 585
 Met Asp Gly Arg Thr Val Leu Val Ile Ala His Arg Leu Ser Thr
 590 595 600
 Ile Lys Asn Ala Asn Met Val Ala Val Leu Asp Glu Gly Lys Ile
 605 610 615
 Thr Glu Tyr Gly Lys His Glu Glu Leu Leu Ser Lys Pro Asn Gly
 620 625 630
 Ile Tyr Arg Lys Leu Met Asn Lys Glu Ser Phe Ile Ser Ala
 635 640 645
 650 655

<210> 5
 <211> 1592
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3200979CD1

<400> 5
 Met Val Lys Lys Glu Ile Ser Val Arg Glu Glu Ile Glu Ala Leu
 1 5 10 15
 Leu Tyr Lys Asn Phe Leu Lys Lys Trp Arg Ile Lys Arg Glu Phe
 20 25 30
 Ile Gly Leu Tyr Leu Cys Ile Phe Ser Glu His Phe Arg Ala Thr
 35 40 45
 Arg Phe Pro Glu Glu Pro Pro Lys Val Leu Gly Ser Val Asp Glu
 50 55 60
 Phe Asn Asp Ser Gly Leu Val Val Ala Tyr Thr Pro Val Ser Asn
 65 70 75
 Ile Thr Glu Arg Ile Met Asn Lys Met Ala Leu Ala Ser Phe Met
 80 85 90
 Lys Gly Arg Thr Val Ile Gly Thr Pro Asp Glu Glu Thr Met Asp
 95 100 105
 Ile Glu Leu Pro Lys Lys Tyr His Glu Met Val Gly Val Ile Phe
 110 115 120
 Ser Asp Thr Phe Ser Tyr Arg Leu Lys Phe Asn Trp Gly Tyr Arg
 125 130 135
 Ile Pro Val Ile Lys Glu His Ser Glu Tyr Thr Glu His Cys Trp
 140 145 150
 Ala Met His Gly Glu Ile Phe Cys Tyr Leu Ala Lys Tyr Trp Leu
 155 160 165
 Lys Gly Phe Val Ala Phe Glu Ala Ala Ile Asn Ala Ala Ile Ile
 170 175 180
 Glu Val Thr Thr Asn His Ser Val Met Glu Glu Leu Thr Ser Val
 185 190 195
 Ile Gly Ile Asn Met Lys Ile Pro Pro Ile Ser Lys Gly Glu
 200 205 210
 Ile Met Asn Glu Trp Phe His Phe Thr Cys Leu Val Ser Phe Ser

9/85

Ser Phe Ile Tyr Phe Ala Ser Leu Asn Val Ala Arg Glu Arg Gly 225
230 235
Lys Phe Lys Lys Leu Met Thr Val Met Gly Leu Arg Glu Ser Ala 240
245
Phe Trp Leu Ser Trp Gly Leu Thr Tyr Ile Cys Phe Ile Phe Ile 255
260 265
Met Ser Ile Phe Met Ala Leu Val Ile Thr Ser Ile Pro Ile Val 270
275
Phe His Thr Gly Phe Met Val Ile Phe Thr Leu Tyr Ser Leu Tyr 285
290 295
Gly Leu Ser Leu Ile Ala Leu Ala Phe Leu Met Ser Val Leu Ile 300
305
Arg Lys Pro Met Leu Ala Gly Leu Ala Gly Phe Leu Phe Thr Val 315
320 325
Phe Trp Gly Cys Leu Gly Phe Thr Val Leu Tyr Arg Gln Leu Pro 330
335 340
Leu Ser Leu Gly Trp Val Leu Ser Leu Ser Pro Phe Ala Phe 345
350 355
Thr Ala Gly Met Ala Gln Ile Thr His Leu Asp Asn Tyr Leu Ser 360
365 370
Gly Val Ile Phe Pro Asp Pro Ser Gly Asp Ser Tyr Lys Met Ile 375
380 385
Ala Thr Phe Phe Ile Leu Ala Phe Asp Thr Leu Phe Tyr Leu Ile 390
395 400
Phe Thr Leu Tyr Phe Glu Arg Val Leu Pro Asp Lys Asp Gly His 405
410 415
Gly Asp Ser Pro Leu Phe Phe Leu Lys Ser Ser Phe Trp Ser Lys 420
425 430
His Gln Asn Thr His His Glu Ile Phe Glu Asn Glu Ile Asn Pro 435
440 445
Glu His Ser Ser Asp Asp Ser Phe Glu Pro Val Ser Pro Glu Phe 450
455 460
His Gly Lys Glu Ala Ile Arg Ile Arg Asn Val Ile Lys Glu Tyr 465
470 475
Asn Gly Lys Thr Gly Lys Val Glu Ala Leu Gln Gly Ile Phe Phe 480
485 490
Asp Ile Tyr Glu Gly Gln Ile Thr Ala Ile Leu Gly His Asn Gly 495
500 505
Ala Gly Lys Ser Thr Leu Leu Asn Ile Leu Ser Gly Leu Ser Val 510
515 520
Ser Thr Glu Gly Ser Ala Thr Ile Tyr Asn Thr Gln Leu Ser Glu 525
530 535
Ile Thr Asp Met Glu Glu Ile Arg Lys Asn Ile Gly Phe Cys Pro 540
545 550
Gln Phe Asn Phe Gln Phe Asp Phe Leu Thr Val Arg Glu Asn Leu 555
560 565
Arg Val Phe Ala Lys Ile Lys Gly Ile Gln Pro Lys Glu Val Glu 570
575 580
Gln Glu Val Leu Leu Leu Asp Glu Pro Thr Ala Gly Leu Asp Pro 585
590 595
Phe Ser Arg His Arg Val Trp Ser Leu Leu Lys Glu His Lys Val 600
605 610
Asp Arg Leu Ile Leu Phe Ser Thr Gln Phe Met Asp Glu Ala Asp 615
620 625
Ile Leu Ala Asp Arg Lys Val Phe Leu Ser Asn Gly Lys Leu Lys 630
635 640
Cys Ala Gly Ser Ser Leu Phe Leu Lys Arg Lys Trp Gly Ile Gly 645
650 655
Tyr His Leu Ser Leu His Arg Asn Glu Met Cys Asp Thr Glu Lys 660
665 670
Ile Thr Ser Leu Ile Lys Gln His Ile Pro Asp Ala Lys Leu Thr 675
680 685

Thr Glu Ser Glu Glu Lys Leu Val Tyr Ser Leu Pro Leu Glu Lys 700
705
Thr Asn Lys Phe Pro Asp Leu Tyr Ser Asp Leu Asp Lys Cys Ser 715
720
Asp Gln Gly Ile Arg Asn Tyr Ala Val Ser Val Thr Ser Leu Asn 725
730
Glu Val Phe Leu Asn Leu Glu Gly Lys Ser Ala Ile Asp Glu Pro 735
740 745
Asp Phe Asp Ile Gly Lys Gln Glu Lys Ile His Val Thr Arg Asn 750
755
Thr Gly Asp Glu Ser Glu Met Glu Gln Val Leu Cys Ser Leu Pro 760
770 775
Glu Thr Arg Lys Ala Val Ser Ser Ala Ala Leu Trp Arg Arg Gln 780
785
Ile Tyr Ala Val Ala Thr Leu Arg Phe Leu Lys Leu Arg Arg Glu 790
800 805
Arg Arg Ala Leu Leu Cys Leu Leu Val Leu Gly Ile Ala Phe 810
815
Ile Pro Ile Ile Leu Glu Lys Ile Met Tyr Lys Val Thr Arg Glu 820
830 835
Thr His Cys Trp Glu Phe Ser Pro Ser Met Tyr Phe Leu Ser Leu 840
845 850
Glu Gln Ile Pro Lys Thr Pro Leu Thr Ser Leu Leu Ile Val Asn 855
860 865
Asn Thr Gly Ser Asn Ile Glu Asp Leu Val His Ser Leu Lys Cys 870
875 880
Gln Asp Ile Val Leu Glu Ile Asp Asp Phe Arg Asn Arg Asn Gly 885
890 895
Ser Asp Asp Pro Ser Tyr Asn Gly Ala Ile Ile Val Ser Gly Asp 900
905 910
Gln Lys Asp Tyr Arg Phe Ser Val Ala Cys Asn Thr Lys Lys Ser 915
920 925
Asn Cys Phe Pro Val Leu Met Gly Ile Val Ser Asn Ala Leu Ile 930
935 940
Gly Ile Phe Asn Phe Thr Glu Leu Ile Gln Met Glu Ser Thr Ser 945
950 955
Phe Phe Arg Asp Asp Ile Val Leu Asp Leu Gly Phe Ile Asp Gly 960
965 970
Ser Ile Phe Leu Leu Leu Ile Thr Asn Cys Ile Ser Pro Tyr Ile 975
980 985
Gly Ile Ser Ser Ile Ser Asp Tyr Lys Ile Pro Ser Ser Ile Pro 990
1000
Ser Ile Leu Cys Gln Lys Asn Val Gln Ser Gln Leu Trp Ile Ser 1005
1010 1015
Gly Leu Trp Pro Ser Ala Tyr Trp Cys Gly Gln Ala Leu Val Asp 1020
1025 1030
Ile Pro Leu His Phe Leu Ile Leu Leu Ser Ile His Leu Ile Tyr 1035
1040 1045
Tyr Phe Ser Phe Leu Gly Phe Gln Leu Pro Trp Glu Leu Met Phe 1050
1055 1060
Val Leu Val Val Cys Ile Ile Gly Cys Ala Ala Ser Leu Ile Phe 1065
1070 1075
Leu Met Tyr Val Leu Ser Phe Phe Ile Phe Cys Lys Trp Arg Lys Asn 1080
1085 1090
Asn Gly Phe Trp Ser Phe Gly Phe Phe Ile Val Leu Ile Cys Val 1095
1100 1105
Ser Thr Ile Leu Val Ser Thr Lys Tyr Glu Lys Pro Asn Leu Ile 1110
1115 1120
Leu Cys Met Ile Phe Ile Pro Ser Phe Thr Phe Leu Asp Met Ser 1125
1130 1135
Leu Leu Ile Gln Leu Asn Phe Met Tyr Met Arg Asn Leu Asp Ser 1140
1145 1150
Leu Asp Asn Arg Ile Asn Glu Val Asn Lys Thr Ile Leu Leu Thr 1155

1160 1165 1170
 Asn Leu Ile Pro Tyr Leu Gln Ser Val Ile Phe Leu Phe Val Ile
 1175 1180 1185
 Arg Cys Leu Glu Met Lys Tyr Gly Asn Gln Ile Met Asn Lys Asp
 1190 1195 1200
 Pro Val Phe Arg Ile Ser Pro Arg Ser Arg Gly Thr His Thr Asn
 1205 1210 1215
 Pro Glu Glu Pro Glu Glu Asp Val Gln Ala Glu Arg Val Gln Ala
 1220 1225 1230
 Ala Asn Ala Leu Thr Thr Pro Asn Leu Glu Glu Pro Val Ile
 1235 1240 1245
 Thr Ala Ser Cys Leu His Lys Glu Tyr Tyr Glu Thr Lys Lys Ser
 1250 1255 1260
 Cys Phe Ser Thr Thr Lys Lys Lys Ala Ala Ile Arg Asn Val Ser
 1265 1270 1275
 Phe Cys Val Lys Lys Gly Glu Val Leu Gly Leu Glu His Asn
 1280 1285 1290
 Gly Ala Gly Lys Ser Thr Ser Ile Lys Met Ile Thr Gly Cys Thr
 1295 1300 1305
 Val Pro Thr Ala Gly Val Val Val Leu Gln Gly Asn Arg Ala Ser
 1310 1315 1320
 Val Arg Gln Gln Arg Asn Asn Ser Leu Lys Phe Leu Gly Tyr Cys
 1325 1330 1335
 Pro Gln Glu Asn Ser Leu Tyr Pro Lys Leu Thr Met Lys Glu His
 1340 1345 1350
 Leu Glu Leu Tyr Ala Ala Val Lys Gly Leu Gly Lys Asp Ala
 1355 1360 1365
 Ala Leu Ser Ile Ser Arg Leu Val Glu Ala Leu Lys Leu Gln Glu
 1370 1375 1380
 Gln Leu Lys Ala Pro Val Lys Thr Leu Ser Glu Gly Ile Lys Arg
 1385 1390 1395
 Lys Leu Cys Phe Val Leu Ser Ile Leu Gly Asn Pro Ser Val Val
 1400 1405 1410
 Leu Leu Asp Glu Pro Phe Thr Gly Met Asp Pro Glu Gly Gln Gln
 1415 1420 1425
 Gln Met Trp Gln Ile Leu Gln Ala Thr Ile Lys Asn Gln Gln Arg
 1430 1435 1440
 Gly Thr Leu Leu Thr Thr His Tyr Met Ser Glu Ala Lys Ser Leu
 1445 1450 1455
 Cys Asp Arg Val Ala Ile Met Val Ser Gly Thr Leu Arg Cys Ile
 1460 1465 1470
 Gly Ser Ile Gln His Leu Lys Asn Lys Phe Gly Lys Asp Tyr Leu
 1475 1480 1485
 Leu Glu Ile Lys Met Lys Glu Pro Thr Gln Val Glu Ala Leu His
 1490 1495 1500
 Thr Glu Ile Leu Lys Leu Phe Pro Gln Ala Ala Trp Gln Glu Arg
 1505 1510 1515
 Tyr Ser Ser Leu Met Ala Tyr Lys Leu Pro Val Glu Asp Val His
 1520 1525 1530
 Pro Leu Ser Arg Ala Phe Phe Lys Leu Glu Ala Met Lys Gln Thr
 1535 1540 1545
 Phe Asn Leu Glu Glu Tyr Ser Leu Ser Gln Ala Thr Leu Glu Gln
 1550 1555 1560
 Val Phe Leu Glu Leu Cys Lys Glu Gln Glu Leu Gly Asn Val Asp
 1565 1570 1575
 Asp Lys Ile Asp Thr Thr Val Glu Trp Tyr Leu Leu Pro Gln Gln
 1580 1585 1590
 Asp Pro

<210> 6
 <211> 382
 <212> PRT

12/85

<213> Homo sapiens
 <220>
 <221> misc_feature
 <223> Incyte ID No: 6754139CD1
 <400> 6
 Met Asp Glu Arg Asn Gln Val Leu Thr Leu Tyr Leu Trp Ile Arg
 1 5 10 15
 Gln Glu Trp Thr Asp Ala Tyr Leu Arg Trp Asp Pro Asn Ala Tyr
 20 25 30
 Gly Gly Leu Asp Ala Ile Arg Ile Pro Ser Ser Leu Val Trp Arg
 35 40 45
 Pro Asp Ile Val Leu Tyr Asn Lys Ala Asp Ala Gln Pro Pro Gly
 50 55 60
 Ser Ala Ser Thr Asn Val Val Leu Arg His Asp Gly Ala Val Arg
 65 70 75
 Trp Asp Ala Pro Ala Ile Thr Arg Ser Ser Cys Arg Val Asp Val
 80 85 90
 Ala Ala Phe Pro Phe Asp Ala Gln His Cys Gly Leu Thr Phe Gly
 95 100 105
 Ser Trp Thr His Gly Gly His Gln Val Asp Val Arg Pro Arg Gly
 110 115 120
 Ala Ala Ala Ser Leu Ala Asp Phe Val Glu Asn Val Glu Trp Arg
 125 130 135
 Val Leu Gly Met Pro Ala Arg Arg Val Leu Thr Tyr Gly Cys
 140 145 150
 Cys Ser Glu Pro Tyr Pro Asp Val Thr Phe Thr Leu Leu Leu Arg
 155 160 165
 Arg Arg Ala Ala Tyr Val Cys Asn Leu Leu Pro Cys Val
 170 175 180
 Leu Ile Ser Leu Leu Ala Pro Leu Ala Phe His Leu Pro Ala Asp
 185 190 195
 Ser Gly Glu Lys Val Ser Leu Gly Val Thr Val Leu Leu Ala Leu
 200 205 210
 Thr Val Phe Gln Leu Leu Ala Glu Ser Met Pro Pro Ala Glu
 215 220 225
 Ser Val Pro Leu Ile Gly Lys Tyr Tyr Met Ala Thr Met Thr Met
 230 235 240
 Val Thr Phe Ser Thr Ala Leu Thr Ile Leu Ile Met Asn Leu His
 245 250 255
 Tyr Cys Gly Pro Ser Val Arg Pro Val Pro Ala Trp Ala Arg Ala
 260 265 270
 Leu Leu Leu Gly His Leu Ala Arg Gly Leu Cys Val Arg Glu Arg
 275 280 285
 Gly Glu Pro Cys Gly Gln Ser Arg Pro Pro Glu Leu Ser Pro Ser
 290 295 300
 Pro Gln Ser Pro Glu Gly Gly Ala Gly Pro Pro Ala Gly Pro Cys
 305 310 315
 His Glu Pro Arg Cys Leu Cys Arg Gln Glu Ala Leu Leu His His
 320 325 330
 Val Ala Thr Ile Ala Asn Thr Phe Arg Ser His Arg Ala Ala Gln
 335 340 345
 Arg Cys His Glu Asp Trp Lys Arg Leu Ala Arg Val Met Asp Arg
 350 355 360
 Phe Phe Leu Ala Ile Phe Phe Ser Met Ala Leu Val Met Ser Leu
 365 370 375
 Leu Val Leu Val Gln Ala Leu
 380

<210> 7
 <211> 1115
 <212> PRT

13/85

<213> Homo sapiens
<220>
<221> misc_feature
<223> Incyte ID No: 6996659CD1
<400> 7
Met Arg Arg Leu Ser Leu Trp Trp Leu Leu Ser Arg Val Cys Leu 10 15
1 5
Leu Leu Pro Pro Cys Ala Leu Val Leu Ala Gly Val Pro Ser 20 30
Ser Ser Ser His Pro Gln Pro Cys Gln Ile Leu Lys Arg Ile Gly 25 35
His Ala Val Arg Val Gly Ala Val His Leu Gln Pro Trp Thr Thr 40 45
50 55 60
Ala Pro Arg Ala Ala Ser Arg Ala Pro Asp Asp Ser Arg Ala Gly 65 70 75
Ala Gln Arg Asp Glu Pro Glu Pro Gly Thr Arg Arg Ser Pro Ala 80 85 90
Pro Ser Pro Gly Ala Arg Trp Leu Gly Ser Thr Leu His Gly Arg 95 100 105
Gly Pro Pro Gly Ser Arg Lys Pro Gly Glu Gly Ala Arg Ala Glu 110 115 120
Ala Leu Trp Pro Arg Asp Ala Leu Leu Phe Ala Val Asp Asn Leu 125 130 135
Asn Arg Val Glu Gly Leu Leu Pro Tyr Asn Leu Ser Leu Glu Val 140 145 150
Val Met Ala Ile Glu Ala Gly Leu Gly Asp Leu Pro Leu Leu Pro 155 160 165
Phe Ser Ser Pro Ser Ser Pro Trp Ser Ser Asp Pro Phe Ser Phe 170 175 180
Leu Gln Ser Val Cys His Thr Val Val Val Gln Gly Val Ser Ala 185 190 195
Leu Leu Ala Phe Pro Gln Ser Gln Gly Glu Met Met Glu Leu Asp 200 205 210
Leu Val Ser Leu Leu His Ile Pro Val Ile Ser Ile Val Arg 215 220 225
His Glu Phe Pro Arg Glu Ser Gln Asn Pro Leu His Leu Gln Leu 230 235 240
Ser Leu Glu Asn Ser Leu Ser Ser Asp Ala Asp Val Thr Val Ser 245 250 255
Ile Leu Thr Met Asn Asn Trp Tyr Asn Phe Ser Leu Leu Leu Cys 260 265 270
Gln Glu Asp Trp Asn Ile Thr Asp Phe Leu Leu Leu Thr Gln Asn 275 280 285
Asn Ser Lys Phe His Leu Gly Ser Ile Ile Asn Ile Thr Ala Asn 290 295 300
Leu Pro Ser Thr Gln Asp Leu Leu Ser Phe Leu Gln Ile Gln Leu 305 310 315
Glu Ser Ile Lys Asn Ser Thr Pro Thr Val Val Met Phe Gly Cys 320 325 330
Asp Met Glu Ser Ile Arg Arg Ile Phe Glu Ile Thr Thr Gln Phe 335 340 345
Gly Val Met Pro Glu Leu Arg Trp Val Leu Gly Asp Ser Gln 350 355 360
Asn Val Glu Glu Leu Arg Thr Glu Gly Leu Pro Leu Gly Leu Ile 365 370 375
Ala His Gly Lys Thr Thr Gln Ser Val Phe Glu His Tyr Val Gln 380 385 390
Asp Ala Met Glu Leu Val Ala Arg Ala Val Ala Thr Ala Thr Met 395 400 405
Ile Gln Pro Glu Leu Ala Leu Ile Pro Ser Thr Met Asn Cys Met 410 415 420

Glu Val Glu Thr Thr Asn Leu Thr Ser Gly Gln Tyr Leu Ser Arg 425 430 435
Phe Leu Ala Asn Thr Thr Phe Arg Gly Leu Ser Gly Ser Ile Arg 440 445 450
Val Lys Gly Ser Thr Ile Val Ser Ser Glu Asn Asn Phe Phe Ile 455 460 465
Trp Asn Leu Gln His Asp Pro Met Gly Lys Pro Met Trp Thr Arg 470 475 480
Leu Gly Ser Trp Gln Gly Gly Lys Ile Val Met Asp Tyr Gly Ile 485 490 495
Trp Pro Glu Gln Ala Gln Arg His Lys Thr His Phe Gln His Pro 500 505 510
Ser Lys Leu His Leu Arg Val Thr Leu Leu Ile Glu His Pro Phe 515 520 525
Val Phe Thr Arg Glu Val Asp Asp Glu Gly Leu Cys Pro Ala Gly 530 535 540
Gln Leu Cys Leu Asp Pro Met Thr Asn Asp Ser Ser Thr Leu Asp 545 550 555
Ser Leu Phe Ser Ser Leu His Ser Ser Asn Asp Thr Val Pro Ile 560 565 570
Lys Phe Lys Lys Cys Cys Tyr Gly Tyr Cys Ile Asp Leu Leu Glu 575 580 585
Lys Ile Ala Glu Asp Met Asn Phe Asp Phe Asp Leu Tyr Ile Val 590 595 600
Gly Asp Gly Lys Tyr Gly Ala Trp Lys Asn Gly His Trp Thr Gly 605 610 615
Leu Val Gly Asp Leu Leu Arg Gly Thr Ala His Met Ala Val Thr 620 625 630
Ser Phe Ser Ile Asn Thr Ala Arg Ser Gln Val Ile Asp Phe Thr 635 640 645
Ser Pro Phe Phe Ser Thr Ser Leu Gly Ile Leu Val Arg Thr Arg 650 655 660
Asp Thr Ala Ala Pro Ile Gly Ala Phe Met Trp Pro Leu His Trp 665 670 675
Thr Met Trp Leu Gly Ile Phe Val Ala Leu His Ile Thr Ala Val 680 685 690
Phe Leu Thr Leu Tyr Glu Trp Lys Ser Pro Phe Gly Leu Thr Ser 695 700 705
Lys Gly Arg Asn Arg Ser Lys Val Phe Ser Phe Ser Ala Leu 710 715 720
Asn Ile Cys Tyr Ala Leu Leu Phe Gly Arg Thr Val Ala Ile Lys 725 730 735
Pro Pro Lys Cys Trp Thr Gly Arg Phe Leu Met Asn Leu Trp Ala 740 745 750
Ile Phe Cys Met Phe Cys Leu Ser Thr Tyr Thr Ala Asn Leu Ala 755 760 765
Ala Val Met Val Gly Glu Lys Ile Tyr Glu Glu Leu Ser Gly Ile 770 775 780
His Asp Pro Lys Leu His His Pro Ser Gln Gly Phe Arg Phe Gly 785 790 795
Thr Val Arg Glu Ser Ser Ala Glu Asp Tyr Val Arg Gln Ser Phe 800 805 810
Pro Glu Met His Glu Tyr met Arg Arg Tyr Asn Val Pro Ala Thr 815 820 825
Pro Asp Gly Val Glu Tyr Leu Lys Asn Asp Pro Glu Lys Leu Asp 830 835 840
Ala Phe Ile Met Asp Lys Ala Leu Leu Asp Tyr Glu Val Ser Ile 845 850 855
Asp Ala Asp Cys Lys Leu Leu Thr Val Gly Lys Pro Phe Ala Ile 860 865 870
Glu Gly Tyr Gly Ile Gly Leu Pro Pro Asn Ser Pro Leu Thr Ala 875 880 885
Asn Ile Ser Glu Leu Ile Ser Gln Tyr Lys Ser His Gly Phe Met 890 895 900

890 895 900
Asp Met Leu His Asp Lys Trp Tyr Arg Val Val Pro Cys Gly Lys
905 910 915
Arg Ser Phe Ala Val Thr Glu Thr Leu Glu Met Gly Ile Lys His
920 925 930
Phe Ser Gly Leu Phe Val Leu Leu Cys Ile Gly Phe Gly Leu Ser
935 940 945
Ile Leu Thr Thr Ile Gly Glu His Ile Val Tyr Arg Leu Leu Leu
950 955 960
Pro Arg Ile Lys Asn Lys Ser Lys Leu Glu Tyr Trp Leu His Thr
965 970 975
Ser Glu Arg Leu His Arg Ala Ile Asn Thr Ser Phe Ile Glu Glu
980 985 990
Lys Glu Glu His Phe Lys Thr Lys Arg Val Glu Lys Arg Ser Asn
995 1000 1005
Val Gly Pro Arg Glu Leu Thr Val Trp Asn Thr Ser Asn Leu Ser
1010 1015 1020
His Asp Asn Arg Arg Lys Tyr Ile Phe Ser Asp Glu Glu Gly Glu
1025 1030 1035
Asn Glu Leu Gly Ile Arg Ile His Glu Asn Ile Pro Leu Pro Pro
1040 1045 1050
Arg Arg Arg Glu Leu Pro Ala Leu Arg Thr Thr Asn Gly Lys Ala
1055 1060 1065
Asp Ser Leu Asn Val Ser Arg Asn Ser Val Met Glu Glu Lys Ala
1070 1075 1080
Glu Leu Glu Lys Glu Ile Glu Val Ile Arg Glu Glu Leu Glu Leu
1085 1090 1095
Ala Val Ser Arg Lys Thr Glu Leu Glu Tyr Tyr Glu Arg Thr Ser
1100 1105 1110
Arg Thr Cys Glu Ser
1115

<210> 8
<211> 295
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 7472747CD1

<400> 8
Met Pro Ser Ala Gly Leu Cys Ser Cys Trp Gly Gly Arg Val Leu
1 5 10 15
Pro Leu Leu Leu Ala Tyr Val Cys Tyr Leu Leu Leu Gly Ala Thr
20 25 30
Ile Phe Glu Leu Leu Glu Arg Glu Ala Glu Ala Glu Ser Arg Asp
35 40 45
Glu Phe Glu Leu Glu Lys Leu Arg Phe Leu Glu Asn Tyr Thr Cys
50 55 60
Leu Asp Glu Trp Ala Met Glu Glu Phe Val Glu Val Ile Met Glu
65 70 75
Ala Trp Val Lys Gly Val Asn Pro Lys Gly Asn Ser Thr Asn Pro
80 85 90
Ser Asn Trp Asp Phe Gly Ser Ser Phe Phe Ala Gly Thr Val
95 100 105
Val Thr Thr Ile Gly Tyr Gly Asn Leu Ala Pro Ser Thr Glu Ala
110 115 120
Gly Glu Val Phe Cys Val Phe Tyr Ala Leu Leu Gly Ile Pro Leu
125 130 135
Asn Val Ile Phe Leu Asn His Leu Gly Thr Gly Leu Arg Ala His
140 145 150
Leu Ala Ala Ile Glu Arg Trp Glu Asp Arg Pro Arg Arg Ser Glu

16/85

155 160 165
Glu Val Leu Glu Glu Val Leu Gly Leu Ala Leu Phe Leu Thr Leu Gly
170 175 180
Thr Leu Val Ile Leu Ile Phe Pro Pro Pro Met Val Phe Ser His Val
185 190 195
Glu Gly Trp Ser Phe Ser Glu Gly Phe Tyr Phe Ala Phe Ile Thr
200 205 210
Leu Ser Thr Ile Gly Phe Gly Asp Tyr Val Ala Gly Thr Asp Pro
215 220 225
Ser Lys His Tyr Ile Ser Val Tyr Arg Ser Leu Ala Ala Ile Trp
230 235 240
Ile Leu Leu Gly Leu Ala Trp Leu Ala Leu Ile Leu Pro Leu Gly
245 250 255
Pro Leu Leu Leu His Arg Cys Cys Glu Leu Trp Leu Leu Ser Arg
260 265 270
Gly Leu Gly Val Lys Asp Gly Ala Ala Ser Asp Pro Ser Gly Leu
275 280 285
Pro Arg Pro Glu Lys Ile Pro Ile Ser Ala
290 295

<210> 9
<211> 384
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 7474121CD1

<400> 9
Met Glu Val Ser Gly His Pro Glu Ala Arg Arg Cys Cys Pro Glu
1 5 10 15
Ala Leu Gly Lys Leu Phe Pro Gly Leu Cys Phe Leu Cys Phe Leu
20 25 30
Val Thr Tyr Ala Leu Val Gly Ala Val Val Phe Ser Ala Ile Gly
35 40 45
Asp Gly Glu Val Leu Val Ala Ala Asp Asp Gly Glu Phe Glu Lys
50 55 60
Phe Leu Glu Glu Leu Cys Arg Ile Leu Asn Cys Ser Glu Thr Val
65 70 75
Val Glu Asp Arg Lys Glu Asp Leu Glu Glu His Leu Glu Lys Val
80 85 90
Lys Pro Glu Trp Phe Asn Arg Thr Thr His Trp Ser Phe Leu Ser
95 100 105
Ser Leu Phe Phe Cys Cys Thr Val Phe Ser Thr Val Gly Tyr Gly
110 115 120
Tyr Ile Tyr Pro Val Thr Arg Leu Gly Lys Tyr Leu Cys Met Leu
125 130 135
Tyr Ala Leu Phe Gly Ile Pro Leu Met Phe Leu Val Leu Thr Asp
140 145 150
Thr Gly Asp Ile Leu Ala Thr Ile Leu Ser Thr Ser Tyr Asn Arg
155 160 165
Phe Arg Lys Phe Pro Phe Phe Thr Arg Pro Leu Leu Ser Lys Trp
170 175 180
Cys Pro Lys Ser Leu Phe Lys Lys Lys Pro Asp Pro Lys Pro Ala
185 190 195
Asp Glu Ala Val Pro Glu Ile Ile Ile Ser Ala Glu Glu Leu Pro
200 205 210
Gly Pro Lys Leu Gly Thr Cys Pro Ser Arg Pro Ser Cys Ser Met
215 220 225
Glu Leu Phe Glu Arg Ser His Ala Leu Glu Lys Glu Asn Thr Leu
230 235 240
Glu Leu Pro Pro Glu Ala Met Glu Arg Ser Asn Ser Cys Pro Glu

17/85

Leu Val Leu Gly Arg Leu Ser Tyr Ser Ile Ile Ser Asn Leu Asp 255
245 260 265 270
Glu Val Gly Gln Gln Val Glu Arg Leu Asp Ile Pro Leu Pro Ile 285
275 280 285
Ile Ala Leu Ile Val Phe Ala Tyr Ile Ser Cys Ala Ala Ala Ile 300
290 295 300
Leu Pro Phe Trp Glu Thr Gln Leu Asp Phe Glu Asn Ala Phe Tyr 315
305 310 315
Phe Cys Phe Val Thr Leu Thr Thr Ile Gly Phe Gly Asp Thr Val 330
320 325 330
Leu Glu His Pro Asn Phe Phe Leu Phe Phe Ser Ile Tyr Ile Ile 345
335 340 345
Val Gly Met Glu Ile Val Phe Ile Ala Phe Lys Leu Val Gln Asn 355
350 360
Arg Leu Ile Asp Ile Tyr Lys Asn Val Met Leu Phe Phe Ala Lys 375
365 370
Gly Lys Phe Tyr His Leu Val Lys Lys 380

<210> 10

<211> 769

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7475615CD1

<400> 10

Met Val Ser Pro Lys Met Tyr Leu Ser Thr Glu Ile Arg Asn Thr 15
1 10 15
Phe Arg Leu Pro Ala Pro Gln Thr His Leu Gln Pro Cys Pro Ala 20
20 25 30
Gly Phe Ala His Pro Leu Leu Val Asn Ala Pro Asp Met Ser Gln 40
35 40 45
Pro Arg Pro Arg Tyr Val Val Asp Arg Ala Ala Tyr Ser Leu Thr 50
50 55 60
Leu Phe Asp Asp Glu Phe Glu Lys Lys Asp Arg Thr Tyr Pro Val 65
65 70 75
Gly Glu Lys Leu Arg Asn Ala Phe Arg Cys Ser Ser Ala Lys Ile 80
80 85 90
Lys Ala Val Val Phe Gly Leu Leu Pro Val Leu Ser Trp Leu Pro 95
95 100 105
Lys Tyr Lys Ile Lys Asp Tyr Ile Ile Pro Asp Leu Leu Gly Gly 110
110 115 120
Leu Ser Gly Gly Ser Ile Gln Val Pro Gln Gly Met Ala Phe Ala 125
125 130 135
Leu Leu Ala Asn Leu Pro Ala Val Asn Gly Leu Tyr Ser Ser Phe 140
140 145 150
Phe Pro Leu Leu Thr Tyr Phe Phe Leu Gly Gly Val His Gln Met 155
155 160 165
Val Pro Gly Thr Phe Ala Val Ile Ser Ile Leu Val Gly Asn Ile 170
170 175 180
Cys Leu Gln Leu Ala Pro Glu Ser Lys Phe Gln Val Phe Asn Asn 185
185 190 195
Ala Thr Asn Glu Ser Tyr Val Asp Thr Ala Ala Met Glu Ala Glu 200
200 205 210
Arg Leu His Val Ser Ala Thr Leu Ala Cys Leu Thr Ala Ile Ile 215
215 220 225
Gln Met Gly Leu Gly Phe Met Gln Phe Gly Phe Val Ala Ile Tyr 230
230 235 240
Leu Ser Glu Ser Phe Ile Arg Gly Phe Met Thr Ala Ala Gly Leu 245

Gln Ile Leu Ile Ser Val Leu Lys Tyr Ile Phe Gly Leu Thr Ile 255
245 260 265 270
Pro Ser Tyr Thr Gly Pro Gly Ser Ile Val Phe Thr Phe Ile Asp 285
275 280 285
Ile Cys Lys Asn Leu Pro His Thr Asn Ile Ala Ser Leu Ile Phe 300
290 295 300
Ala Leu Ile Ser Gly Ala Phe Leu Val Leu Val Lys Glu Leu Asn 315
305 310 315
Ala Arg Tyr Met His Lys Ile Arg Phe Pro Ile Pro Thr Glu Met 330
320 325 330
Ile Val Val Val Val Ala Thr Ala Ile Ser Gly Gly Cys Lys Met 345
335 340 345
Pro Lys Lys Tyr His Met Gln Ile Val Gly Glu Ile Gln Arg Gly 355
350 360
Phe Pro Thr Pro Val Ser Pro Val Val Ser Gln Trp Lys Asp Met 375
365 370
Ile Gly Thr Ala Phe Ser Leu Ala Ile Val Ser Tyr Val Ile Asn 385
380 390
Leu Ala Met Gly Arg Thr Leu Ala Asn Lys His Gly Tyr Asp Val 405
395 400 405
Asp Ser Asn Gln Glu Met Ile Ala Leu Gly Cys Ser Asn Phe Phe 420
410 415 420
Gly Ser Phe Phe Lys Ile His Val Ile Cys Cys Ala Leu Ser Val 435
425 430 435
Thr Leu Ala Val Asp Gly Ala Gly Gly Lys Ser Gln Ser Val Leu 445
440 450
Gly Ala Leu Ile Ala Val Asn Leu Lys Asn Ser Leu Lys Gln Leu 465
455 460 465
Thr Asp Pro Tyr Tyr Leu Trp Arg Lys Ser Lys Leu Asp Cys Cys 475
470 480
Ile Trp Val Val Ser Phe Leu Ser Ser Phe Phe Leu Ser Leu Pro 495
485 490 495
Tyr Gly Val Ala Val Gly Val Ala Phe Ser Val Leu Val Val Val 510
500 505
Phe Gln Thr Gln Phe Arg Asn Gly Tyr Ala Leu Ala Gln Val Met 525
515 520 525
Asp Thr Asp Ile Tyr Val Asn Pro Lys Thr Tyr Asn Arg Ala Gln 540
530 535
Asp Ile Gln Gly Ile Lys Ile Ile Thr Tyr Cys Ser Pro Leu Tyr 555
545 550 555
Phe Ala Asn Ser Glu Ile Phe Arg Gln Lys Val Ile Ala Lys Thr 570
560 565
Val Ser Leu Gln Glu Leu Gln Gln Asp Phe Glu Asn Ala Pro Pro 585
575 580 585
Thr Asp Pro Asn Asn Asn Gln Thr Pro Ala Asn Gly Thr Ser Val 595
590 600
Ser Tyr Ile Thr Phe Ser Pro Asp Ser Ser Ser Pro Ala Gln Ser 615
605 610 615
Glu Pro Pro Ala Ser Ala Glu Ala Pro Gly Glu Pro Ser Asp Met 630
620 625 630
Leu Ala Ser Val Pro Pro Phe Val Thr Phe His Thr Leu Ile Leu 645
635 640 645
Asp Met Ser Gly Val Ser Phe Val Asp Leu Met Gly Ile Lys Ala 660
650 655 660
Leu Ala Lys Leu Ser Ser Thr Tyr Gly Lys Ile Gly Val Lys Val 675
665 670 675
Phe Leu Val Asn Ile His Ala Gln Val Tyr Asn Asp Ile Ser His 685
680 690
Gly Gly Val Phe Glu Asp Gly Ser Leu Glu Cys Lys His Val Phe 705
695 700
Pro Ser Ile His Asp Ala Val Leu Phe Ala Gln Ala Asn Ala Arg 715
710 715 720

Asp Val Thr Pro Gly His Asn Phe Gln Gly Ala Pro Gly Asp Ala
 725 730 735
 Gln Leu Ser Leu Tyr Asp Ser Gln Glu Asp Ile Arg Ser Tyr Trp
 740 745 750
 Asp Leu Gln Gln Met Phe Gly Ser Met Phe His Ala Gln Thr
 755 760 765
 Leu Thr Ala Leu

<210> 11
 <211> 882
 <212> PRT
 <213> Homo sapiens

<220>
 <221> mIsoc.feature
 <223> Incyte ID No: 7475656CD1

<400> 11
 Met Gln Gly Gly Lys Pro Asn Ser Ser Ser Asn Ser Arg Asp
 1 5 10 15
 Asp Gly Asn Ser Val Phe Pro Ala Lys Ala Ser Ala Pro Gly Ala
 20 25 30
 Gly Pro Ala Ala Gln Lys Arg Leu Gly Thr Pro Pro Gly Gly
 35 40 45
 Gly Gly Ala Gly Ala Lys Gln His Gly Asn Ser Val Cys Phe Lys
 50 55 60
 Val Asp Gly Gly Gly Gln Glu Pro Ala Gly Gly Phe Gln Asp
 65 70 75
 Ala Gln Gly Pro Arg Arg Gln Tyr Gly Phe Met Gln Arg Gln Phe
 80 85 90
 Thr Ser Met Leu Gln Pro Gly Val Asn Lys Phe Ser Leu Arg Met
 95 100 105
 Phe Gly Ser Gln Lys Ala Val Gln Lys Gln Gln Arg Val Lys
 110 115 120
 Thr Ala Gly Phe Tyr Ile Ile His Pro Tyr Ser Asp Phe Arg Phe
 125 130 135
 Tyr Trp Asp Leu Ile Met Leu Ile Met Met Val Gly Asn Leu Val
 140 145 150
 Ile Ile Pro Val Gly Ile Thr Phe Phe Thr Gln Thr Thr Thr
 155 160 165
 Pro Trp Ile Ile Phe Asn Val Ala Ser Asp Thr Val Phe Leu Leu
 170 175 180
 Asp Leu Ile Met Asn Phe Arg Thr Gly Thr Val Asn Gln Asp Ser
 185 190 195
 Ser Gln Ile Ile Leu Asp Pro Lys Val Ile Lys Met Asn Tyr Leu
 200 205 210
 Lys Ser Trp Phe Val Val Asp Phe Ile Ser Ser Ile Pro Val Asp
 215 220 225
 Tyr Ile Phe Leu Ile Val Gln Lys Gly Met Asp Ser Gln Val Tyr
 230 235 240
 Lys Thr Ala Arg Ala Leu Arg Ile Val Arg Phe Thr Lys Ile Leu
 245 250 255
 Ser Leu Leu Arg Leu Leu Arg Leu Ser Arg Leu Ile Arg Tyr Ile
 260 265 270
 His Gln Trp Gln Glu Ile Phe His Met Thr Tyr Asp Leu Ala Ser
 275 280 285
 Ala Val Val Arg Ile Phe Asn Leu Ile Gly Met Met Leu Leu Leu
 290 295 300
 Cys His Trp Asp Gly Cys Leu Gln Phe Leu Val Pro Leu Leu Gln
 305 310 315
 Asp Phe Pro Pro Asp Cys Trp Val Ser Leu Asn Gln Met Val Asn
 320 325 330

20/85

Asp Ser Trp Gly Lys Gln Tyr Ser Tyr Ala Leu Phe Lys Ala Met
 335 340 345
 Ser His Met Leu Cys Ile Gly Tyr Gly Ala Gln Ala Pro Val Ser
 350 355 360
 Met Ser Asp Leu Trp Ile Thr Met Leu Ser Met Ile Val Gly Ala
 365 370 375
 Thr Cys Tyr Ala Met Phe Val Gly His Ala Thr Ala Leu Ile Gln
 380 385 390
 Ser Leu Asp Ser Ser Arg Arg Gln Tyr Gln Lys Tyr Lys Gln
 395 400 405
 Val Gln Gln Tyr Met Ser Phe His Lys Leu Pro Ala Asp Met Arg
 410 415 420
 Gln Lys Ile His Asp Tyr Tyr Gln His Arg Tyr Gln Gly Lys Ile
 425 430 435
 Phe Asp Gln Glu Asn Ile Leu Asn Gln Leu Asn Asp Pro Leu Arg
 440 445 450
 Gln Gln Ile Val Asn Phe Asn Cys Arg Lys Leu Val Ala Thr Met
 455 460 465
 Pro Leu Phe Ala Asn Ala Asp Pro Asn Phe Val Thr Ala Met Leu
 470 475 480
 Ser Lys Leu Arg Arg Phe Gln Val Phe Gln Pro Gly Asp Tyr Ile Ile
 485 490 495
 Arg Gln Gly Ala Val Gly Lys Lys Met Tyr Phe Ile Gln His Gly
 500 505 510
 Val Ala Gly Val Ile Thr Lys Ser Ser Lys Gln Met Lys Leu Thr
 515 520 525
 Asp Gly Ser Tyr Phe Gly Gln Ile Cys Leu Leu Thr Lys Gly Arg
 530 535 540
 Arg Thr Ala Ser Val Arg Ala Asp Thr Tyr Cys Arg Leu Tyr Ser
 545 550 555
 Leu Ser Val Asp Asn Phe Asn Gln Val Leu Gln Glu Tyr Pro Met
 560 565 570
 Met Arg Arg Ala Phe Gln Thr Val Ala Ile Asp Arg Leu Asp Arg
 575 580 585
 Ile Gly Lys Lys Asn Ser Ile Leu Leu Gln Lys Phe Gln Lys Asp
 590 595 600
 Leu Asn Thr Gly Val Phe Asn Asn Gln Gln Asn Gln Ile Leu Lys
 605 610 615
 Gln Ile Val Lys His Asp Arg Gln Met Val Gln Ala Ile Ala Pro
 620 625 630
 Ile Asn Tyr Pro Gln Met Thr Thr Leu Asn Ser Thr Ser Ser Thr
 635 640 645
 Thr Thr Pro Thr Ser Arg Met Arg Thr Gln Ser Pro Pro Val Tyr
 650 655 660
 Thr Ala Thr Ser Leu Ser His Ser Asn Leu His Ser Pro Ser Pro
 665 670 675
 Ser Thr Gln Thr Pro Gln Pro Ser Ala Ile Leu Ser Pro Cys Ser
 680 685 690
 Tyr Thr Thr Ala Val Cys Ser Pro Pro Val Gln Ser Pro Leu Ala
 695 700 705
 Ala Arg Thr Phe His Tyr Ala Ser Pro Thr Ala Ser Gln Leu Ser
 710 715 720
 Leu Met Gln Gln Gln Pro Gln Gln Gln Val Gln Gln Ser Gln Pro
 725 730 735
 Pro Gln Thr Gln Pro Gln Gln Pro Ser Pro Gln Pro Gln Thr Thr
 740 745 750
 Gly Ser Ser Thr Pro Lys Asn Gln Val His Lys Ser Thr Gln Ala
 755 760 765
 Leu His Asn Thr Asn Leu Thr Arg Gln Val Arg Pro Leu Ser Ala
 770 775 780
 Ser Gln Pro Ser Leu Pro His Gln Val Ser Thr Leu Ile Ser Arg
 785 790 795
 Pro His Pro Thr Val Gly Gln Ser Leu Ala Ser Ile Pro Gln Pro

21/85

Val Thr Ala Val Pro Gly Thr Gly Leu Gln Ala Gly Gly Arg Ser 800 805 810
815 820 825
Thr Val Pro Gln Arg Val Thr Leu Phe Arg Gln Met Ser Ser Gly 830 835 840
Ala Ile Pro Pro Asn Arg Gly Val Pro Pro Ala Pro Pro Pro 845 850 855
Ala Ala Ala Leu Pro Arg Glu Ser Ser Val Leu Asn Thr Asp 860 865 870
Pro Asp Ala Glu Lys Pro Arg Phe Ala Ser Asn Leu 875 880

<210> 12
<211> 1547
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 7480632CD1

<400> 12
Met Val Lys Lys Glu Ile Ser Val Arg Gln Gln Ile Gln Ala Leu 1
1 5 10 15
Leu Tyr Lys Asn Phe Leu Lys Lys Trp Arg Ile Lys Arg Glu Phe 20 25 30
Leu Glu Glu Trp Thr Ile Thr Leu Phe Leu Glu Gly Leu Tyr Lys Cys 35 40 45
Ile Phe Ser Glu His Phe Arg Ala Thr Arg Phe Pro Glu Gln Pro 50 55 60
Pro Lys Val Leu Gly Ser Val Asp Gln Phe Asn Asp Ser Gly Leu 65 70 75
Val Val Ala Tyr Thr Pro Val Ser Asn Ile Thr Gln Arg Ile Met 80 85 90
Asn Lys Met Ala Leu Ala Ser Phe Met Lys Gly Arg Thr Val Ile 95 100 105
Gly Thr Pro Asp Glu Glu Thr Met Asp Ile Glu Leu Pro Lys Lys 110 115 120
Tyr His Glu Met Val Gly Val Ile Phe Ser Asp Thr Phe Ser Tyr 125 130 135
Arg Leu Lys Phe Asn Trp Gly Tyr Arg Ile Pro Val Ile Lys Glu 140 145 150
His Ser Glu Tyr Thr Gly His Cys Trp Ala Met His Gly Glu Ile 155 160 165
Phe Cys Tyr Leu Ala Lys Tyr Trp Leu Lys Gly Phe Val Ala Phe 170 175 180
Gln Ala Ala Ile Asn Ala Ala Ile Ile Glu Val Thr Thr Asn His 185 190 195
Ser Val Met Glu Glu Leu Thr Ser Val Ile Gly Ile Asn Met Lys 200 205 210
Ile Pro Pro Phe Ile Ser Lys Gly Glu Ile Met Asn Glu Trp Phe 215 220 225
His Phe Thr Cys Leu Val Ser Phe Ser Phe Ile Tyr Phe Ala 230 235 240
Ser Leu Asn Val Ala Arg Glu Arg Gly Lys Phe Lys Lys Leu Met 245 250 255
Thr Val Met Gly Leu Arg Glu Ser Ala Phe Trp Leu Ser Trp Gly 260 265 270
Leu Thr Tyr Ile Cys Phe Ile Phe Ile Met Ser Ile Phe Met Ala 275 280 285
Leu Val Ile Thr Ser Ile Pro Ile Val Phe His Thr Gly Phe Met 290 295 300
Val Ile Phe Thr Leu Tyr Ser Leu Tyr Gly Leu Ser Leu Val Ala

Leu Ala Phe Leu Met Ser Val Leu Ile Arg Lys Pro Met Leu Ala 305 310 315
320 325 330
Gly Leu Ala Gly Phe Leu Phe Thr Val Phe Trp Gly Cys Leu Gly 335 340 345
Phe Thr Val Leu Tyr Arg Gln Leu Pro Leu Ser Leu Gly Trp Val 350 355 360
Leu Ser Leu Leu Ser Pro Phe Ala Phe Thr Ala Gly Met Ala Gln 365 370 375
Ile Thr His Leu Asp Asn Tyr Leu Ser Gly Val Ile Phe Pro Asp 380 385 390
Pro Ser Gly Asp Ser Tyr Lys Met Ile Ala Thr Phe Phe Ile Leu 395 400 405
Ala Phe Asp Thr Leu Phe Tyr Leu Ile Phe Thr Leu Tyr Phe Glu 410 415 420
Arg Val Leu Pro Gly Lys Asp Gly His Gly Asp Ser Pro Leu Phe 425 430 435
Phe Leu Lys Ser Phe Thr Ser Lys His Gln Asn Thr His His 440 445 450
Glu Ile Phe Glu Asn Glu Ile Asn Pro Glu His Ser Ser Asp Asp 455 460 465
Ser Phe Glu Pro Val Ser Pro Glu Phe His Gly Lys Glu Ala Ile 470 475 480
Arg Ile Arg Asn Val Ile Lys Glu Tyr Asn Glu Lys Thr Gly Lys 485 490 495
Val Glu Ala Leu Gln Gly Ile Phe Phe Asp Ile Tyr Glu Gly Gln 500 505 510
Ile Thr Ala Ile Leu Gly His Asn Gly Ala Gly Lys Ser Thr Leu 515 520 525
Leu Asn Ile Leu Ser Gly Leu Ser Val Ser Thr Glu Gly Ser Ala 530 535 540
Thr Ile Tyr Asn Thr Gln Leu Ser Glu Ile Thr Asp Met Glu Glu 545 550 555
Ile Arg Lys Asn Ile Gly Phe Cys Pro Gln Phe Asn Phe Gln Phe 560 565 570
Asp Phe Leu Thr Val Arg Glu Asn Leu Arg Val Phe Ala Lys Ile 575 580 585
Lys Gly Ile Gln Pro Lys Glu Val Glu Glu Val Leu Leu Leu 590 595 600
Asp Glu Pro Thr Ala Gly Leu Asp Pro Phe Ser Arg His Arg Val 605 610 615
Trp Ser Leu Leu Lys Glu His Lys Val Asp Arg Leu Ile Leu Phe 620 625 630
Ser Thr Gln Phe Met Asp Glu Ala Asp Ile Leu Ala Asp Arg Lys 635 640 645
Val Phe Leu Ser Asn Gly Lys Leu Lys Cys Ala Gly Ser Ser Leu 650 655 660
Phe Leu Lys Arg Lys Trp Gly Ile Gly Tyr His Leu Ser Leu His 665 670 675
Arg Asn Glu Met Cys Asp Thr Glu Lys Ile Thr Ser Leu Ile Lys 680 685 690
Gln His Ile Pro Asp Ala Lys Leu Thr Thr Glu Ser Glu Glu Lys 695 700 705
Leu Val Tyr Ser Leu Pro Leu Glu Lys Thr Asn Lys Phe Pro Asp 710 715 720
Leu Tyr Ser Asp Leu Asp Lys Cys Ser Asp Gln Gly Ile Arg Asn 725 730 735
Tyr Ala Val Ser Val Thr Ser Leu Asn Glu Val Phe Leu Asn Leu 740 745 750
Glu Gly Lys Ser Ala Ile Asp Glu Pro Asp Phe Asp Ile Gly Lys 755 760 765
Gln Glu Lys Ile His Val Thr Arg Asn Thr Gly Asp Glu Ser Glu 770 775 780

Met Glu Gln Val Leu Cys Ser Leu Pro Glu Thr Arg Lys Ala Val 1250
 785 790 795
 Ser Ser Ala Ala Leu Trp Arg Arg Gln Ile Tyr Ala Val Ala Thr 1275
 800 805 810
 Leu Arg Phe Leu Leu Arg Arg Glu Arg Arg Ala Leu Leu Cys 1280
 815 820 825
 Leu Leu Leu Val Leu Gly Ile Ala Phe Ile Pro Ile Ile Leu Glu 1290
 830 835 840
 Lys Ile Met Tyr Lys Val Thr Arg Glu Thr His Cys Trp Glu Phe 1300
 845 850 855
 Ser Pro Ser Met Tyr Phe Leu Ser Leu Glu Gln Ile Pro Lys Thr 1310
 860 865 870
 Pro Leu Thr Ser Leu Leu Ile Val Asn Asn Thr Gly Ser Asn Ile 1320
 875 880 885
 Glu Asp Leu Val His Ser Leu Lys Cys Gln Asp Ile Val Leu Glu 1330
 890 895 900
 Ile Asp Asp Phe Arg Asn Arg Asn Gly Ser Asp Asp Pro Ser Tyr 1340
 905 910 915
 Asn Gly Ala Ile Ile Val Ser Gly Asp Gln Lys Asp Tyr Arg Phe 1350
 920 925 930
 Ser Val Ala Cys Asn Thr Lys Lys Leu Asn Cys Phe Pro Val Leu 1360
 935 940 945
 Met Gly Ile Val Ser Asn Ala Leu Met Gly Ile Phe Asn Phe Thr 1370
 950 955 960
 Glu Leu Ile Gln Met Glu Ser Thr Ser Phe Phe Tyr Ile Thr 1380
 965 970 975
 Thr Lys Ser Phe Gln Thr Lys Ile Pro Ser Ser Ile Pro Ser Ile 1390
 980 985 990
 Leu Cys Gln Lys Asn Val Gln Ser Gln Leu Trp Ile Ser Gly Leu 1400
 995 1000 1005
 Trp Pro Ser Ala Tyr Trp Cys Gly Gln Ala Leu Val Asp Ile Pro 1410
 1010 1015 1020
 Leu Tyr Phe Leu Ile Leu Phe Ser Ile His Leu Ile Tyr Tyr Phe 1420
 1025 1030 1035
 Ile Phe Leu Gly Phe Gln Leu Ser Trp Glu Leu Met Phe Val Leu 1430
 1040 1045 1050
 Val Val Cys Ile Ile Gly Cys Ala Val Ser Leu Ile Phe Leu Thr 1440
 1055 1060 1065
 Tyr Val Leu Ser Phe Ile Phe Arg Lys Trp Arg Lys Asn Asn Phe 1450
 1070 1075 1080
 Phe Trp Ser Phe Gly Phe Phe Ile Val Ser Ile Tyr Thr Asp Phe 1460
 1085 1090 1095
 Ser Phe His Tyr Asn Val Ser Arg Cys Asp Phe Leu Phe Ile Phe 1470
 1100 1105 1110
 Ile Phe Val Cys Leu Phe Ile Ala His His Phe Ser Phe Cys Ser 1480
 1115 1120 1125
 Pro Tyr Leu Gln Ser Val Ile Phe Leu Phe Val Ile Arg Cys Leu 1490
 1130 1135 1140
 Glu Met Lys Tyr Gly Asn Glu Ile Met Asn Lys Asp Pro Val Phe 1500
 1145 1150 1155
 Arg Ile Ser Pro Arg Ser Arg Glu Thr His Pro Asn Pro Glu Glu 1510
 1160 1165 1170
 Pro Glu Glu Glu Asp Glu Asp Val Gln Ala Glu Arg Val Gln Ala 1520
 1175 1180 1185
 Ala Asn Ala Leu Thr Ala Pro Asn Leu Glu Glu Pro Val Ile 1530
 1190 1195 1200
 Thr Ala Ser Cys Leu His Lys Glu Tyr Tyr Glu Thr Lys Lys Ser 1540
 1205 1210 1215
 Cys Phe Ser Thr Arg Lys Lys Lys Ile Ala Ile Arg Asn Val Ser 1550
 1220 1225 1230
 Phe Cys Val Lys Lys Gly Glu Val Val Leu Leu Glu Gly His Asn 1560
 1235 1240 1245
 Gly Ala Gly Lys Ser Thr Ser Ile Lys Met Ile Thr Gly Cys Thr 1570

Lys Pro Thr Ala Gly Val Val Leu Gln Gly Ser Arg Ala Ser 1250
 1265 1270 1275
 Val Arg Gln Gln His Asp Asn Ser Leu Lys Phe Leu Gly Tyr Cys 1280
 1285 1290 1295
 Pro Gln Glu Asn Ser Leu Trp Pro Lys Leu Thr Met Lys Glu His 1300
 1295 1300 1305
 Leu Glu Leu Tyr Ala Val Lys Gly Leu Gly Lys Glu Asp Ala 1310
 1315 1320 1325
 Ala Leu Ser Ile Ser Arg Leu Val Glu Ala Leu Lys Leu Gln Glu 1330
 1335 1340 1345
 Gln Leu Lys Ala Pro Val Lys Thr Leu Ser Glu Gly Ile Lys Arg 1350
 1355 1360 1365
 Lys Leu Cys Phe Val Leu Ser Ile Leu Gly Asn Pro Ser Val Val 1370
 1375 1380 1385
 Leu Leu Asp Glu Pro Phe Thr Gly Met Asp Pro Glu Gly Gln Gln 1390
 1395 1400 1405
 Gln Met Trp Gln Ile Leu Gln Ala Thr Val Lys Asn Lys Glu Arg 1410
 1415 1420 1425
 Gly Thr Leu Leu Thr Thr His Tyr Met Ser Glu Ala Glu Ala Val 1430
 1435 1440 1445
 Cys Asp Arg Met Ala Met Met Val Ser Gly Thr Leu Arg Cys Ile 1450
 1455 1460 1465
 Gly Ser Ile Gln His Leu Lys Asn Lys Phe Gly Arg Asp Tyr Leu 1470
 1475 1480 1485
 Leu Glu Ile Lys Met Lys Glu Pro Thr Thr Val Glu Ala Leu His 1490
 1495 1500 1505
 Thr Glu Ile Leu Lys Leu Phe Pro Gln Ala Ala Trp Gln Glu Arg 1510
 1515 1520 1525
 Tyr Ser Ser Leu Met Ala Tyr Lys Leu Pro Val Glu Asp Val His 1530
 1535 1540 1545
 Pro Leu Ser Arg Ala Phe Phe Lys Leu Glu Ala Met Lys Gln Thr 1550
 1555 1560 1565
 Phe Asn Leu Glu Glu Tyr Ser Leu Ser Gln Ala Thr Leu Glu Gln 1570
 1575 1580 1585
 Val Phe Leu Glu Leu Cys Lys Glu Gln Glu Leu Gly Asn Val Asp 1590
 1595 1600 1605
 Asp Lys Ile Asp Thr Thr Val Glu Trp Lys Leu Leu Pro Gln Glu 1610
 1615 1620 1625
 Asp Pro 1630
 1635 1640 1645
 <210> 13
 <211> 698
 <212> PRT
 <213> Homo sapiens
 <220>
 <221> misc_feature
 <223> Incyte ID No: 6952742CD1
 <400> 13
 Met Asp Glu Ser Pro Glu Pro Leu Gln Gln Gly Arg Gly Pro Val 1
 5 10 15
 Pro Val Arg Arg Gln Arg Pro Ala Pro Arg Gly Leu Arg Glu Met 20
 25 30 35
 Leu Lys Ala Arg Leu Trp Cys Ser Cys Ser Cys Ser Val Leu Cys 40
 45 50 55
 Val Arg Ala Leu Val Gln Asp Leu Leu Pro Ala Thr Arg Trp Leu 60
 65 70 75
 Arg Gln Tyr Arg Pro Arg Glu Tyr Leu Ala Asp Val Met Ser 75
 Gly Leu Val Ile Gly Ile Ile Leu Ala Ile Ala Tyr Ser Leu Leu

Ala Gly Leu Gln Pro Ile Tyr Ser Leu Tyr Thr Ser Phe Phe Ala 90
80
95
Asn Leu Ile Tyr Phe Leu Met Gly Thr Ser Arg His Val Ser Val 105
110
Gly Ile Phe Ser Leu Leu Cys Leu Met Val Gly Gln Val Val Asp 120
125
Arg Glu Leu Gln Leu Ala Gly Phe Asp Pro Ser Gln Asp Gly Leu 135
140
Gln Pro Gly Ala Asn Ser Ser Thr Leu Asn Gly Ser Ala Ala Met 150
155
Leu Asp Cys Gly Arg Asp Cys Tyr Ala Ile Arg Val Ala Thr Ala 165
170
Leu Thr Leu Met Thr Gly Leu Tyr Gln Val Leu Met Gly Val Leu 180
185
Arg Leu Gly Phe Val Ser Ala Tyr Leu Ser Gln Pro Leu Leu Asp 195
200
Gly Phe Ala Met Gly Ala Ser Val Thr Ile Leu Thr Ser Gln Leu 210
215
Lys His Leu Leu Gly Val Arg Ile Pro Arg His Gln Gly Pro Gly 225
230
Met Val Val Leu Thr Trp Leu Ser Leu Leu Arg Gly Ala Gly Gln 240
245
Ala Asn Val Cys Asp Val Val Thr Ser Thr Val Cys Leu Ala Val 255
260
Leu Leu Ala Ala Lys Glu Leu Ser Asp Arg Tyr Arg His Arg Leu 270
275
Arg Val Pro Leu Pro Thr Glu Leu Leu Val Ile Val Val Ala Thr 285
290
Leu Val Ser His Phe Gly Gln Leu His Lys Arg Phe Gly Ser Ser 300
305
Val Ala Gly Asp Ile Pro Thr Gly Phe Met Pro Pro Gln Val Pro 315
320
Glu Pro Arg Leu Met Gln Arg Val Ala Leu Asp Ala Val Ala Leu 330
335
Ala Leu Val Ala Ala Phe Ser Ile Ser Leu Ala Glu Met Phe 345
350
Ala Arg Ser His Gly Tyr Ser Val Arg Ala Asn Gln Glu Leu Leu 360
365
Ala Val Gly Cys Cys Asn Val Leu Pro Ala Phe Leu His Cys Phe 375
380
Ala Thr Ser Ala Ala Leu Ala Lys Ser Leu Val Lys Thr Ala Thr 390
395
Gly Cys Arg Thr Gln Leu Ser Ser Val Val Ser Ala Thr Val Val 405
410
Leu Leu Val Leu Leu Ala Leu Ala Pro Leu Phe His Asp Leu Gln 420
425
Arg Ser Val Leu Ala Cys Val Ile Val Val Ser Leu Arg Gly Ala 435
440
Leu Arg Lys Val Trp Asp Leu Pro Arg Leu Trp Arg Met Ser Pro 450
455
Ala Asp Ala Leu Val Trp Ala Gly Thr Val Ala Thr Cys Met Leu 465
470
Val Ser Thr Glu Ala Gly Leu Leu Ala Gly Val Ile Leu Ser Leu 480
485
Leu Ser Leu Ala Gly Arg Thr Gln Ser His Gly Thr Ala Leu Leu 495
500
Ala Arg Ile Gly Asp Thr Ala Phe Tyr Glu Asp Ala Thr Glu Phe 510
515
Glu Gly Leu Val Pro Glu Pro Gly Val Arg Val Phe Arg Phe Gly 525
530
Gly Pro Leu Tyr Tyr Ala Asn Lys Asp Phe Phe Leu Gln Ser Leu 540
545
550
555

Tyr Ser Leu Thr Gly Leu Asp Ala Gly Cys Met Ala Ala Arg Arg 570
565
Lys Glu Gly Gly Ser Glu Thr Gly Val Gly Glu Gly Gly Pro Ala 585
590
Gln Gly Glu Asp Leu Gly Pro Val Ser Thr Arg Ala Ala Leu Val 600
595
Pro Ala Ala Ala Gly Phe His Thr Val Ile Asp Cys Ala Pro 615
620
Leu Leu Phe Leu Asp Ala Ala Gly Val Ser Thr Leu Gln Asp Leu 630
625
Arg Arg Asp Tyr Gly Ala Leu Gly Ile Ser Leu Leu Leu Ala Cys 645
635
Cys Ser Pro Pro Val Arg Asp Ile Leu Ser Arg Gly Gly Phe Leu 655
650
Gly Glu Gly Pro Gly Asp Thr Ala Glu Glu Gln Leu Phe Leu 660
665
Ser Val His Asp Ala Val Gln Thr Ala Arg Ala Arg His Arg Glu 675
680
Leu Glu Ala Thr Asp Ala His Leu 685
695

<210> 14
<211> 766
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<222> Incyte ID No: 7478795CD1

<400> 14
Met Arg Leu Trp Lys Ala Val Val Thr Leu Ala Phe Met Ser 15
1
Val Asp Ile Cys Val Thr Thr Ala Ile Tyr Val Phe Ser His Leu 20
25
Asp Arg Ser Leu Leu Glu Asp Ile Arg His Phe Asn Ile Phe Asp 30
35
Ser Val Leu Asp Leu Trp Ala Ala Cys Leu Tyr Arg Ser Cys Leu 40
50
Leu Leu Gly Ala Thr Ile Gly Val Ala Lys Asn Ser Ala Leu Gly 55
60
Pro Arg Arg Leu Arg Ala Ser Trp Leu Val Ile Thr Leu Val Cys 70
80
Leu Phe Val Gly Ile Tyr Ala Met Val Lys Leu Leu Phe Ser 85
90
Glu Val Arg Arg Pro Ile Arg Asp Pro Trp Phe Trp Ala Leu Phe 100
110
Val Trp Thr Tyr Ile Ser Leu Gly Ala Ser Phe Leu Leu Trp Trp 115
125
Leu Leu Ser Thr Val Arg Pro Gly Thr Gln Ala Leu Glu Pro Gly 130
140
Ala Ala Thr Glu Ala Glu Gly Phe Pro Gly Ser Gly Arg Pro Pro 145
155
Pro Glu Gln Ala Ser Gly Ala Thr Leu Gln Lys Leu Leu Ser Tyr 160
170
Thr Lys Pro Asp Val Ala Phe Leu Val Ala Ala Ser Phe Phe Leu 175
185
Ile Val Ala Ala Leu Leu Gly Glu Thr Phe Leu Pro Tyr Tyr Gly 190
200
Arg Ala Ile Asp Gly Ile Val Ile Gln Lys Ser Met Asp Gln Phe 205
215
Ser Thr Ala Val Ile Val Cys Leu Leu Ala Ile Gly Ser Ser 220
230
235
240
245

Phe Ala Ala Gly Ile Arg Gly Gly Ile Phe Thr Leu Ile Phe Ala
 245 250 255
 Arg Leu Asn Ile Arg Leu Arg Asn Cys Leu Phe Arg Ser Leu Val
 260 265 270
 Ser Gln Glu Thr Ser Phe Asp Glu Asn Arg Thr Gly Asp Leu
 275 280 285
 Ile Ser Arg Leu Thr Ser Asp Thr Thr Met Val Ser Asp Leu Val
 290 295 300
 Ser Gln Asn Ile Asn Val Phe Leu Arg Asn Thr Val Lys Val Thr
 305 310 315
 Gly Val Val Val Phe Met Phe Ser Leu Ser Trp Gln Leu Ser Leu
 320 325 330
 Val Thr Phe Met Gly Phe Pro Ile Ile Met Met Val Ser Asn Ile
 335 340 345
 Tyr Gly Lys Tyr Tyr Lys Arg Leu Ser Lys Glu Val Gln Asn Ala
 350 355 360
 Leu Ala Arg Ala Ser Asn Thr Ala Glu Glu Thr Ile Ser Ala Met
 365 370 375
 Lys Thr Val Arg Ser Phe Ala Asn Glu Glu Glu Ala Glu Val
 380 385 390
 Tyr Leu Arg Lys Leu Gln Gln Val Tyr Lys Leu Asn Arg Lys Glu
 395 400 405
 Ala Ala Ala Tyr Met Tyr Tyr Val Trp Gly Ser Gly Leu Thr Leu
 410 415 420
 Leu Val Val Gln Val Ser Ile Leu Tyr Tyr Gly Gly His Leu Val
 425 430 435
 Ile Ser Gly Gln Met Thr Ser Gly Asn Leu Ile Ala Phe Ile Ile
 440 445 450
 Tyr Glu Phe Val Leu Gly Asp Cys Met Glu Ser Val Gly Ser Val
 455 460 465
 Tyr Ser Gly Leu Met Gln Gly Val Gly Ala Ala Glu Lys Val Phe
 470 475 480
 Glu Phe Ile Asp Arg Gln Pro Thr Met Val His Asp Gly Ser Leu
 485 490 495
 Ala Pro Asp His Leu Glu Gly Arg Val Asp Phe Glu Asn Val Thr
 500 505 510
 Phe Thr Tyr Arg Thr Arg Pro His Thr Gln Val Leu Gln Asn Val
 515 520 525
 Ser Phe Ser Leu Ser Pro Gly Lys Val Thr Ala Leu Val Gly Pro
 530 535 540
 Ser Gly Ser Gly Ser Ser Cys Val Asn Ile Leu Glu Asn Phe
 545 550 555
 Tyr Pro Leu Glu Gly Arg Val Leu Leu Asp Gly Lys Pro Ile
 560 565 570
 Ser Ala Tyr Asp His Lys Tyr Leu His Arg Val Ile Ser Leu Val
 575 580 585
 Ser Gln Glu Pro Val Leu Phe Ala Arg Ser Ile Thr Asp Asn Ile
 590 595 600
 Ser Tyr Gly Leu Pro Thr Val Pro Phe Glu Met Val Val Glu Ala
 605 610 615
 Ala Gln Lys Ala Asn Ala His Gly Phe Ile Met Glu Leu Gln Asp
 620 625 630
 Gly Tyr Ser Thr Thr Glu Thr Gly Glu Lys Ala Gln Leu Ser Gly
 635 640 645
 Gly Gln Lys Gln Arg Val Ala Met Ala Ala Leu Val Arg Asn
 650 655 660
 Pro Pro Val Leu Ile Leu Asp Glu Ala Thr Ser Ala Leu Asp Ala
 665 670 675
 Glu Ser Glu Tyr Leu Ile Gln Gln Ala Ile His Gly Asn Leu Gln
 680 685 690
 Lys His Thr Val Leu Ile Ile Ala His Arg Leu Ser Thr Val Glu
 695 700 705
 His Ala His Leu Ile Val Val Leu Asp Lys Gly Arg Val Val Gln

28/85

Gln Gly Thr His Gln Gln Leu Leu Ala Gln Gly Leu Tyr Ala
 710 715 720
 Lys Leu Val Gln Arg Gln Met Leu Gly Leu Gln Pro Ala Ala Asp
 725 730 735
 Phe Thr Ala Gly His Asn Glu Pro Val Ala Asn Gly Ser His Lys
 740 745 750
 Ala 755 760 765

 <210> 15
 <211> 450
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> misc_feature
 <223> Incyte ID No: 656293CD1

 <400> 15
 Met Gly Leu Arg Ser His His Leu Ser Leu Gly Leu Leu Leu
 1 5 10 15
 Phe Leu Leu Pro Ala Glu Cys Leu Gly Ala Glu Gly Arg Leu Ala
 20 25 30
 Leu Lys Leu Phe Arg Asp Leu Phe Ala Asn Tyr Thr Ser Ala Leu
 35 40 45
 Arg Pro Val Ala Asp Thr Asp Gln Thr Leu Asn Val Thr Leu Glu
 50 55 60
 Val Thr Leu Ser Gln Ile Ile Asp Met Asp Glu Arg Asn Gln Val
 65 70 75
 Leu Thr Leu Tyr Leu Trp Ile Arg Gln Glu Trp Thr Asp Ala Tyr
 80 85 90
 Leu Arg Trp Asp Pro Asn Ala Tyr Gly Gly Leu Asp Ala Ile Arg
 95 100 105
 Ile Pro Ser Ser Leu Val Trp Arg Pro Asp Ile Val Leu Tyr Asn
 110 115 120
 Lys Ala Asp Ala Gln Pro Gly Ser Ala Ser Thr Asn Val Val
 125 130 135
 Leu Arg His Asp Gly Ala Val Arg Trp Asp Ala Pro Ala Ile Thr
 140 145 150
 Arg Ser Ser Cys Arg Val Asp Val Ala Ala Phe Pro Phe Asp Ala
 155 160 165
 Gln His Cys Gly Leu Thr Phe Gly Ser Trp Thr His Gly Gly His
 170 175 180
 Gln Leu Asp Val Arg Pro Arg Gly Ala Ala Ser Leu Ala Asp
 185 190 195
 Phe Val Glu Asn Val Glu Trp Arg Val Leu Gly Met Pro Ala Arg
 200 205 210
 Arg Arg Val Leu Thr Tyr Gly Cys Cys Ser Glu Pro Tyr Pro Asp
 215 220 225
 Val Thr Phe Thr Leu Leu Leu Arg Arg Ala Ala Ala Tyr Val
 230 235 240
 Cys Asn Leu Leu Leu Pro Cys Val Leu Ile Ser Leu Leu Ala Pro
 245 250 255
 Leu Ala Phe His Leu Pro Ala Asp Ser Gly Glu Lys Val Ser Leu
 260 265 270
 Gly Val Thr Val Leu Leu Ala Leu Thr Val Phe Gln Leu Leu
 275 280 285
 Ala Glu Ser Met Pro Pro Ala Glu Ser Val Pro Leu Ile Gly Lys
 290 295 300
 Tyr Tyr Met Ala Thr Met Thr Met Val Thr Phe Ser Thr Ala Leu
 305 310 315
 Thr Ile Leu Ile Met Asn Leu His Tyr Cys Gly Pro Ser Val Arg

29/85

320 Pro Val Pro Ala Trp Ala Arg Ala Leu Leu Gly His Leu Ala 330
 335 Arg Gly Leu Cys Val Arg Glu Arg Gly Glu Pro Cys Gly Gln Ser 345
 350 Arg Pro Glu Leu Ser Pro Ser Pro Gln Ser Pro Glu Gly Gly 360
 365 Ala Gly Pro Ala Gly Pro Cys His Glu Pro Arg Cys Leu Cys 375
 380 Arg Gln Glu Ala Leu Leu His Val Ala Thr Ile Ala Asn Thr 385
 395 Phe Arg Ser His Arg Ala Ala Gln Arg Cys His Glu Asp Trp Lys 405
 410 Arg Leu Ala Arg Val Met Asp Arg Phe Phe Leu Ala Ile Phe Phe 420
 425 Ser Met Ala Leu Val Met Ser Leu Leu Val Leu Val Gln Ala Leu 435
 440
 <210> 16
 <211> 260
 <212> PRT
 <213> Homo sapiens
 <220>
 <221> misc_feature
 <223> Incyte ID No: 7473957CD1

Leu Glu Phe Gln Ala 260
 <210> 17
 <211> 506
 <212> PRT
 <213> Homo sapiens
 <220>
 <221> misc_feature
 <223> Incyte ID No: 7474111CD1
 <400> 17
 Met Ser Ser Glu Pro Glu Leu Gly Ser Gly Gln Phe Leu Glu Lys Ala 15
 1 Leu Gln Thr Pro Ser Val Pro Ala Pro Glu Ser Thr Leu Gly Phe 25
 Glu Pro Gly Leu Leu Lys Gly Ala Leu Gly Thr Ala Gln Phe Ile 30
 35 Pro Met Ala Gln Gly Arg Thr Arg Glu Gln Ala Ser Arg Arg Trp 45
 50 Ala Pro Arg Ser Pro Ala Leu Arg Thr Pro Pro Arg His Tyr Gly 60
 65 Pro Glu Arg Arg Gly Arg Thr Ala Ser Arg Gly Gly Glu Pro Glu 75
 80 Val Gln Gly Gly Ala Pro Gly Asn Pro Ser Pro Ser Lys Pro Gly 90
 95 Ser Pro Gln Gly Val Gly Pro Ala Ala Trp Glu Arg Ala Pro Arg 105
 110 Pro Arg Cys Ala Gln Pro Ser Gly Ala Arg Val Gly Glu Arg Thr 120
 125 Gln Pro Arg Ser Gln Pro Val Gly Leu Ser Arg Gly Ala Gly Glu 135
 140 Asp Ser Pro Ala Thr Arg Ser Gly Ala Ala Ser Val Val Leu Asn 150
 155 Val Gly Gly Ala Arg Tyr Ser Leu Ser Arg Glu Leu Lys Asp 165
 170 Phe Pro Leu Arg Arg Val Ser Arg Leu His Gly Cys Arg Ser Glu 180
 185 Arg Asp Val Leu Glu Val Cys Asp Asp Tyr Asp Arg Glu Arg Asn 195
 200 Glu Tyr Phe Phe Asp Arg His Ser Glu Ala Phe Gly Phe Ile Leu 210
 215 Leu Tyr Ala Ala Pro Ser Arg Arg Trp Leu Glu Arg Met Arg Arg 225
 230 Thr Phe Glu Glu Pro Thr Ser Ser Leu Ala Ala Gln Ile Leu Ala 240
 245 Ser Val Ser Val Val Phe Val Ile Val Ser Met Val Val Leu Cys 255
 260 Ala Ser Thr Leu Pro Asp Trp Arg Asn Ala Ala Ala Asp Asn Arg 270
 275 Ser Leu Asp Asp Arg Ser Arg Ile Ile Glu Ala Ile Cys Ile Gly 285
 290 Trp Phe Thr Ala Glu Cys Ile Val Arg Phe Ile Val Ser Lys Asn 300
 305 Lys Cys Glu Phe Val Lys Arg Pro Leu Asn Ile Ile Asp Leu Leu 315
 320 Ala Ile Thr Pro Tyr Ile Ser Val Leu Met Thr Val Phe Thr 330
 335 Gly Glu Asn Ser Gln Leu Gln Arg Ala Gly Val Thr Leu Arg Val 345
 350 Leu Arg Met Met Arg Ile Phe Trp Val Ile Lys Leu Ala Arg His 360
 365

Phe Ile Gly Leu Gln Thr Leu Gly Leu Thr Leu Lys Arg Cys Tyr
380
Arg Gln Met Val Met Leu Leu Val Phe Ile Cys Val Ala Met Ala
395
Ile Phe Ser Ala Leu Ser Gln Leu Leu Gln His Gly Leu Asp Leu
410
Glu Thr Ser Asn Lys Asp Phe Thr Ser Ile Pro Ala Ala Cys Trp
425
Trp Val Ile Ile Ser Met Thr Thr Val Gly Tyr Gly Asp Met Tyr
440
Pro Ile Thr Val Pro Gly Arg Ile Leu Gly Val Cys Val Val
455
Ser Gly Ile Val Leu Leu Ala Leu Pro Ile Thr Phe Ile Tyr His
470
Ser Phe Val Gln Cys Tyr His Gln Leu Lys Phe Arg Ser Ala Arg
485
Tyr Ser Arg Ser Leu Ser Thr Gln Phe Leu Asn 500

<210> 18
<211> 506
<212> PRT
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No: 7480826CD1

<400> 18
Met Lys Lys Ala Gln Met Gly Arg Phe Ser Ile Ser Pro Asp Gln
1
1
Asp Ser Ser Ser Tyr Ser Ser Asn Ser Asp Phe Asn Tyr Ser Tyr
20
Pro Thr Lys Gln Ala Ala Leu Lys Ser His Tyr Ala Asp Val Asp
35
Pro Gln Asn Gln Asn Phe Leu Leu Gln Ser Asn Leu Gly Lys Lys
50
Lys Tyr Gln Thr Gln Phe His Pro Gly Thr Thr Ser Phe Gly Met
65
Ser Val Phe Asn Leu Ser Asn Ala Ile Val Gly Ser Gly Ile Leu
80
Gly Leu Ser Tyr Ala Met Ala Asn Thr Gly Ile Ala Leu Phe Ile
95
Ile Leu Leu Thr Phe Val Ser Ile Phe Ser Leu Tyr Ser Val His
110
Leu Leu Leu Lys Thr Ala Asn Gln Gly Ser Ser Leu Leu Tyr Gln
125
Gln Leu Gly Tyr Lys Ala Phe Gly Leu Val Gly Lys Leu Ala Ala
140
Ser Gly Ser Ile Thr Met Gln Asn Ile Gly Ala Met Ser Ser Tyr
155
Leu Phe Ile Val Lys Tyr Gln Leu Pro Leu Val Ile Gln Ala Leu
170
Thr Asn Ile Gln Asp Lys Thr Gly Leu Trp Tyr Leu Asn Gly Asn
185
Tyr Leu Val Leu Leu Val Ser Leu Val Val Ile Leu Pro Leu Ser
200
Leu Phe Arg Asn Leu Gly Tyr Leu Gly Tyr Thr Ser Gly Leu Ser
215
Leu Leu Cys Met Val Phe Phe Leu Ile Val Ile Cys Lys Lys
230
Phe Gln Val Pro Cys Pro Val Gln Ala Ala Leu Ile Ile Asn Gln
245

Thr Ile Asn Thr Thr Leu Thr Gln Pro Thr Ala Leu Val Pro Ala
260
Leu Ser His Asn Val Thr Gln Asn Asp Ser Cys Arg Pro His Tyr
275
Phe Ile Phe Asn Ser Gln Thr Val Tyr Ala Val Pro Ile Leu Ile
290
Phe Ser Phe Val Cys His Pro Ala Val Leu Pro Ile Tyr Gln Gln
305
Leu Lys Asp Arg Ser Arg Arg Arg Met Met Asn Val Ser Lys Ile
320
Ser Phe Phe Ala Met Phe Leu Met Tyr Leu Leu Ala Ala Leu Phe
335
Gly Tyr Leu Thr Thr Phe Tyr Gln His Val Gln Ser Gln Leu Leu His
350
Thr Tyr Ser Ser Ile Leu Gly Thr Asp Ile Leu Leu Leu Ile Val
365
Arg Leu Ala Val Leu Met Ala Val Thr Leu Thr Val Pro Val Val
380
Ile Phe Pro Ile Arg Ser Ser Val Thr His Leu Leu Cys Ala Ser
395
Lys Asp Phe Ser Trp Trp Arg His Ser Leu Ile Thr Val Ser Ile
410
Leu Ala Phe Thr Asn Leu Leu Val Ile Phe Val Pro Thr Ile Arg
425
Asp Ile Phe Gly Phe Ile Gly Ala Ser Ala Ala Ser Met Leu Ile
440
Phe Ile Leu Pro Ser Ala Phe Tyr Ile Lys Leu Val Lys Lys Gln
455
Pro Met Lys Ser Val Gln Lys Ile Gly Ala Leu Phe Phe Leu Leu
470
Ser Gly Val Leu Val Met Thr Gly Ser Met Ala Leu Ile Val Leu
485
Asp Trp Val His Asn Ala Pro Gly Gly His 505

<210> 19
<211> 315
<212> PRT
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No: 6025572CD1

<400> 19
Met His Arg Gln Pro Ala Lys Lys Lys Ala Gln Lys Arg Leu Phe
1
1
Asp Ala Ser Ser Phe Gly Lys Asp Leu Leu Ala Gly Gly Val Ala
20
Ala Ala Val Ser Lys Thr Ala Val Ala Pro Ile Gln Arg Val Lys
35
Leu Leu Leu Gln Val Gln Ala Ser Ser Lys Gln Ile Ser Pro Gln
50
Ala Arg Tyr Lys Gly Met Val Asp Cys Leu Val Arg Ile Pro Arg
65
Gln Gln Gly Phe Phe Ser Phe Trp Arg Gly Asn Leu Ala Asn Val
80
Ile Arg Tyr Phe Pro Thr Gln Ala Leu Asn Phe Ala Phe Lys Asp
95
Lys Tyr Lys Gln Leu Phe Met Ser Gly Val Asn Lys Gln Lys Gln
110
Phe Trp Arg Trp Phe Leu Ala Asn Leu Ala Ser Gly Gly Ala Ala
125

Gly Ala Thr Ser Leu Cys Val Val Tyr Pro Leu Asp Phe Ala Arg
140 145 150
Thr Arg Leu Gly Val Asp Ile Gly Lys Pro Glu Arg Gln
155 160 165
Phe Lys Gly Leu Gly Asp Cys Ile Met Lys Ile Ala Lys Ser Asp
170 175 180
Gly Ile Ala Gly Leu Tyr Gln Gly Phe Gly Val Ser Val Gln Gly
185 190 195
Ile Ile Val Tyr Arg Ala Ser Tyr Phe Gly Ala Tyr Asp Thr Val
200 205 210
Lys Gly Leu Leu Pro Lys Pro Lys Lys Thr Pro Phe Leu Val Ser
215 220 225
Phe Phe Ile Ala Gln Val Val Thr Thr Cys Ser Gly Ile Leu Ser
230 235 240
Tyr Pro Phe Asp Thr Val Arg Arg Arg Met Met Met Gln Ser Gly
245 250 255
Glu Ala Lys Arg Gln Tyr Lys Gly Thr Leu Asp Cys Phe Val Lys
260 265 270
Ile Tyr Gln His Glu Gly Ile Ser Ser Phe Phe Arg Gly Ala Phe
275 280 285
Ser Asn Val Leu Arg Gly Thr Gly Gly Ala Leu Val Leu Val
290 295 300
Tyr Asp Lys Ile Lys Glu Phe Phe His Ile Asp Ile Gly Gly Arg
305 310 315

<210> 20

<211> 540

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5686561CD1

<400> 20

Met Val Pro Ala Gly Trp Val Arg Gly Leu Glu Leu Ser Leu Trp
5 10 15
Gly Gly Asp Pro Val Val Pro Trp Ser Cys Arg Phe Cys Ser Gln
20 25 30
Gln Asp Asp Gly Gln Asp Arg Glu Arg Leu Thr Tyr Phe Gln Asn
35 40 45
Leu Pro Glu Ser Leu Thr Ser Leu Leu Val Leu Leu Thr Thr Ala
50 55 60
Asn Asn Pro Asp Val Met Ile Pro Ala Tyr Ser Lys Asn Arg Ala
65 70 75
Tyr Ala Ile Phe Phe Ile Val Phe Thr Val Ile Gly Ser Leu Phe
80 85 90
Leu Met Asn Leu Leu Thr Ala Ile Ile Tyr Ser Gln Phe Arg Gly
95 100 105
Tyr Leu Met Lys Ser Leu Gln Thr Ser Leu Phe Arg Arg Arg Leu
110 115 120
Gly Thr Arg Ala Ala Phe Glu Val Leu Ser Ser Met Val Gly Gly
125 130 135
Gly Gly Ala Phe Pro Gln Ala Val Gly Val Lys Pro Gln Asn Leu
140 145 150
Leu Gln Val Leu Gln Lys Val Gln Leu Asp Ser Ser His Lys Gln
155 160 165
Ala Met Met Glu Lys Val Arg Ser Tyr Asp Ser Val Leu Leu Ser
170 175 180
Ala Glu Glu Phe Gln Lys Leu Phe Asn Glu Leu Asp Arg Ser Val
185 190 195
Val Lys Glu His Pro Pro Arg Pro Glu Tyr Gln Ser Pro Phe Leu

34/85

Gln Ser Ala Gln Phe Leu Phe Gly His Tyr Tyr Phe Asp Tyr Leu
200 205 210
Gly Asn Leu Ile Ala Leu Ala Asn Leu Val Ser Ile Cys Val Phe
215 220 225
Leu Val Leu Asp Ala Asp Val Leu Pro Ala Glu Arg Asp Phe
230 235 240
Ile Leu Gly Ile Leu Asn Cys Val Phe Ile Val Tyr Tyr Leu Leu
245 250 255
Glu Met Leu Leu Lys Val Phe Ala Leu Gly Leu Arg Gly Tyr Leu
260 265 270
Ser Tyr Pro Ser Asn Val Phe Asp Gly Leu Leu Thr Val Val Leu
275 280 285
Leu Val Leu Glu Ile Ser Thr Leu Ala Val Tyr Arg Leu Pro His
290 295 300
Pro Gly Trp Arg Pro Glu Met Val Gly Leu Ser Leu Trp Asp
305 310 315
Met Thr Arg Met Leu Asn Met Leu Ile Val Phe Arg Phe Leu Arg
320 325 330
Ile Ile Pro Ser Met Lys Pro Met Ala Val Val Ala Ser Thr Val
335 340 345
Leu Gly Leu Val Gln Asn Met Arg Ala Phe Gly Gly Ile Leu Val
350 355 360
Val Val Tyr Tyr Val Phe Ala Ile Ile Gly Ile Asn Leu Phe Arg
365 370 375
Gly Val Ile Val Ala Leu Pro Gly Asn Ser Ser Leu Ala Pro Ala
380 385 390
Asn Gly Ser Ala Pro Cys Gly Ser Phe Phe Gln Leu Glu Tyr Trp
395 400 405
Ala Asn Asn Phe Asp Asp Phe Ala Ala Leu Val Thr Leu Trp
410 415 420
Asn Leu Met Val Val Asn Asn Trp Gln Val Phe Leu Asp Ala Tyr
425 430 435
Arg Arg Tyr Ser Gly Pro Trp Ser Lys Ile Tyr Phe Val Leu Trp
440 445 450
Trp Leu Val Ser Ser Val Ile Trp Val Asn Leu Phe Leu Ala Leu
455 460 465
Ile Leu Glu Asn Phe Leu His Lys Trp Asp Pro Arg Ser His Leu
470 475 480
Gln Pro Leu Ala Gly Thr Pro Glu Ala Thr Tyr Gln Met Thr Val
485 490 495
Glu Leu Leu Phe Arg Asp Ile Leu Glu Glu Pro Gly Glu Asp Glu
500 505 510
Leu Thr Glu Arg Leu Ser Gln His Pro His Leu Trp Leu Cys Arg
515 520 525
530 535 540

<210> 21

<211> 322

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1553725CD1

<400> 21

Met Glu Ala Asp Leu Ser Gly Phe Asn Ile Asp Ala Pro Arg Trp
1 5 10 15
Asp Gln Arg Thr Phe Leu Gly Arg Val Lys His Phe Leu Asn Ile
20 25 30
Thr Asp Pro Arg Thr Val Phe Val Ser Glu Arg Glu Leu Asp Trp
35 40 45

35/85

Ala Lys Val Met Val Glu Lys Ser Arg Met Gly Val Val Pro Pro
 50 55 60
 Gly Thr Gln Val Glu Gln Leu Leu Tyr Ala Lys Lys Leu Tyr Asp
 65 70 75
 Ser Ala Phe His Pro Asp Thr Gly Glu Lys Met Asn Val Ile Gly
 80 85 90
 Arg Met Ser Phe Gln Leu Pro Gly Met Ile Ile Thr Gly Phe
 95 100 105
 Met Leu Gln Phe Tyr Arg Thr Met Pro Ala Val Ile Phe Trp Gln
 110 115 120
 Trp Val Asn Gln Ser Phe Asn Ala Leu Val Asn Tyr Thr Asn Arg
 125 130 135
 Asn Ala Ala Ser Pro Thr Ser Val Arg Gln Met Ala Leu Ser Tyr
 140 145 150
 Phe Thr Ala Thr Thr Ala Val Ala Thr Ala Val Gly Met Asn
 155 160 165
 Met Leu Thr Lys Lys Ala Pro Pro Leu Val Gly Arg Trp Val Pro
 170 175 180
 Phe Ala Ala Val Ala Ala Asn Cys Val Asn Ile Pro Met Met
 185 190 195
 Arg Gln Gln Glu Leu Ile Lys Gly Ile Cys Val Lys Asp Arg Asn
 200 205 210
 Glu Asn Glu Ile Gly His Ser Arg Arg Ala Ala Ile Ile Gly Ile
 215 220 225
 Thr Gln Val Val Ile Ser Arg Ile Thr Met Ser Ala Pro Gly Met
 230 235 240
 Ile Leu Leu Pro Val Ile Met Glu Arg Leu Glu Lys Leu His Phe
 245 250 255
 Met Gln Lys Val Lys Val Leu His Ala Pro Leu Gln Val Met Leu
 260 265 270
 Ser Gly Cys Phe Leu Ile Phe Met Val Pro Val Ala Cys Gly Lys
 275 280 285
 Phe Pro Gln Lys Cys Glu Leu Pro Val Ser Tyr Leu Glu Pro Lys
 290 295 300
 Leu Gln Asp Thr Ile Lys Ala Lys Tyr Gly Glu Leu Glu Pro
 305 310 315
 Val Tyr Phe Asn Lys Gly Leu
 320

<210> 22
 <211> 417
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc.feature
 <223> Incyte ID No: 1695770CD1

<400> 22
 Met Thr Thr Leu Val Pro Ala Thr Leu Ser Phe Leu Leu Leu Tyr
 1 5 10 15
 Thr Leu Pro Gly Gln Val Leu Leu Arg Val Ala Leu Ala Lys Glu
 20 25 30
 Glu Val Lys Ser Gly Thr Lys Gly Ser Gln Pro Met Ser Pro Ser
 35 40 45
 Asp Phe Leu Asp Lys Leu Met Gly Arg Thr Ser Gly Tyr Asp Ala
 50 55 60
 Arg Ile Arg Pro Asn Phe Lys Gly Pro Pro Val Asn Val Thr Cys
 65 70 75
 Asn Ile Phe Ile Asn Ser Phe Ser Ser Val Thr Lys Thr Thr Met
 80 85 90
 Asp Tyr Arg Val Asn Val Phe Leu Arg Gln Gln Trp Asn Asp Pro
 95 100 105

36/85

Arg Leu Ser Tyr Arg Glu Tyr Pro Asp Asp Ser Leu Asp Leu Asp
 110 115 120
 Pro Ser Met Leu Asp Ser Ile Trp Lys Pro Asp Leu Phe Phe Ala
 125 130 135
 Asn Glu Lys Gly Ala Asn Phe His Glu Val Thr Thr Asp Asn Lys
 140 145 150
 Leu Leu Arg Ile Phe Lys Asn Gly Asn Val Leu Tyr Ser Ile Arg
 155 160 165
 Leu Thr Leu Ile Leu Ser Cys Leu Met Asp Leu Lys Asn Phe Pro
 170 175 180
 Met Asp Ile Gln Thr Cys Thr Met Gln Leu Glu Ser Phe Gly Tyr
 185 190 195
 Thr Met Lys Asp Leu Val Phe Glu Trp Leu Glu Asp Ala Pro Ala
 200 205 210
 Val Gln Val Ala Glu Gly Leu Thr Leu Pro Gln Phe Ile Leu Arg
 215 220 225
 Asp Glu Lys Asp Leu Gly Cys Cys Thr Lys His Tyr Asn Thr Gly
 230 235 240
 Lys Phe Thr Cys Ile Glu Val Lys Phe His Leu Glu Arg Gln Met
 245 250 255
 Gly Tyr Tyr Leu Ile Gln Met Tyr Ile Pro Ser Leu Leu Ile Val
 260 265 270
 Ile Leu Ser Trp Val Ser Phe Trp Ile Asn Met Asp Ala Ala Pro
 275 280 285
 Ala Arg Val Gly Leu Gly Ile Thr Thr Val Leu Thr Met Thr Thr
 290 295 300
 Gln Ser Ser Gly Ser Arg Ala Ser Leu Pro Lys Val Ser Tyr Val
 305 310 315
 Lys Ala Ile Asp Ile Trp Met Ala Val Cys Leu Leu Phe Val Phe
 320 325 330
 Ala Ala Leu Leu Glu Tyr Ala Ala Ile Asn Phe Val Ser Arg Gln
 335 340 345
 His Lys Glu Phe Ile Arg Leu Arg Arg Arg Gln Arg Arg
 350 355 360
 Leu Glu Glu Asp Ile Ile Gln Glu Ser Arg Phe Tyr Phe Arg Gly
 365 370 375
 Tyr Gly Leu Gly His Cys Leu Gln Ala Arg Asp Gly Gly Pro Met
 380 385 390
 Glu Gly Ser Gly Ile Tyr Ser Pro Gln Pro Pro Ala Pro Leu Leu
 395 400 405
 Arg Glu Gly Glu Thr Thr Arg Lys Leu Tyr Val Asp
 410 415

<210> 23
 <211> 1864
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc.feature
 <223> Incyte ID No: 4672222CD1

<400> 23
 Met Ser Gln Lys Ser Trp Ile Glu Ser Thr Leu Thr Lys Arg Glu
 1 5 10 15
 Cys Val Tyr Ile Ile Pro Ser Ser Lys Asp Pro His Arg Cys Leu
 20 25 30
 Pro Gly Cys Gln Ile Cys Gln Gln Leu Val Arg Cys Phe Cys Gly
 35 40 45
 Arg Leu Val Lys Gln His Ala Cys Phe Thr Ala Ser Leu Ala Met
 50 55 60
 Lys Tyr Ser Asp Val Lys Leu Gly Asp His Phe Asn Gln Ala Ile
 65 70 75

37/85

Glu Glu Trp Ser Val Glu Lys His Thr Glu Gln Ser Pro Thr Asp 80 85 90
Ala Tyr Gly Val Ile Asn Phe Gln Gly Ser His Ser Tyr Arg 95
Ala Lys Tyr Val Arg Leu Ser Tyr Asp Thr Lys Pro Glu Val Ile 110 115 120
Leu Gln Leu Leu Lys Glu Trp Gln Met Glu Leu Pro Lys Leu 125 130 135
Val Ile Ser Val His Gly Gly Met Gln Lys Phe Glu Leu His Pro 140 145 150
Arg Ile Lys Gln Leu Leu Gly Lys Gly Ile Lys Ala Ala Val 155 160 165
Thr Thr Gly Ala Trp Ile Leu Thr Gly Gly Val Asn Thr Gly Val 170 175 180
Ala Lys His Val Gly Asp Ala Leu Lys Gln His Ala Ser Arg Ser 185 190 195
Ser Arg Lys Ile Cys Thr Ile Gly Ile Ala Pro Trp Gly Val Ile 200 205 210
Glu Asn Arg Asn Asp Leu Val Gly Arg Asp Val Val Ala Pro Tyr 215 220 225
Gln Thr Leu Leu Asn Pro Leu Ser Lys Leu Asn Val Leu Asn Asn 230 235 240
Leu His Ser His Phe Ile Leu Val Asp Asp Gly Thr Val Gly Lys 245 250 255
Tyr Gly Ala Glu Val Arg Leu Arg Arg Glu Leu Glu Lys Thr Ile 260 265 270
Asn Gln Gln Arg Ile His Ala Arg Ile Gly Gln Gly Val Pro Val 275 280 285
Val Ala Leu Ile Phe Glu Gly Gly Pro Asn Val Ile Leu Thr Val 290 295 300
Leu Glu Tyr Leu Gln Glu Ser Pro Pro Val Pro Val Val Val Cys 305 310 315
Glu Gly Thr Gly Arg Ala Ala Asp Leu Leu Ala Tyr Ile His Lys 320 325 330
Gln Thr Glu Glu Gly Gly Asn Leu Pro Asp Ala Ala Glu Pro Asp 335 340 345
Ile Ile Ser Thr Ile Lys Lys Thr Phe Asn Phe Gly Gln Asn Glu 350 355 360
Ala Leu His Leu Phe Gln Thr Leu Met Glu Cys Met Lys Arg Lys 365 370 375
Glu Leu Ile Thr Val Phe His Ile Gly Ser Asp Glu His Gln Asp 380 385 390
Ile Asp Val Ala Ile Leu Thr Ala Leu Lys Lys Gly Thr Asn Ala 395 400 405
Ser Ala Phe Asp Gln Leu Ile Leu Thr Leu Ala Trp Asp Arg Val 410 415 420
Asp Ile Ala Lys Asn His Val Phe Val Tyr Gly Gln Gln Trp Leu 425 430 435
Val Gly Ser Leu Glu Gln Ala Met Leu Asp Ala Leu Val Met Asp 440 445 450
Arg Val Ala Phe Val Lys Leu Leu Ile Glu Asn Gly Val Ser Met 455 460 465
His Lys Phe Leu Thr Ile Pro Arg Leu Glu Leu Tyr Asn Thr 470 475 480
Lys Gln Gly Pro Thr Asn Pro Met Leu Phe His Leu Val Arg Asp 485 490 495
Val Lys Gln Gly Asn Leu Pro Pro Gly Tyr Lys Ile Thr Leu Ile 500 505 510
Asp Ile Gly Leu Val Ile Glu Tyr Leu Met Gly Gly Thr Tyr Arg 515 520 525
Cys Thr Tyr Thr Arg Lys Arg Phe Arg Leu Ile Tyr Asn Ser Leu 530 535 540
Gly Gly Asn Asn Arg Arg Ser Gly Arg Asn Thr Ser Ser Thr

Pro Gln Leu Arg 545 550 555
Lys Ser His Glu Ser Phe Gly Asn Arg Ala Asp 560 565 570
Lys Lys Glu Lys Met Arg His Asn His Phe Ile Lys Thr Ala Gln 575 580 585
Pro Tyr Arg Pro Lys Ile Asp Thr Val Met Glu Glu Gly Lys Lys 590 595 600
Lys Arg Thr Lys Asp Glu Ile Val Asp Ile Asp Asp Pro Glu Thr 605 610 615
Lys Arg Phe Pro Tyr Pro Leu Asn Glu Leu Leu Ile Trp Ala Cys 620 625 630
Leu Met Lys Arg Gln Val Met Ala Arg Phe Leu Trp Gln His Gly 635 640 645
Glu Glu Ser Met Ala Lys Ala Leu Val Ala Cys Lys Ile Tyr Arg 650 655 660
Ser Met Ala Tyr Glu Ala Lys Gln Ser Asp Leu Val Asp Asp Thr 665 670 675
Ser Glu Glu Leu Lys Gln Tyr Ser Asn Asp Phe Gly Gln Leu Ala 680 685 690
Val Glu Leu Leu Gln Ser Phe Arg Gln Asp Glu Thr Met Ala 695 700 705
Met Lys Leu Leu Thr Tyr Glu Leu Lys Asn Trp Ser Asn Ser Thr 710 715 720
Cys Leu Lys Leu Ala Val Ser Ser Arg Leu Arg Pro Phe Val Ala 725 730 735
His Thr Cys Thr Gln Met Leu Ser Asp Met Trp Met Gly Arg 740 745 750
Leu Asn Met Arg Lys Asn Ser Trp Tyr Lys Val Ile Leu Ser Ile 755 760 765
Leu Val Pro Pro Ala Ile Leu Leu Leu Glu Tyr Lys Thr Lys Ala 770 775 780
Glu Met Ser His Ile Pro Gln Ser Gln Asp Ala His Gln Met Thr 785 790 795
Met Asp Asp Ser Glu Asn Asn Phe Gln Asn Ile Thr Glu Glu Ile 800 805 810
Pro Met Glu Val Phe Lys Glu Val Arg Ile Leu Asp Ser Asn Glu 815 820 825
Gly Lys Asn Glu Met Glu Ile Gln Met Lys Ser Lys Lys Leu Pro 830 835 840
Ile Thr Arg Lys Phe Tyr Ala Phe Tyr His Ala Pro Ile Val Lys 845 850 855
Phe Trp Phe Asn Thr Leu Ala Tyr Leu Gly Phe Leu Met Leu Tyr 860 865 870
Thr Phe Val Val Leu Val Gln Met Glu Gln Leu Pro Ser Val Gln 875 880 885
Glu Trp Ile Val Ile Ala Tyr Ile Phe Thr Tyr Ala Ile Glu Lys 890 895 900
Val Arg Glu Ile Phe Met Ser Glu Ala Gly Lys Val Asn Gln Lys 905 910 915
Ile Lys Val Trp Phe Ser Asp Tyr Phe Asn Ile Ser Asp Thr Ile 920 925 930
Ala Ile Ile Ser Phe Phe Ile Gly Phe Gly Leu Arg Phe Gly Ala 935 940 945
Lys Trp Asn Phe Ala Asn Ala Tyr Asp Asn His Val Phe Val Ala 950 955 960
Gly Arg Leu Ile Tyr Cys Leu Asn Ile Ile Phe Trp Tyr Val Arg 965 970 975
Leu Leu Asp Phe Leu Ala Val Asn Gln Gln Ala Gly Pro Tyr Val 980 985 990
Met Met Ile Gly Lys Met Val Ala Asn Met Phe Tyr Ile Val Val 995 1000 1005
Ile Met Ala Leu Val Leu Leu Ser Phe Gly Val Pro Arg Lys Ala 1010 1015 1020

Ile Leu Tyr Pro His Glu Ala Pro Ser Thr Thr Leu Ala Lys Asp 1025 1030 1035
 Ile Val Phe His Pro Tyr Trp Met Ile Phe Gly Glu Val Tyr Ala 1040 1045 1050
 Tyr Glu Ile Asp Val Cys Ala Asn Asp Ser Val Ile Pro Gln Ile 1055 1060 1065
 Cys Gly Pro Gly Thr Trp Leu Thr Pro Phe Leu Gln Ala Val Tyr 1070 1075 1080
 Leu Phe Val Gln Tyr Ile Ile Met Val Asn Leu Leu Ile Ala Phe 1085 1090 1095
 Phe Asn Asn Val Tyr Leu Gln Val Lys Ala Ile Ser Asn Ile Val 1100 1105 1110
 Trp Lys Tyr Gln Arg Tyr His Phe Ile Met Ala Tyr His Glu Lys 1115 1120 1125
 Pro Val Leu Pro Pro Leu Ile Ile Leu Ser His Ile Val Ser 1130 1135 1140
 Leu Phe Cys Cys Ile Cys Lys Arg Arg Lys Asp Lys Thr Ser 1145 1150 1155
 Asp Gly Pro Lys Leu Phe Leu Thr Glu Asp Gln Lys Lys Leu 1160 1165 1170
 His Asp Phe Glu Glu Gln Cys Val Glu Met Tyr Phe Asn Glu Lys 1175 1180 1185
 Asp Asp Lys Phe His Ser Gly Ser Glu Arg Ile Arg Val Thr 1190 1195 1200
 Phe Glu Arg Val Glu Gln Met Cys Ile Gln Ile Lys Glu Val Gly 1205 1210 1215
 Asp Arg Val Asn Tyr Ile Lys Arg Ser Leu Gln Ser Leu Asp Ser 1220 1225 1230
 Gln Ile Gly His Leu Gln Asp Leu Ser Ala Leu Thr Val Asp Thr 1235 1240 1245
 Leu Lys Thr Leu Thr Ala Gln Lys Ala Ser Glu Ala Ser Lys Val 1250 1255 1260
 His Asn Glu Ile Thr Arg Glu Leu Ser Ile Ser Lys His Leu Ala 1265 1270 1275
 Gln Asn Leu Ile Asp Asp Gly Pro Val Arg Pro Ser Val Trp Lys 1280 1285 1290
 Lys His Gly Val Val Asn Thr Leu Ser Ser Ser Leu Pro Gln Gly 1295 1300 1305
 Asp Leu Glu Ser Asn Asn Pro Phe His Cys Asn Ile Leu Met Lys 1310 1315 1320
 Asp Asp Lys Asp Pro Gln Cys Asn Ile Phe Gly Gln Asp Leu Pro 1325 1330 1335
 Ala Val Pro Gln Arg Lys Glu Phe Asn Phe Pro Glu Ala Gly Ser 1340 1345 1350
 Ser Ser Gly Ala Leu Phe Pro Ser Ala Val Ser Pro Pro Glu Leu 1355 1360 1365
 Arg Gln Arg Leu His Gly Val Glu Leu Lys Ile Phe Asn Lys 1370 1375 1380
 Asn Gln Lys Leu Gly Ser Ser Ser Thr Ser Ile Pro His Leu Ser 1385 1390 1395
 Ser Pro Pro Thr Lys Phe Phe Val Ser Thr Pro Ser Gln Pro Ser 1400 1405 1410
 Cys Lys Ser His Leu Glu Thr Gly Thr Lys Asp Gln Glu Thr Val 1415 1420 1425
 Cys Ser Lys Ala Thr Glu Gly Asp Asn Thr Glu Phe Gly Ala Phe 1430 1435 1440
 Val Gly His Arg Asp Ser Met Asp Leu Gln Arg Phe Lys Glu Thr 1445 1450 1455
 Ser Asn Lys Ile Lys Ile Leu Ser Asn Asn Asn Thr Ser Glu Asn 1460 1465 1470
 Thr Leu Lys Arg Val Ser Ser Leu Ala Gly Phe Thr Asp Cys His 1475 1480 1485
 Arg Thr Ser Ile Pro Val His Ser Lys Gln Glu Lys Ile Ser Arg

40/85

Arg Pro Ser Thr Thr Asp Thr His Glu Val Asp Ser Lys Ala Ala 1490 1495 1500
 Leu Ile Pro Asp Trp Leu Gln Asp Arg Pro Ser Asn Arg Glu Met 1505 1510 1515
 Pro Ser Glu Glu Gly Thr Leu Asn Gly Leu Thr Ser Pro Phe Lys 1520 1525 1530
 Pro Ala Met Asp Thr Asn Tyr Tyr Ser Ala Val Glu Arg Asn 1535 1540 1545
 Asn Leu Met Arg Leu Ser Gln Ser Ile Pro Phe Thr Pro Val Pro 1550 1555 1560
 Pro Arg Gly Glu Pro Val Thr Val Tyr Arg Leu Glu Ser Ser 1565 1570 1575
 Pro Asn Ile Leu Asn Asn Ser Met Ser Ser Trp Ser Gln Leu Gly 1580 1585 1590
 Leu Cys Ala Lys Ile Glu Phe Leu Ser Lys Glu Glu Met Gly Gly 1595 1600 1605
 Gly Leu Arg Arg Ala Val Lys Val Gln Cys Thr Trp Ser Glu His 1610 1615 1620
 Asp Ile Leu Lys Ser Gly His Leu Tyr Ile Ile Lys Ser Phe Leu 1625 1630 1635
 Pro Glu Val Val Asn Thr Trp Ser Ser Ile Tyr Lys Glu Asp Thr 1640 1645 1650
 Val Leu His Leu Cys Leu Arg Glu Ile Gln Gln Arg Ala Ala 1655 1660 1665
 Gln Lys Leu Thr Phe Ala Phe Asn Gln Met Lys Pro Lys Ser Ile 1670 1675 1680
 Pro Tyr Ser Pro Arg Phe Leu Glu Val Phe Leu Leu Tyr Cys His 1685 1690 1695
 Ser Ala Gly Gln Trp Phe Ala Val Glu Glu Cys Met Thr Gly Glu 1700 1705 1710
 Phe Arg Lys Tyr Asn Asn Asn Asn Gly Asp Glu Ile Ile Pro Thr 1715 1720 1725
 Asn Thr Leu Glu Glu Ile Met Leu Ala Phe Ser His Trp Thr Tyr 1730 1735 1740
 Glu Tyr Thr Arg Gly Glu Leu Leu Val Leu Asp Leu Gln Gly Val 1745 1750 1755
 Gly Glu Asn Leu Thr Asp Pro Ser Val Ile Lys Ala Glu Glu Lys 1760 1765 1770
 Arg Ser Cys Asp Met Val Phe Gly Pro Ala Asn Leu Gly Glu Asp 1775 1780 1785
 Ala Ile Lys Asn Phe Arg Ala Lys His His Cys Asn Ser Cys Cys 1790 1795 1800
 Arg Lys Leu Lys Leu Pro Asp Leu Lys Arg Asn Asp Tyr Thr Pro 1805 1810 1815
 Asp Lys Ile Ile Phe Pro Gln Asp Glu Pro Ser Asp Leu Asn Leu 1820 1825 1830
 Gln Pro Gly Asn Ser Thr Lys Glu Ser Glu Ser Thr Asn Ser Val 1835 1840 1845
 Arg Leu Met Leu 1850 1855 1860

<210> 24
 <211> 1237
 <212> PRT
 <213> Homo sapiens
 <220>
 <221> misc-feature
 <223> Incyte ID No: 6176128CD1
 <400> 24
 Met Ala Arg Ala Lys Leu Pro Arg Ser Pro Ser Glu Gly Lys Ala

41/85

1	Gly	Pro	Gly	Ala	Pro	Ala	Gly	Ala	Ala	Pro	Glu	Glu	Pro	15
20	His	Gly	Leu	Ser	Pro	Leu	Leu	Pro	Ala	Arg	Gly	Gly	Ser	30
35	Gly	Ser	Asp	Val	Gly	Gln	Arg	Leu	Pro	Val	Glu	Asp	Phe	45
50	Asp	Ser	Ser	Leu	Ser	Gln	Val	Gln	Val	Glu	Phe	Tyr	Val	60
65	Asn	Thr	Phe	Lys	Glu	Arg	Leu	Lys	Leu	Phe	Phe	Ile	Lys	75
80	Arg	Ser	Ser	Leu	Arg	Ile	Arg	Leu	Phe	Asn	Phe	Ser	Leu	90
95	Leu	Thr	Cys	Leu	Leu	Tyr	Ile	Val	Arg	Val	Leu	Leu	Asp	105
110	Ala	Leu	Gly	Ile	Gly	Trp	Gly	Cys	Pro	Arg	Gln	Asn	Tyr	120
125	Phe	Asn	Asp	Ser	Ser	Glu	Ile	Asn	Trp	Ala	Pro	Ile	Leu	135
140	Val	Glu	Arg	Lys	Met	Thr	Leu	Trp	Ala	Ile	Gln	Val	Ile	150
155	Ile	Ile	Ser	Phe	Leu	Glu	Thr	Met	Leu	Ile	Tyr	Leu	Ser	165
170	Lys	Gly	Asn	Ile	Trp	Glu	Gln	Ile	Phe	Arg	Val	Ser	Phe	180
185	Glu	Met	Ile	Asn	Thr	Leu	Pro	Phe	Ile	Ile	Thr	Ile	Phe	195
200	Pro	Leu	Arg	Asn	Leu	Phe	Ile	Pro	Val	Phe	Leu	Asn	Cys	210
215	Ala	Lys	His	Ala	Leu	Glu	Asn	Met	Ile	Asn	Asp	Phe	His	225
230	Ile	Leu	Arg	Thr	Gln	Ser	Ala	Met	Phe	Asn	Gln	Val	Leu	240
245	Phe	Cys	Thr	Leu	Leu	Cys	Leu	Val	Phe	Thr	Gly	Thr	Cys	255
260	Gln	His	Leu	Glu	Arg	Ala	Gly	Glu	Asn	Leu	Ser	Leu	Thr	270
275	Phe	Tyr	Phe	Cys	Ile	Val	Thr	Phe	Ser	Thr	Val	Gly	Tyr	285
290	Val	Thr	Pro	Lys	Ile	Trp	Pro	Ser	Gln	Leu	Leu	Val	Ile	295
305	Ile	Cys	Val	Ala	Leu	Val	Val	Leu	Pro	Leu	Gln	Phe	Glu	310
320	Val	Tyr	Leu	Trp	Met	Glu	Arg	Gln	Lys	Ser	Gly	Gly	Asn	315
335	Arg	His	Arg	Ala	Gln	Thr	Glu	Lys	His	Val	Val	Leu	Cys	320
350	Ser	Leu	Lys	Ile	Asp	Leu	Leu	Met	Asp	Phe	Leu	Asn	Glu	335
365	Ala	His	Pro	Arg	Leu	Gln	Asp	Tyr	Tyr	Val	Ile	Leu	Cys	345
380	Thr	Glu	Met	Asp	Val	Gln	Val	Arg	Arg	Val	Leu	Gln	Ile	350
395	Trp	Ser	Gln	Arg	Val	Ile	Tyr	Leu	Gln	Gly	Ser	Ala	Leu	355
410	Gln	Asp	Leu	Met	Arg	Ala	Lys	Met	Asp	Asn	Gly	Glu	Ala	360
425	Ile	Leu	Ser	Ser	Arg	Asn	Glu	Val	Asp	Arg	Thr	Ala	Ala	365
440	Gln	Thr	Ile	Leu	Arg	Ala	Trp	Ala	Val	Lys	Asp	Phe	Ala	370
455	Cys	Pro	Leu	Tyr	Val	Gln	Ile	Leu	Lys	Pro	Glu	Asn	Lys	375
470														480
485	Val	Lys	Phe	Ala	Asp	His	Val	Val	Cys	Glu	Glu	Cys	Lys	495
500	Ala	Met	Leu	Ala	Leu	Asn	Cys	Ile	Cys	Pro	Ala	Thr	Ser	505
515	Ile	Thr	Leu	Leu	Val	His	Thr	Ser	Arg	Gly	Gln	Glu	Gln	520
530	Ser	Pro	Glu	Gln	Trp	Gln	Arg	Met	Tyr	Gly	Arg	Cys	Ser	525
545	Glu	Val	Tyr	His	Ile	Arg	Met	Gly	Asp	Ser	Lys	Phe	Phe	535
560	Tyr	Glu	Gly	Lys	Ser	Phe	Thr	Tyr	Ala	Ala	Phe	His	Ala	540
575	Lys	Tyr	Gly	Val	Cys	Leu	Ile	Gly	Leu	Lys	Arg	Glu	Asp	555
590	Ser	Ile	Leu	Leu	Asn	Pro	Gly	Pro	Arg	His	Ile	Leu	Ala	565
605	Asp	Thr	Cys	Phe	Tyr	Ile	Asn	Ile	Thr	Lys	Glu	Glu	Asn	570
620	Phe	Ile	Phe	Lys	Gln	Glu	Glu	Lys	Arg	Lys	Arg	Ala	Phe	575
635	Gly	Gln	Gly	Leu	His	Glu	Gly	Pro	Ala	Arg	Leu	Pro	Val	580
650	Ile	Ile	Ala	Ser	Met	Gly	Thr	Val	Ala	Met	Asp	Leu	Gln	585
665	Glu	His	Arg	Pro	Thr	Gln	Ser	Gly	Gly	Gly	Gly	Gly	Ser	590
680	Leu	Ala	Leu	Pro	Thr	Glu	Asn	Gly	Ser	Gly	Ser	Arg	Arg	595
695	Ile	Ala	Pro	Val	Leu	Glu	Leu	Ala	Asp	Ser	Ser	Ala	Leu	600
710	Cys	Asp	Leu	Leu	Ser	Asp	Gln	Ser	Glu	Asp	Glu	Val	Thr	605
725	Asp	Asp	Glu	Gly	Leu	Ser	Val	Val	Glu	Tyr	Val	Lys	Gly	610
740	Pro	Asn	Ser	Pro	Tyr	Ile	Gly	Ser	Ser	Pro	Thr	Leu	Cys	615
755	Leu	Pro	Val	Lys	Ala	Pro	Phe	Cys	Cys	Leu	Arg	Leu	Asp	620
770	Cys	Lys	His	Asn	Ser	Tyr	Glu	Asp	Ala	Lys	Ala	Tyr	Gly	625
785	Asn	Lys	Leu	Ile	Ile	Val	Ser	Ala	Glu	Thr	Ala	Gly	Asn	630
800	Tyr	Asn	Phe	Ile	Val	Pro	Leu	Arg	Ala	Tyr	Tyr	Arg	Ser	635
815	Glu	Leu	Asn	Pro	Ile	Val	Leu	Leu	Leu	Asp	Asn	Lys	Pro	640
830	His	Phe	Leu	Glu	Ala	Ile	Cys	Cys	Phe	Pro	Met	Val	Tyr	645
845	Glu	Gly	Ser	Val	Asp	Asn	Leu	Asp	Ser	Leu	Leu	Cln	Cys	650
860	Ile	Tyr	Ala	Asp	Asn	Leu	Val	Val	Val	Asp	Lys	Glu	Ser	655
875	Ser	Ala	Glu	Glu	Asp	Tyr	Met	Ala	Asp	Ala	Lys	Thr	Ile	660
890	Val	Gln	Thr	Met	Phe	Arg	Leu	Phe	Pro	Ser	Leu	Ser	Ile	665
905	Glu	Leu	Thr	His	Pro	Ser	Asn	Met	Arg	Phe	Met	Gln	Phe	670
920	Lys	Asp	Ser	Tyr	Ser	Leu	Ala	Leu	Ser	Lys	Leu	Glu	Lys	675
935	Arg	Glu	Asn	Gly	Ser	Asn	Leu	Ala	Phe	Met	Phe	Arg	Leu	680
950	Ala	Ala	Gly	Arg	Val	Phe	Ser	Ile	Ser	Ile	Ser	Met	Leu	685

Tyr Gln Ser Phe Val Lys Asp Tyr Met Ile Thr Ile Thr Arg Leu 950 955 960
 965 970 975
 Leu Leu Gly Leu Asp Thr Thr Pro Gly Ser Gly Tyr Leu Cys Ala 980 985 990
 Met Lys Ile Thr Gln Gly Asp Leu Trp Ile Arg Thr Tyr Gly Arg 995 1000 1005
 Leu Phe Gln Lys Leu Cys Ser Ser Ala Gln Ile Pro Ile Gly 1010 1015 1020
 Ile Tyr Arg Thr Gln Ser His Val Phe Ser Thr Ser Gln Pro His 1025 1030 1035
 Gln Leu Arg Ala Gln Ser Gln Ile Ser Val Asn Val Gln Asp Cys 1040 1045 1050
 Gln Asp Thr Arg Gln Val Lys Gly Pro Trp Gly Ser Arg Ala Gly 1055 1060 1065
 Thr Gly Gly Ser Ser Gln Gly Arg His Thr Gly Gly Asp Pro 1070 1075 1080
 Ala Gln His Pro Leu Leu Arg Arg Lys Ser Leu Gln Trp Ala Arg 1085 1090 1095
 Arg Leu Ser Arg Lys Ala Pro Lys Gln Ala Gly Arg Ala Ala 1100 1105 1110
 Ala Gln Trp Ile Ser Gln Gln Arg Leu Ser Leu Tyr Arg Arg Ser 1115 1120 1125
 Gln Arg Gln Gln Leu Ser Gln Leu Val Lys Asn Arg Met Lys His 1130 1135 1140
 Leu Gly Leu Pro Thr Thr Gly Tyr Gln Asp Val Ala Asn Leu Thr 1145 1150 1155
 Ala Ser Asp Val Met Asn Arg Val Asn Leu Gly Tyr Leu Gln Asp 1160 1165 1170
 Gln Met Asn Asp His Gln Asn Thr Leu Ser Tyr Val Leu Ile Asn 1175 1180 1185
 Pro Pro Asp Thr Arg Leu Gln Pro Ser Asp Ile Val Tyr Leu 1190 1195 1200
 Ile Arg Ser Asp Pro Leu Ala His Val Ala Ser Ser Ser Gln Ser 1205 1210 1215
 Arg Lys Ser Ser Cys Ser His Lys Leu Ser Ser Cys Asn Pro Gln 1220 1225 1230
 Thr Arg Asp Gln Thr Gln Leu 1235

<210> 25
 <211> 539
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No: 7673418CD1

<400> 25
 Met Ala Ser Ala Leu Ser Tyr Val Ser Lys Phe Lys Ser Phe Val 1 5 10 15
 Ile Leu Phe Val Thr Pro Leu Leu Leu Leu Pro Leu Val Ile Leu 20 25 30
 Met Pro Ala Lys Phe Val Arg Cys Ala Tyr Val Ile Ile Leu Met 35 40 45
 Ala Ile Tyr Trp Cys Thr Gln Val Ile Pro Leu Ala Val Thr Ser 50 55 60
 Leu Met Pro Val Leu Leu Phe Pro Leu Phe Gln Ile Leu Asp Ser 65 70 75
 Arg Gln Val Cys Val Gln Tyr Met Lys Asp Thr Asn Met Leu Phe 80 85 90
 Leu Gly Gly Leu Ile Val Ala Val Ala Val Gln Arg Trp Asn Leu

44/85

His Lys Arg Ile Ala Leu Arg Thr Leu Leu Trp Val Gly Ala Lys 95 100 105
 110 115 120
 Pro Ala Arg Leu Met Leu Gly Phe Met Gly Val Thr Ala Leu Leu 125 130 135
 Ser Met Trp Ile Ser Asn Thr Ala Thr Thr Ala Met Met Val Pro 140 145 150
 Ile Val Gln Ala Ile Leu Gln Gln Met Gln Ala Thr Ser Ala Ala 155 160 165
 Thr Gln Ala Gly Leu Leu Val Asp Lys Gly Lys Ala Lys Gln 170 175 180
 Leu Pro Ala Asn Ser Ala Val Pro Thr Thr Gly Ser Gln Val Ile 185 190 195
 Phe Gln Gly Pro Thr Leu Gly Gln Gln Asp Gln Gln Arg Lys 200 205 210
 Arg Leu Cys Lys Ala Met Thr Leu Cys Ile Cys Tyr Ala Ala Ser 215 220 225
 Ile Gly Gly Thr Ala Thr Leu Thr Gly Thr Gly Pro Asn Val Val 230 235 240
 Leu Leu Gly Gln Met Asn Gln Leu Phe Pro Asp Ser Lys Asp Leu 245 250 255
 Val Asn Phe Ala Ser Trp Phe Ala Phe Ala Phe Pro Asn Met Leu 260 265 270
 Val Met Leu Leu Phe Ala Trp Leu Trp Leu Gln Phe Val Tyr Met 275 280 285
 Arg Phe Asn Phe Lys Ser Trp Gly Cys Gly Leu Gln Ser Lys 290 295 300
 Lys Asn Gln Lys Ala Ala Leu Lys Val Leu Gln Gln Gln Tyr Arg 305 310 315
 Lys Leu Gly Pro Leu Ser Phe Ala Gln Ile Asn Val Leu Ile Cys 320 325 330
 Phe Phe Leu Leu Val Ile Leu Trp Phe Ser Arg Asp Pro Gly Phe 335 340 345
 Met Pro Gly Trp Leu Thr Val Ala Trp Val Gln Gln Arg Lys Thr 350 355 360
 Pro Phe Tyr Pro Pro Pro Leu Leu Asp Trp Lys Val Thr Gln Gln 365 370 375
 Lys Val Pro Trp Gly Ile Val Leu Leu Gly Gly Gly Phe Ala 380 385 390
 Leu Ala Lys Gly Ser Gln Ala Ser Gly Leu Ser Val Trp Met Gly 395 400 405
 Lys Gln Met Gln Pro Leu His Ala Val Pro Pro Ala Ala Ile Thr 410 415 420
 Leu Ile Leu Ser Leu Leu Val Ala Val Phe Thr Gln Cys Thr Ser 425 430 435
 Asn Val Ala Thr Thr Thr Leu Phe Leu Pro Ile Phe Ala Ser Met 440 445 450
 Ser Arg Ser Ile Gly Leu Asn Pro Leu Tyr Ile Met Leu Pro Cys 455 460 465
 Thr Leu Ser Ala Ser Phe Ala Phe Met Leu Pro Val Ala Thr Pro 470 475 480
 Pro Asn Ala Ile Val Phe Thr Tyr Gly His Leu Lys Val Ala Asp 485 490 495
 Met Val Lys Thr Gly Val Ile Met Asn Ile Ile Gly Val Phe Cys 500 505 510
 Val Phe Leu Ala Val Asn Thr Trp Gly Arg Ala Ile Phe Asp Leu 515 520 525
 Asp His Phe Pro Asp Trp Ala Asn Val Thr His Ile Gln Thr 530 535

<210> 26
 <211> 755
 <212> PRT

45/85

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7474129CD1

<400> 26

Met Lys Ala His Pro Lys Glu Met Val Pro Leu Met Gly Lys Arg
 1 5 10 15
 Val Ala Ala Pro Ser Gly Asn Pro Ala Val Leu Pro Glu Lys Arg
 20 25 30
 Pro Ala Glu Ile Thr Pro Thr Lys Lys Ser Ile Ser Gly Asn Cys
 35 40 45
 Asp Asp Met Asp Ser Pro Gln Ser Pro Gln Asp Asp Val Thr Glu
 50 55 60
 Thr Pro Ser Asn Pro Asn Ser Pro Ser Ala Gln Leu Ala Lys Glu
 65 70 75
 Glu Gln Arg Arg Lys Lys Arg Arg Leu Lys Lys Arg Ile Phe Ala
 80 85 90
 Ala Val Ser Glu Gly Cys Val Glu Glu Leu Val Glu Leu Val
 95 100 105
 Glu Leu Gln Glu Leu Cys Arg Arg Arg His Asp Glu Asp Val Pro
 110 115 120
 Asp Phe Leu Met His Lys Leu Thr Ala Ser Asp Thr Gly Lys Thr
 125 130 135
 Cys Leu Met Lys Ala Leu Leu Asn Ile Asn Pro Asn Thr Lys Glu
 140 145 150
 Ile Val Arg Ile Leu Leu Ala Phe Ala Glu Asn Asp Ile Leu
 155 160 165
 Gly Arg Phe Ile Asn Ala Glu Tyr Thr Glu Ala Tyr Glu Gly
 170 175 180
 Gln Thr Ala Leu Asn Ile Ala Ile Glu Arg Arg Gln Gly Asp Ile
 185 190 195
 Ala Ala Leu Leu Ile Ala Ala Gly Ala Asp Val Asn Ala His Ala
 200 205 210
 Lys Gly Ala Phe Asn Pro Lys Tyr Gln His Glu Gly Phe Tyr
 215 220 225
 Phe Gly Glu Thr Pro Leu Ala Leu Ala Cys Thr Asn Gln Pro
 230 235 240
 Glu Ile Val Gln Leu Leu Met Glu His Glu Gln Thr Asp Ile Thr
 245 250 255
 Ser Arg Asp Ser Arg Gly Asn Asn Ile Leu His Ala Leu Val Thr
 260 265 270
 Val Ala Glu Asp Phe Lys Thr Gln Asn Asp Val Val Lys Arg Met
 275 280 285
 Tyr Asp Met Ile Leu Leu Arg Ser Gly Asn Thr Glu Leu Glu Thr
 290 295 300
 Thr Arg Asn Asn Asp Gly Leu Thr Pro Leu Gln Leu Ala Ala Lys
 305 310 315
 Met Gly Lys Ala Glu Ile Leu Lys Tyr Ile Leu Ser Arg Glu Ile
 320 325 330
 Lys Glu Lys Arg Leu Arg Ser Leu Ser Arg Lys Phe Thr Asp Thr
 335 340 345
 Ala Tyr Gly Pro Val Ser Ser Ser Leu Tyr Asp Leu Thr Asn Val
 350 355 360
 Asp Thr Thr Asp Asn Ser Val Leu Glu Ile Thr Val Tyr Asn
 365 370 375
 Thr Asn Ile Asp Asn Arg His Glu Met Leu Thr Leu Glu Pro Leu
 380 385 390
 His Thr Leu Leu His Met Lys Trp Lys Lys Phe Ala Lys His Met
 395 400 405
 Phe Phe Leu Ser Phe Cys Phe Tyr Phe Thr Asn Ile Thr Leu
 410 415 420

46/85

Thr Leu Val Ser Tyr Arg Pro Arg Glu Glu Ala Ile Pro
 425 430 435
 His Pro Leu Ala Leu Thr His Lys Met Gly Trp Leu Gln Leu
 440 445 450
 Gly Arg Met Phe Val Leu Ile Trp Ala Met Cys Ile Ser Val Lys
 455 460 465
 Glu Gly Ile Ala Ile Phe Leu Leu Arg Pro Ser Asp Leu Gln Ser
 470 475 480
 Ile Leu Ser Asp Ala Trp Phe His Phe Val Phe Ile Gln Ala
 485 490 495
 Val Leu Val Ile Leu Ser Val Phe Leu Tyr Leu Phe Ala Tyr Lys
 500 505 510
 Glu Tyr Leu Ala Cys Leu Val Leu Ala Met Ala Leu Gly Trp Ala
 515 520 525
 Asn Met Leu Tyr Thr Arg Gly Phe Gln Ser Met Gly Met Tyr
 530 535 540
 Ser Val Met Ile Gln Lys Val Ile Leu His Asp Val Leu Lys Phe
 545 550 555
 Leu Phe Val Tyr Ile Val Phe Leu Leu Gly Phe Gly Val Ala Leu
 560 565 570
 Ala Ser Leu Ile Glu Lys Cys Pro Lys Asp Asn Lys Asp Cys Ser
 575 580 585
 Ser Tyr Gly Ser Phe Ser Asp Ala Val Leu Glu Leu Phe Lys Leu
 590 595 600
 Thr Ile Gly Leu Gly Asp Leu Asn Ile Gln Gln Asn Ser Lys Tyr
 605 610 615
 Pro Ile Leu Phe Leu Phe Leu Ile Thr Tyr Val Ile Leu Thr
 620 625 630
 Phe Val Leu Leu Asn Met Leu Ile Ala Leu Met Gly Glu Thr
 635 640 645
 Val Glu Asn Val Ser Lys Glu Ser Glu Arg Ile Trp Arg Leu Gln
 650 655 660
 Arg Ala Arg Thr Ile Leu Glu Phe Glu Lys Met Leu Pro Glu Trp
 665 670 675
 Leu Arg Ser Arg Phe Arg Met Gly Glu Leu Cys Lys Val Ala Glu
 680 685 690
 Asp Asp Phe Arg Leu Cys Leu Arg Ile Asn Glu Val Lys Trp Thr
 695 700 705
 Glu Trp Lys Thr His Val Ser Phe Leu Asn Glu Asp Pro Gly Pro
 710 715 720
 Val Arg Arg Thr Asp Phe Asn Lys Ile Gln Asp Ser Ser Arg Asn
 725 730 735
 Asn Ser Lys Thr Thr Leu Asn Ala Phe Glu Glu Val Glu Glu Phe
 740 745 750
 Pro Glu Thr Ser Val
 755

<210> 27

<211> 301

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7481414CD1

<400> 27

Met Lys Ser His Pro Ala Ile Gln Ala Ala Ile Asp Leu Thr Ala
 1 5 10 15
 Gly Ala Ala Gly Gly Ala Cys Val Leu Thr Gly Gln Pro Phe
 20 25 30
 Asp Thr Ile Lys Val Lys Met Gln Thr Phe Pro Gln Leu Tyr Lys
 35 40 45

47/85

Gly Leu Ala Asp Cys Phe Leu Lys Thr Tyr Asn Gln Val Gly Ile
50 55 60
Arg Gly Leu Tyr Arg Gly Thr Ser Pro Ala Leu Leu Ala Tyr Val
65 70 75
Thr Gln Gly Ser Val Leu Phe Met Cys Phe Gly Phe Cys Gln Gln
80 85 90
Phe Val Arg Lys Val Ala Arg Val Gln Gln Asn Ala Gln Leu Asn
95 100 105
Asp Leu Gln Thr Ala Thr Ala Gly Ser Leu Ala Ser Ala Phe Ala
110 115 120
Ala Leu Ala Leu Cys Pro Thr Gln Leu Val Lys Cys Arg Leu Gln
125 130 135
Thr Met Tyr Gln Met Lys Met Ser Gly Lys Ile Ala Gln Ser Tyr
140 145 150
Asn Thr Ile Trp Ser Met Val Lys Ser Ile Phe Met Lys Asp Gly
155 160 165
Pro Leu Gly Phe Tyr Arg Gly Leu Ser Thr Thr Leu Ala Gln Gln
170 175 180
Ile Pro Gly Tyr Phe Phe Gly Gly Tyr Gln Ile Ser Arg
185 190 195
Ser Phe Phe Ala Ser Gly Gly Ser Lys Asp Gln Leu Gly Pro Val
200 205 210
Pro Leu Met Leu Ser Gly Gly Phe Ala Gly Ile Cys Leu Trp Leu
215 220 225
Ile Ile Phe Pro Val Asp Cys Ile Lys Ser Arg Ile Gln Val Leu
230 235 240
Ser Met Phe Gly Lys Pro Ala Gly Leu Ile Gln Thr Phe Ile Ser
245 250 255
Val Val Arg Asn Gln Gly Ile Ser Ala Leu Tyr Ser Gly Leu Lys
260 265 270
Ala Thr Leu Ile Arg Ala Ile Pro Ser Asn Ala Ala Leu Phe Leu
275 280 285
Val Tyr Gln Tyr Ser Arg Lys Met Met Asn Met Val Gln Gln
290 295 300
Tyr

<210> 28

<211> 515

<212> PRT

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No: 7481461CD1

<400> 28

Met Val Leu Ser Gln Gln Gln Pro Asp Ser Ala Arg Gly Thr Ser
5 10 15
Gln Ala Gln Pro Leu Gly Pro Ala Pro Thr Gly Ala Ala Pro Pro
20 25 30
Pro Gly Pro Gly Pro Ser Asp Ser Pro Gln Ala Ala Val Gln Lys
35 40 45
Val Gln Val Gln Leu Ala Gly Pro Ala Thr Ala Gln Pro His Gln
50 55 60
Pro Pro Gln Pro Pro Gln Gly Gly Trp Gly Trp Leu Val Met Leu
65 70 75
Ala Ala Met Trp Cys Asn Gly Ser Val Phe Gly Ile Gln Asn Ala
80 85 90
Cys Gly Val Leu Phe Val Ser Met Leu Gln Thr Phe Gly Ser Lys
95 100 105
Asp Asp Asp Lys Met Val Phe Lys Thr Ala Trp Val Gly Ser Leu
110 115 120

48/85

Ser Met Gly Met Ile Phe Phe Cys Cys Pro Ile Val Ser Val Phe
125 130 135
Thr Asp Leu Phe Gly Cys Arg Lys Thr Ala Val Val Gly Ala Ala
140 145 150
Val Gly Phe Val Gly Leu Met Ser Ser Ser Phe Val Ser Ser Ile
155 160 165
Gln Pro Leu Tyr Leu Thr Tyr Gly Ile Ile Phe Ala Cys Gly Cys
170 175 180
Ser Phe Ala Tyr Gln Pro Ser Leu Val Ile Leu Gly His Tyr Phe
185 190 195
Lys Lys Arg Leu Gly Leu Val Asn Gly Ile Val Thr Ala Gly Ser
200 205 210
Ser Val Phe Thr Ile Leu Leu Pro Leu Leu Asp Val Leu Ile
215 220 225
Asp Ser Val Gly Leu Phe Tyr Thr Leu Arg Val Leu Cys Ile Phe
230 235 240
Met Phe Val Leu Leu Ala Gly Phe Thr Tyr Arg Pro Leu Ala
245 250 255
Thr Ser Thr Lys Asp Lys Gln Ser Gly Ser Gly Ser Ser Leu
260 265 270
Phe Ser Arg Lys Lys Phe Ser Pro Pro Lys Lys Ile Phe Asn Phe
275 280 285
Ala Ile Phe Lys Val Thr Ala Tyr Ala Val Trp Ala Val Gly Ile
290 295 300
Pro Leu Ala Leu Phe Gly Tyr Phe Val Pro Tyr Val His Leu Met
305 310 315
Lys His Val Asn Gln Arg Phe Gln Asp Gln Lys Asn Lys Gln Val
320 325 330
Val Leu Met Cys Ile Gly Val Thr Ser Gly Val Arg Leu Leu
335 340 345
Phe Gly Arg Ile Ala Asp Tyr Val Pro Gly Val Lys Lys Val Tyr
350 355 360
Leu Gln Val Leu Ser Phe Phe Ile Gly Leu Met Ser Met Met
365 370 375
Ile Pro Leu Cys Ser Ile Phe Gly Ala Leu Ile Ala Val Cys Leu
380 385 390
Ile Met Gly Leu Phe Asp Gly Cys Phe Ile Ser Ile Met Ala Pro
395 400 405
Ile Ala Phe Gln Leu Val Gly Ala Gln Asp Val Ser Gln Ala Ile
410 415 420
Gly Phe Leu Leu Gly Phe Met Ser Ile Pro Met Thr Val Gly Pro
425 430 435
Pro Ile Ala Gly Leu Leu Arg Asp Lys Leu Gly Ser Tyr Asp Val
440 445 450
Ala Phe Tyr Leu Ala Gly Val Pro Pro Leu Ile Gly Gly Ala Val
455 460 465
Leu Cys Phe Ile Pro Trp Ile His Ser Lys Lys Gln Arg Gln Ile
470 475 480
Ser Lys Thr Thr Tyr Lys Gln Lys Met Gln Lys Met Leu Gln Asn
485 490 495
Gln Asn Ser Leu Leu Ser Ser Ser Gly Met Phe Lys Lys Gln
500 505 510
Ser Asp Ser Ile Ile

<210> 29

<211> 1519

<212> PRT

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No: 7472541CD1

49/85

<400> 29
Met Ala Leu Ser Val Asp Ser Ser Trp His Arg Trp Gln Trp Arg 15
1 Val Arg Asp Gly Phe Pro His Cys Pro Ser Glu Thr Thr Pro Leu 20
20 Leu Ser Pro Glu Lys Gly Arg Gln Ser Tyr Asn Leu Thr Gln Gln 30
35 Arg Val Val Phe Pro Asn Asn Ser Ile Phe His Gln Asp Trp Glu 45
50 Glu Val Ser Arg Arg Tyr Pro Gly Asn Arg Thr Cys Thr Thr Lys 60
65 Tyr Thr Leu Phe Thr Phe Leu Pro Arg Asn Leu Phe Glu Gln Phe 75
80 His Arg Trp Ala Asn Leu Tyr Phe Leu Phe Leu Val Ile Leu Ser 90
95 Trp Met Pro Ser Met Glu Val Phe His Arg Glu Ile Thr Met Leu 105
110 Pro Leu Ala Ile Val Leu Phe Val Ile Met Ile Lys Asp Gly Met 120
125 Glu Asp Phe Lys Arg His Arg Phe Asp Lys Ala Ile Asn Cys Ser 135
140 Asn Ile Arg Ile Tyr Glu Arg Lys Glu Gln Thr Tyr Val Gln Lys 150
155 Cys Trp Lys Asp Val Arg Val Gly Asp Phe Ile Gln Met Lys Cys 165
170 Asn Glu Ile Val Pro Ala Asp Ile Leu Leu Leu Phe Ser Ser 180
185 Pro Asn Gly Ile Cys His Leu Glu Thr Ala Ser Leu Asp Gly Glu 195
200 Thr Asn Leu Lys Gln Arg Arg Val Val Lys Gly Phe Ser Gln Gln 210
215 Glu Val Gln Phe Glu Pro Glu Leu Phe His Asn Thr Ile Val Cys 225
230 Glu Lys Pro Asn Asn His Leu Asn Lys Phe Lys Gly Tyr Met Glu 240
245 His Pro Asp Gln Thr Arg Thr Gly Phe Gly Cys Glu Ser Leu Leu 255
260 Leu Arg Gly Cys Thr Ile Arg Asn Thr Glu Met Ala Val Gly Ile 270
275 Val Ile Tyr Ala Gly His Glu Thr Lys Ala Met Leu Asn Asn Ser 285
290 Gly Pro Arg Tyr Lys Arg Ser Lys Ile Glu Arg Arg Met Asn Ile 300
305 Asp Ile Phe Phe Cys Ile Gly Ile Leu Ile Leu Met Cys Leu Ile 315
320 Gly Ala Val Gly His Ser Ile Trp Asn Gly Thr Phe Glu Glu His 330
335 Pro Pro Phe Asp Val Pro Asp Ala Asn Gly Ser Phe Leu Pro Ser 345
350 Ala Leu Gly Gly Phe Tyr Met Phe Leu Thr Met Ile Ile Leu Leu 360
365 Gln Val Leu Ile Pro Ile Ser Leu Tyr Val Ser Ile Glu Leu Val 375
380 Lys Leu Gly Gln Val Phe Phe Leu Ser Asn Asp Leu Asp Leu Tyr 390
395 Asp Glu Glu Thr Asp Leu Ser Ile Gln Cys Arg Ala Leu Asn Ile 405
410 Ala Glu Asp Leu Gly Gln Ile Gln Tyr Ile Phe Ser Asp Lys Thr 420
425 Gly Thr Leu Thr Glu Asn Lys Met Val Phe Arg Arg Cys Thr Ile 435
440 Met Gly Ser Glu Tyr Ser His Gln Glu Asn Ala Lys Arg Leu Glu 450
455

Thr Pro Lys Glu Leu Asp Ser Asp Gly Glu Glu Trp Thr Gln Tyr 470
475 Gln Cys Leu Ser Phe Ser Ala Arg Trp Ala Gln Asp Pro Ala Thr 480
485 Met Arg Ser Gln Lys Gly Ala Gln Pro Leu Arg Arg Ser Gln Ser 495
500 Ala Arg Val Pro Ile Gln Gly His Tyr Arg Gln Arg Ser Met Gly 510
515 His Arg Glu Ser Ser Gln Pro Pro Val Ala Phe Ser Ser Ser Ile 525
530 Glu Lys Asp Val Thr Pro Asp Lys Asn Leu Leu Thr Lys Val Arg 540
545 Asp Ala Ala Leu Trp Leu Glu Thr Leu Ser Asp Ser Arg Pro Ala 555
560 Lys Ala Ser Leu Ser Thr Thr Ser Ser Ile Ala Asp Phe Phe Leu 570
575 Ala Leu Thr Ile Cys Asn Ser Val Met Val Ser Thr Thr Thr Glu 585
590 Pro Arg Gln Arg Trp Asp Asp Gln Lys Ile Val Glu Asn Asp His 600
605 Cys Gln Cys Leu Glu Phe Gln Gly Trp Arg Lys Ile Ser Gly Phe 615
620 Thr Tyr Cys Lys Ser Thr Phe Ile Phe Arg Ile Arg Gln Leu Gly 630
635 Ile Ile Ser Asn Ile Glu Ser Asn Ile Pro Leu Ser Phe Phe Gly 645
650 His Lys Val Thr Ile Lys Pro Ser Ser Lys Ala Leu Gly Thr Ser 660
665 Leu Glu Lys Ile Gln Gln Leu Phe Gln Lys Leu Lys Leu Ser 675
680 Leu Ser Gln Ser Phe Ser Ser Thr Ala Pro Ser Asp Thr Asp Leu 690
695 Gly Glu Ser Leu Gly Ala Asn Val Ala Thr Thr Asp Ser Asp Glu 705
710 Arg Asp Asp Ala Ser Val Cys Ser Gly Gly Asp Ser Thr Asp Asp 720
725 Gly Gly Tyr Arg Ser Ser Met Trp Asp Gln Gly Asp Ile Leu Glu 735
740 Ser Gly Ser Gly Thr Ser Leu Glu Glu Ala Leu Glu Ala Pro Ala 750
755 Thr Asp Leu Ala Arg Pro Glu Phe Cys Tyr Glu Ala Glu Ser Pro 765
770 Asp Glu Ala Ala Leu Val His Ala His Ala Tyr Ser Phe Thr 780
785 Leu Val Ser Arg Thr Pro Glu Gln Val Thr Val Arg Leu Pro Gln 795
800 Gly Thr Cys Leu Thr Phe Ser Leu Leu Cys Thr Leu Gly Phe Asp 810
815 Ser Val Arg Lys Arg Met Ser Val Val Val Arg His Pro Leu Thr 825
830 Gly Glu Ile Val Val Tyr Thr Lys Gly Ala Asp Ser Val Ile Met 840
845 Asp Leu Leu Glu Asp Pro Ala Cys Val Pro Asp Ile Asn Met Glu 855
860 Lys Lys Leu Arg Lys Ile Arg Ala Arg Thr Gln Lys His Leu Asp 870
875 Leu Tyr Ala Arg Asp Gly Leu Arg Thr Leu Cys Ile Ala Lys Lys 885
890 Val Val Ser Glu Glu Asp Phe Arg Arg Trp Ala Ser Phe Arg Arg 900
905 Glu Ala Glu Ala Ser Leu Asp Asn Arg Asp Glu Leu Leu Met Glu 915
920 Thr Ala Gln His Leu Glu Asn Gln Leu Thr Leu Leu Gly Ala Thr 930

Gly Ile Glu Asp Arg Leu Gln Glu Gly Val Pro Asp Thr Ile Ala 935 940 945
 Thr Leu Arg Glu Ala Gly Ile Gln Leu Trp Val Leu Thr Gly Asp 950 955 960
 Lys Gln Glu Thr Ala Val Asn Ile Ala His Ser Cys Arg Leu Leu 965 970 975
 Asn Gln Thr Asp Thr Val Tyr Thr Ile Asn Thr Glu Asn Gln Glu 980 985 990
 Thr Cys Glu Ser Ile Leu Asn Cys Ala Leu Glu Leu Lys Gln 995 1000 1005
 Phe Arg Glu Leu Gln Lys Pro Asp Arg Lys Leu Phe Gly Phe Arg 1010 1015 1020
 Leu Pro Ser Lys Thr Pro Ser Ile Thr Ser Glu Ala Val Val Pro 1025 1030 1035
 Glu Ala Gly Leu Val Ile Asp Gly Lys Thr Leu Asn Ala Ile Phe 1040 1045 1050
 Gln Gly Lys Leu Glu Lys Lys Phe Leu Glu Leu Thr Gln Tyr Cys 1055 1060 1065
 Arg Ser Val Leu Cys Cys Arg Ser Thr Pro Leu Gln Lys Ser Met 1070 1075 1080
 Ile Val Lys Leu Val Arg Asp Lys Leu Arg Val Met Thr Leu Ser 1085 1090 1095
 Ile Gly Asp Gly Ala Asn Asp Val Ser Met Ile Gln Ala Ala Asp 1100 1105 1110
 Ile Gly Ile Gly Ile Ser Gly Gln Glu Gly Met Gln Ala Val Met 1115 1120 1125
 Ser Ser Asp Phe Ala Ile Thr Arg Phe Phe His Leu Lys Lys Leu 1130 1135 1140
 Leu Leu Val His Gly His Trp Cys Tyr Ser Arg Leu Ala Arg Met 1145 1150 1155
 Val Val Tyr Tyr Leu Tyr Lys Asn Val Cys Tyr Val Asn Leu Leu 1160 1165 1170
 Phe Trp Tyr Gln Phe Phe Cys Gly Phe Ser Ser Thr Met Ile 1175 1180 1185
 Asp Tyr Trp Gln Met Ile Phe Phe Asn Leu Phe Thr Ser Leu 1190 1195 1200
 Pro Pro Leu Val Phe Gly Val Leu Asp Lys Asp Ile Ser Ala Glu 1205 1210 1215
 Thr Leu Leu Ala Leu Pro Glu Leu Tyr Lys Ser Gly Gln Asn Ser 1220 1225 1230
 Glu Cys Tyr Asn Leu Ser Thr Phe Trp Ile Ser Met Val Asp Ala 1235 1240 1245
 Phe Tyr Gln Ser Leu Ile Cys Phe Phe Ile Pro Tyr Leu Ala Tyr 1250 1255 1260
 Lys Gly Ser Asp Ile Asp Val Phe Thr Phe Gly Thr Pro Ile Asn 1265 1270 1275
 Thr Ile Ser Leu Thr Thr Ile Leu Leu His Gln Ala Met Glu Met 1280 1285 1290
 Lys Thr Trp Thr Ile Phe His Gly Val Val Leu Glu Gly Ser Phe 1295 1300 1305
 Leu Met Tyr Phe Leu Val Ser Leu Leu Tyr Asn Ala Thr Cys Val 1310 1315 1320
 Ile Cys Asn Ser Pro Thr Asn Pro Tyr Trp Val Met Glu Gly Gln 1325 1330 1335
 Leu Ser Asn Pro Thr Phe Tyr Leu Val Cys Phe Leu Thr Pro Val 1340 1345 1350
 Val Ala Leu Leu Leu Arg Tyr Phe Phe Leu Ser Leu Gln Gly Thr 1355 1360 1365
 Cys Gly Lys Ser Leu Ile Ser Lys Ala Gln Lys Ile Asp Lys Leu 1370 1375 1380
 Pro Pro Asp Lys Arg Asn Leu Glu Ile Gln Ser Trp Arg Ser Arg 1385 1390 1395
 1400 1405 1410

52/85

Gln Arg Pro Ala Pro Val Pro Glu Val Ala Arg Pro Thr His His 1415 1420 1425
 Pro Val Ser Ser Ile Thr Gly Gln Asp Phe Ser Ala Ser Thr Pro 1430 1435 1440
 Lys Ser Ser Asn Pro Pro Lys Arg Lys His Val Glu Glu Ser Val 1445 1450 1455
 Leu His Glu Gln Arg Cys Gly Thr Glu Cys Met Arg Asp Ser 1460 1465 1470
 Cys Ser Gly Asp Ser Ser Ala Gln Leu Ser Ser Gly Glu His Leu 1475 1480 1485
 Leu Gly Pro Asn Arg Ile Met Ala Tyr Ser Gly Gly Gln Thr Asp 1490 1495 1500
 Met Cys Arg Cys Ser Lys Arg Ser Ser His Arg Arg Ser Gln Ser 1505 1510 1515
 Ser Leu Thr Ile

 <210> 30
 <211> 1585
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> mlec_feature
 <223> Incyte ID No: 6999183CD1

 <400> 30
 Met Ser Lys Arg Arg Met Ser Val Gly Gln Gln Thr Trp Ala Leu 5 10 15
 1 Leu Cys Lys Asn Cys Leu Lys Lys Trp Arg Met Lys Arg Gln Thr 20 25 30
 Leu Leu Glu Trp Leu Phe Ser Phe Leu Leu Val Leu Phe Leu Tyr 35 40 45
 Leu Phe Phe Ser Asn Leu His Gln Val His Asp Thr Pro Gln Met 50 55 60
 Ser Ser Met Asp Leu Gly Arg Val Asp Ser Phe Asn Asp Thr Asn 65 70 75
 Tyr Val Ile Ala Phe Ala Pro Glu Ser Lys Thr Thr Gln Glu Ile 80 85 90
 Met Asn Lys Val Ala Ser Ala Pro Phe Leu Met Ala Gly Arg Thr 95 100 105
 Ile Met Gly Trp Pro Asp Glu Lys Ser Met Asp Glu Leu Asp Leu 110 115 120
 Asn Tyr Ser Ile Asp Ala Val Arg Val Ile Phe Thr Asp Thr Phe 125 130 135
 Ser Tyr His Leu Lys Phe Ser Trp Gly His Arg Ile Pro Met Met 140 145 150
 Lys Glu His Arg Asp His Ser Ala His Cys Gln Ala Val Asn Glu 155 160 165
 Lys Met Lys Cys Glu Gly Ser Glu Phe Trp Glu Lys Gly Phe Val 170 175 180
 Ala Phe Gln Ala Ala Ile Asn Ala Ala Ile Ile Glu Ile Ala Thr 185 190 195
 Asn His Ser Val Met Glu Gln Leu Met Ser Val Thr Gly Val His 200 205 210
 Met Lys Ile Leu Pro Phe Val Ala Gln Gly Gly Val Ala Thr Asp 215 220 225
 Phe Phe Ile Phe Cys Ile Ile Ser 235
 Tyr Val Ser Val Asn Val Thr Gln Glu Arg Gln Tyr Ile Thr Ser 240 245 250
 Leu Met Thr Met Met Gly Leu Arg Glu Ser Ala Phe Trp Leu Ser 255 260 265 270

53/85

Trp Gly Leu Met Tyr Ala Gly Phe Ile Leu Ile Met Ala Thr Leu 285
275 Val Val Lys Ser Ala Gln Ile Val Val Leu Thr Gly 285
Met Ala Leu Ile Val Lys Ser Ala Gln Ile Val Val Leu Thr Gly 300
Phe Val Met Val Phe Thr Leu Phe Leu Tyr Tyr Gly Leu Ser Leu 315
305 Ile Thr Leu Ala Phe Leu Met Ser Val Leu Ile Lys Lys Pro Phe 330
320 Leu Thr Gly Leu Val Val Phe Leu Ile Val Phe Trp Gly Ile 345
335 Leu Gly Phe Pro Ala Leu Tyr Thr His Leu Pro Ala Phe Leu Glu 350
350 Trp Thr Leu Cys Leu Leu Ser Pro Phe Ala Phe Thr Val Gly Met 360
365 Ala Gln Leu Ile His Leu Asp Tyr Asp Val Asn Ser Asn Ala His 375
380 Leu Asp Ser Ser Gln Asn Pro Tyr Leu Ile Ile Ala Thr Leu Phe 390
395 Met Leu Val Phe Asp Thr Leu Leu Tyr Leu Val Leu Thr Leu Tyr 405
410 Phe Asp Lys Ile Leu Pro Ala Glu Tyr Gly His Arg Cys Ser Pro 420
425 Leu Phe Phe Leu Lys Ser Cys Phe Trp Phe Gln His Gly Arg Ala 435
440 Asn His Val Val Leu Glu Asn Glu Thr Asp Ser Asp Pro Thr Pro 450
455 Asn Asp Cys Phe Glu Pro Val Ser Pro Glu Phe Cys Gly Lys Glu 465
470 Ala Ile Arg Ile Lys Asn Leu Lys Lys Glu Tyr Ala Gly Lys Cys 480
485 Glu Arg Val Glu Ala Leu Lys Gly Val Val Phe Asp Ile Tyr Glu 495
500 Gly Gln Ile Thr Ala Leu Leu Gly His Ser Gly Ala Gly Lys Thr 510
515 Thr Leu Leu Asn Ile Leu Ser Gly Leu Ser Val Pro Thr Ser Gly 525
530 Ser Val Thr Val Tyr Asn His Thr Leu Ser Arg Met Ala Asp Ile 540
545 Glu Asn Ile Ser Lys Phe Thr Gly Phe Cys Pro Gln Ser Asn Val 555
560 Gln Phe Gly Phe Leu Thr Val Lys Glu Asn Leu Arg Leu Phe Ala 570
575 Lys Ile Lys Gly Ile Leu Pro His Glu Val Glu Lys Glu Val Leu 585
590 Leu Leu Asp Glu Pro Thr Ala Gly Leu Asp Pro Leu Ser Arg His 600
605 Arg Ile Trp Asn Leu Leu Lys Glu Gly Lys Ser Asp Arg Val Ile 615
620 Leu Phe Ser Thr Gln Phe Ile Asp Glu Ala Asp Ile Leu Ala Asp 630
635 Arg Lys Val Phe Ile Ser Asn Gly Lys Leu Lys Cys Ala Gly Ser 645
650 Ser Leu Phe Leu Lys Lys Tyr Gly Ile Gly Tyr His Leu Ser 660
665 Leu His Leu Asn Glu Arg Cys Asp Pro Glu Ser Ile Thr Ser Leu 675
680 Val Lys Gln His Ile Ser Asp Ala Lys Leu Thr Ala Gln Ser Glu 690
695 Glu Lys Leu Val Tyr Ile Leu Pro Leu Glu Arg Thr Asn Lys Phe 705
710 Pro Glu Leu Tyr Arg Asp Leu Asp Arg Cys Ser Asn Gln Gly Ile 720
725 Glu Asp Tyr Gly Val Ser Ile Thr Thr Leu Asn Glu Val Phe Leu 735

740 Lys Leu Glu Gly Lys Ser Thr Ile Asp Glu Ser Asp Ile Gly Ile 750
755 Trp Gly Gln Leu Gln Thr Asp Gly Ala Lys Asp Ile Gly Ser Leu 765
770 Val Glu Leu Glu Gln Val Leu Ser Ser Phe His Glu Thr Arg Lys 780
785 Thr Ile Ser Gly Val Ala Leu Trp Arg Gln Gln Val Cys Ala Ile 795
800 Ala Lys Val Arg Phe Leu Lys Leu Lys Glu Arg Lys Ser Leu 810
815 Trp Thr Ile Leu Leu Leu Phe Gly Ile Ser Phe Ile Pro Gln Leu 825
830 Leu Glu His Leu Phe Tyr Glu Ser Tyr Gln Lys Ser Tyr Pro Trp 840
845 Glu Leu Ser Pro Asn Thr Tyr Phe Leu Ser Pro Gly Gln Gln Pro 855
860 Gln Asp Pro Leu Thr His Leu Val Ile Asn Lys Thr Gly Ser 870
875 Thr Ile Asp Asn Phe Leu His Ser Leu Arg Arg Gln Asn Ile Ala 885
890 Ile Glu Val Asp Ala Phe Gly Thr Arg Asn Gly Thr Asp Asp Pro 900
905 Ser Tyr Asn Gly Ala Ile Ile Val Ser Gly Asp Glu Lys Asp His 915
920 Arg Phe Ser Ile Ala Cys Asn Thr Lys Arg Leu Asn Cys Phe Pro 930
935 Val Leu Leu Asp Val Ile Ser Asn Gly Leu Leu Gly Ile Phe Asn 945
950 Ser Ser Glu His Ile Gln Thr Asp Arg Ser Thr Phe Phe Glu Glu 960
965 His Met Asp Tyr Glu Tyr Gly Tyr Arg Ser Asn Thr Phe Phe Tyr 975
980 Ile Pro Met Ala Ala Ser Phe Phe Thr Pro Tyr Ile Ala Met Ser Ser 990
1000 Ile Gly Asp Tyr Lys Lys Lys Ala His Ser Gln Leu Arg Ile Ser 1005
1010 Gly Leu Tyr Pro Ser Ala Tyr Trp Phe Gly Gln Ala Leu Val Asp 1020
1025 Val Ser Leu Tyr Phe Leu Ile Leu Leu Leu Met Gln Ile Met Asp 1035
1040 Tyr Ile Phe Ser Pro Glu Glu Ile Ile Phe Ile Ile Gln Asn Leu 1050
1055 Leu Ile Gln Ile Leu Cys Ser Ile Gly Tyr Val Ser Ser Leu Val 1065
1070 Phe Leu Thr Tyr Val Ile Ser Phe Ile Phe Arg Asn Gly Arg Lys 1080
1085 Asn Ser Gly Ile Trp Ser Phe Phe Leu Ile Val Val Ile Phe 1095
1100 Ser Ile Val Ala Thr Asp Leu Asn Glu Tyr Gly Phe Leu Gly Leu 1110
1115 Phe Phe Gly Thr Met Leu Ile Pro Pro Phe Thr Leu Ile Gly Ser 1125
1130 Leu Phe Ile Phe Ser Glu Ile Ser Pro Asp Ser Met Asp Tyr Leu 1140
1145 Gly Ala Ser Glu Ser Glu Ile Val Tyr Leu Ala Leu Leu Ile Pro 1155
1160 Tyr Leu His Phe Leu Ile Phe Leu Phe Ile Leu Arg Cys Leu Glu 1170
1175 Met Asn Cys Arg Lys Lys Leu Met Arg Lys Asp Pro Val Phe Arg 1185
1190 Ile Ser Pro Arg Ser Asn Ala Ile Phe Pro Asn Pro Glu Glu Pro 1200
1205 1210 55/85

Glu Gly Glu Glu Asp Ile Gln Met Glu Arg Met Arg Thr Val
1220 1225 1230
Asn Ala Met Ala Val Arg Asp Phe Asp Glu Thr Pro Val Ile Ile
1235 1240 1245
Ala Ser Cys Leu Arg Lys Glu Tyr Ala Gly Lys Lys Asn Cys
1250 1255 1260
Phe Ser Lys Arg Lys Thr Ile Ala Thr Arg Asn Val Ser Phe
1265 1270 1275
Cys Val Lys Lys Glu Val Ile Gly Leu Glu His Asn Gly
1280 1285 1290
Ala Gly Lys Ser Thr Thr Ile Lys Met Ile Thr Gly Asp Thr Lys
1295 1300 1305
Pro Thr Ala Gly Gln Val Ile Leu Lys Gly Ser Gly Gly Glu
1310 1315 1320
Pro Leu Gly Phe Leu Gly Tyr Cys Pro Gln Glu Asn Ala Leu Tyr
1325 1330 1335
Pro Asn Leu Thr Val Arg Gln His Leu Glu Val Tyr Ala Ala Val
1340 1345 1350
Lys Gly Leu Arg Lys Gly Asp Ala Met Ile Ala Ile Thr Arg Leu
1355 1360 1365
Val Asp Ala Leu Lys Leu Gln Asp Gln Leu Lys Ala Pro Val Lys
1370 1375 1380
Thr Leu Ser Glu Gly Ile Lys Arg Lys Leu Cys Phe Val Leu Ser
1385 1390 1395
Ile Leu Gly Asn Pro Ser Val Val Leu Leu Asp Glu Pro Ser Thr
1400 1405 1410
Gly Met Asp Pro Glu Gly Gln Gln Met Tyr Gln Val Ile Arg
1415 1420 1425
Ala Thr Phe Arg Asn Thr Glu Arg Gly Ala Leu Leu Thr Thr His
1430 1435 1440
Tyr Met Ala Glu Ala Glu Ala Val Cys Asp Arg Val Ala Ile Met
1445 1450 1455
Val Ser Gly Arg Lys Arg Cys Ile Gly Ser Ile Gln His Leu Lys
1460 1465 1470
Ser Lys Phe Gly Lys Asp Tyr Leu Leu Leu Met Lys Leu Lys Asn
1475 1480 1485
Leu Ala Gln Met Glu Pro Leu His Ala Glu Ile Leu Arg Leu Phe
1490 1495 1500
Pro Gln Ala Ala Gln Gln Glu Arg Phe Ser Ser Leu Met Val Tyr
1505 1510 1515
Lys Leu Pro Val Gln Asp Val Arg Pro Leu Ser Gln Ala Phe
1520 1525 1530
Lys Leu Glu Ile Val Lys Gln Ser Phe Asp Leu Glu Glu Tyr Ser
1535 1540 1545
Leu Ser Gln Ser Thr Leu Glu Val Phe Leu Glu Leu Ser Lys
1550 1555 1560
Glu Gln Glu Leu Gly Asp Leu Glu Asp Phe Asp Pro Ser Val
1565 1570 1575
Lys Tyr Lys Leu Leu Leu Gln Glu Glu Pro
1580 1585

<210> 31
<211> 1129
<212> DNA
<213> Homo sapiens

<220>
<221> misc.feature
<223> InCyte ID No: 2194064CB1

<400> 31
ggcgccgacg ccttcggat aaacgggctg tccatgggac tgcgcgcgc gctgggacct 60
ggcttcctcg accttcgcga gcaattcgac cgaacgcgcc aggcactcgc gtggatcagc 120

ggccttcgcc tggcgtcgca gcaagcagcc agccctcttg gcaagccct gagacgcgc 180
tggggggccc gccccgttgg tgatgtcttg ggggtctcgc gccctcgcgg gcttcgacct 240
ctcggtcttc gccagcgatg tglctcatct ctaccctgcg ctgggtccctc tggatggctt 300
tggltgggccc ctgggtctcg ccccgccctt aggcaccttc tgggtctact tcccgccgc 360
tccagctctg gcgttggggc tggcgctcac cggcaacggg gccctccgcgc tccctcggc 420
ggcccgcttg caactcttc tggatctt cggctcgccg ggcgtctcgc tccctccgc 480
cggatcaccc ctccacctca cccctctctg ggcctctgct ctacccttg tccctccgc 540
agaccctcca gcccccgc gtatcgcc agctccctc agtctggatc tggatcaacg 600
ccgggtcttc tcaatcttg ctctagagac agccttgatc ggggtggggtt acttcgtcc 660
ttagcgcac tggctccccc agcttggac ccggggcttg gggggatagc gggagcgct 720
gggtgctgccc gtggctcgca tgggtatgca gggcgccgc ctggtctcg gtgtgtctgc 780
agaccagatc tgggtgcccc tcccgcgctt gctggcgcta tccgggtctc tgaatggct 840
ggggctgttg gtgttggggc tggltggcgt ggttgggcggc gaaagagatc gggggggctc 900
ctcgctggcc gccgtctgag ctatggctt gggcgccggc agttaagccc ccgttggttt 960
cgtgtgtacc cccgggtctg tgggtctcg aggttggtg ctggaccacg ggcctggatc 1020
gatcgatag agccttggg gtctctcggg cccctccctg tcaagttaaag accgtgactc 1080
acagatctgc ctactcatct ctctcgccc tgggttctga ggtctctca 1129

<210> 32
<211> 2699
<212> DNA
<213> Homo sapiens

<220>
<221> misc.feature
<223> InCyte ID No: 2744094CB1

<400> 32
agtgctctgg aaagtgtat tatattatc aaacttgac ctgcagggac tgcataaggt 60
gaactaact ctctagaac ctgcctcttg ccagcgactg agctgagac ggtctagatc tccacaag 120
ctatataca tcaacctcag catgctctg aagctggua ggtatccat attcaatagg 180
taagaagag gtatgtctgt gaacttgtt acctatcca agtctagac gtttggagtt 240
ggtagagcca ggaattggaac ccaggtctct ctgactcttg aaccagatcc tcccatctc 300
tccaaagtgt ccaactggg tgggtctgca ggcctatcca ggggttggaa aagaagatgg 360
ccggagactt gattcagag gtctctcgca cctgtttgt gactctgac atcaacctgt 420
tccctccccc aggtctacag gtctctcgca taatttctct ctltgttggt ccgaatctg 480
ctttaagat cttaagctc tggtygtcga gtacttggtt ggaagagggc tgggtgacca 540
atagacagcg agagagaat ggaacatgg caagcttggg caaatctgca gaaatctcc 600
naacttctct ctacagctg gttttagggc tcaagacctt gctctctatc tgttggggg 660
tctgtctccc aaggtcttc acctagaagc ctgagggc atccagggc ctgcacaaca 720
agctcttcaa caaggtcttc cgtctgcca tgaattctt tgaacacatc ccaataggcc 780
ggcttctgaa ctgtctcgca gggagcttg aacagcttgg caagctcttg ccaatcttt 840
caggagcagt ctgtgtctg tcccttaagg tgaatgcgct cctgttgat gtcaatgtggc 900
tgcttcacaa tatccgtta atggagacca taatcatgtt tatctgcttc attatata 960
tgatgttcaa gaagccatc ggtctgttca agagacttga gaactatagc ggtctctct 1020
tatctccca catctccat tctctgcaag gctctgagct catccatgtc tatggaaaa 1080
ctgaagact catctccag tttaagaggc tgaatgagtc gaagatatac taactgtgtg 1140
tggttctaac ttcacaacga tggatggcat tgaagcttga gatcatggac aactctgtg 1200
ccttggtctg tgcctgttc gtgtcttgg gcaattcttc caaccctcac tcccttaag 1260
tcaatgtctg caaatctgtg ctgcagcttg gtttcaagctt caagcgact gcccgatgt 1320
gcttggagac agagagatcag ttacggctg taagaagat actgagatc atgaagatgt 1380
gtgtctcgga agtctctta caatlgaaag gcaacagttg tcccaagggg tggccaagtc 1440
atggggaat catattctac gatctacaa tgaatatagc agacaacga cccacgttgc 1500
ttcagatct caactggac atccgggccc agaatgtgt gggatctgt ggaagagcgg 1560
gctccgggaa gtccctctg ggtatggccc tcttcgctt ggttggagccc atggcaagc 1620
ggatcttcat tgaagcggtg gacatttga gcatcgctt ggaagatc cgttccaagc 1680
tctcagatg cccctcaagat ccagtgctg tctccagagc catcagatct aactcagatc 1740
cctttggcg tccaatctag ccaagatatt ggaatggctt ggaagagcca tctcgacca 1800
agaccatccc aaagtccccc aaaaagctgc atcaagatgt ggttgaabaac gtttgatact 1860
tctctgttgg ggaagagcag ctgctctgca ttgccaaggg tgttgcttgc aacttcaaga 1920
tcactcttat cgaatgaagc acagcttcaa ttgaabaagg taagagaccc ctgaatcgat 1980
gaaataacgg tgaagcttc caaggtctga ccgtgtcgtt catctccac cgtgtcacca 2040
ctgtgcctgaa ctgtgacgcg atccgtgta tgggaabagc gaagtgttga gatttgatc 2100

ggcggagggt actcggggaag aagcctgggt catgtttcgc agccctcatg gccacagcca 2160
 cttcttcaat gggatagaga gatgtgaga cttcatggag gcttgcagct gagctcagag 2220
 gtccacagag gtgcagcttc ggggccaca gctgcgaccc tctgtttg gagatagaga 2280
 cttctctgg agcaggggt aatgtaggg ggggtgggga tgcgtggag gaaacctgg 2340
 aataggctac tggatggctc tcaagacct tcaagcccg aacctataa gacatgggat 2400
 t-cagtgata tgggtgtctc atttaactc atagctgaa taagtgttaa 2460
 agctatagat tttctgacat gcttagaag gtttgcaat gctgtactga 2520
 tataaactga aggaanaact actttcttg tctgccttcc tttgtttct cttttttaa 2580
 tttttttaga cagggttttg atttttgc caggctggag gggatgggg caatctcagc 2640
 tcaetgagc cctgcctcc cagggttaacg caattctctt gctcagact cctgaatag 2699

<210> 33

<211> 6369

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Ineye ID No: 2798241CB1

<400> 33

tgccacaca ccagctcaat ttttatatgt ttagtagaga cagggtttca coagtgtgt 60
 caggctgggc tcaactcct gacttoggta tctgccacc ttgcctccc aaagtgtgtg 120
 gattcacagg gtgcagccc gccccgggic agctatttcc tacatgttc atttgcagt 180
 taactatgga ttgatagaga ctttgggttt ttgttaata cttcacagaa atgtgatcat 240
 tttctcttag caggcttga accaggtttg gttcaggtca taagtttcta cccactttat 300
 ttgaactctga gtgtcattt tagtttatt ttcaanaat ttgcgactac ctttttgctc 360
 tgtctgtgt gtgccttga gtaacagctc tggatttga cagttgtctg ttttttggtt 420
 cagttttcta agcctttgc acactaatag gatggatttt atgtatgtcc agcttgggaa 480
 ttatcacagg aattaaaac aactttttag agtgccttcc ttagactctt tttctttgt 540
 tcccctttt actttttgt tccctgtgyc tttgtttct ttctctcagc cagagagctta 600
 gtgtttatt tctctatgt ttctacatct tgtgcagctgt caaccctat atccaggctc 660
 caa tgggtgg aggtcagagaa gaacagaaa agggattggc cttcatctct tacaacgata 720
 gtctcatatga atagagagaa aggttttctt gctctcaggt gttgccttga ctagcttttt 780
 gttaacttag tctggccctg ttaactatgg atctcaggat ttgtgggata caggagaaat 840
 cagaaagaaa aaaaagattt gctatttcta cttcttccct gactcttaag actctccctg 900
 cccatctccc aattcaagg caaggttctt tctggagacct cctctgtggg cgtttgggaa 960
 gatacacaag gagaaaagt accactgttg atatgttgt atttcaatt tggttctacc 1020
 ctatttcaaa tgcctgtttt acttttcaag gctgcagat ttgtgtcca gcatcttgt 1080
 ccagttctct aagagagaca gcttggagta tgtctaatcc atcttactg tgcctgaac 1140
 agctgtctat ttggccgtta aatattcaat gcagtttact gctgtgctcc ggtttgttt 1200
 gtttgttttt cctctttaat aggttttattc agaaaacatg tccactgcaa ttggggagt 1260
 aggtgttttg agacagacca gaacacttct actgaagaat tacttaata aatgcagaa 1320
 caaaaaggt agtgttcagg aactttttt tccactatt tttttattt ggttaattt 1380
 aattagcatg atgcatacaa taagaanaa tgaagaagtg cctaataag aactcaatcc 1440
 tatggacaag ttactcttt ctaatactat tcttggatat actccagtgta ctaattatc 1500
 aagcagcatc atgcagaag tttctactga tcatctacct gatgtcataa ttactgaaga 1560
 atatacaaat gaacagaaa tgttaacatc cagctctctt aagccagaca acttttagg 1620
 ttgtgttttc aagaactcaa tgcctatga actctgttt ttctcgata tgcctccagt 1680
 atctctctat tatatgatt caagagcttg cttgtcaaaa tcatgtgag tgcctcagta 1740
 gttgcctcta gttttcacag ttttaacagc atctatagat gctgcctatta tacaagttaa 1800
 gaccaatgt tcttttga aggtgttga gtaacataa gctgttatta tggngagaac 1860
 tgcgttgtga gaatatgata cttttcccg agagtaatt ttaataacc tagttatagc 1920
 altttcaact tttagatata ttttggcaat tcaatactga gcgaanaag aaaaanaat 1980
 aaaaagatt tttaagataa tgggaactta tgaactgcga ttttggcttt cttgggtttc 2040
 tctatataca atgcagaag ttttatgic tcatctacct gactctatt gpaacagctc 2100
 ttgtttattt cctcaagla gcagcatgt gatattctg cttttttcc tttatggat 2160
 atcatctgta ttttttgtt taatgtgnc accttttt aaaaaataca aacatgtggg 2220
 aatagtga ttttttgtt ctggtgcttt tggattttt ggcctatga taaactctat 2280
 agaaagttt cccaattgt tagtgggttt ttccagttt tctcagctt gtaatttct 2340
 gatgtgatt gccaggtga tgaatttga aacaaagga gaagataat agtttggaa ctttaacag 2400
 ttgtcgtga ggcacatac ctttaattat tacaattat agtctcac ttaatagat 2460
 attctatgic ctttggctg tctatctga tcaagtcatt ccagggggat ttggtttct 2520

gagatcatc ttatatctt tgaagcttc atattgttca aagagcaaaa gaaattatga 2580
 gggattatca gaggccaatg ttaatggaaa tattagtctt agtgaatta ttgagccagt 2640
 ttttcagaa ttgttagga aagaagccat agaatattgt ggtatctaga agacatacag 2700
 aagaaggggt gaatagttgg aggttttgg aattttgtca ttgtacatat atgaggggtca 2760
 tgaactctg ttaactggcc acagtggaa acgttaagat gacattatga atattctttg 2820
 ttgactctgc ccaactctg atgtgttgc atatatatc agacacagag tctcagaatt 2880
 agatgaaag tttagaagaa gaaaaatgat tggctattgt ccaagtttag atatacact 2940
 tgtatgttt acgttagaag aaaaatttat aattttgtct tcaatcaaa ggtaccagc 3000
 caaataata atctagaag tgcagaaggt ttactagat ttgactatgc agactatca 3060
 agataacca gtaaaaaat taagtgttg tcaaaaaa agctgtcat taggaattgc 3120
 ttttttggg aaccaaga tactgtct agatgaaca acagtggaa tggacctct 3180
 ttctgacat atgttatgga atcttttaa atacagaana gccaatctggg tgaactgttt 3240
 cagtaactat tcaatggat agctgacat tcttcagat aggaagctg tgaataca 3300
 aggaatgtg aatgtgttg gtcttcaat gttctcaaa agtaaatgg ggtatggcta 3360
 ccgctgagc atgtacatag acaaatatg tgcacagaa tctcttctt cactgtttaa 3420
 acaacatata cctggagata cttttatac aagatagac caacaactg tttacagct 3480
 gctctcaag gactggaca aattttcag ttgtttctt gccctagca gtaataaaa 3540
 ttgtgtgtc atttcttat ggttttccat gacacttgg gaagcgtat ttttaagct 3600
 agaagtga gcagaaattg accaagcaga ttatagtga ttactcagc agccactgga 3660
 ggaagaatg gattcaaat ctttgatga aatggaacag agcttaacta tttcttga 3720
 aaccaaggt tcttctatga gaaccatgag ctttggaaa caacagatgt atacaatgc 3780
 aagtttctat tcttctact tgaacatgga aagtaaatca gtgcagatgt ttgtttctt 3840
 gcttttaatt tcttcaag ttcagatttt tatgttttg gttctact ctttcaaaa 3900
 ttgtgtgtt cccatcaaac ttgttcaga cttatattt ctaaaacctg gagacaaac 3960
 acataaatc aaaaagctc tgccttcta aattctgct gactcagata tcaatgtact 4020
 tatagactt ttcaaacgc agacataat ggtgcgatg attaatgca tgcactatgt 4080
 atccgtgtt cccatctg cggctttaa ttgtgtgcat tcaaaaaag actatgttt 4140
 tgaagttga tttcaagta ctatgtttta tctttacct attatagta gttacttag 4200
 taactactat ctttaacta taaatgtgac tgaacacatc cagatctgga taccacct 4260
 ctttcaaga atactgata tagttttaa aattgactg tatttcaag cagcttctc 4320
 ttgaactt gtactgcaa tgcacctta ctttgcatg gaaatcgag agactatca 4380
 caaagatgt tacaatac ttaaaccttc aggtctttg ccatctgat attgtatgg 4440
 acaagttgt ttgtatctc cttattttt tactactct attttgatc taggaactt 4500
 attgcatct catatagat tatatttta tcatgaag ttcttctgt tggtttttg 4560
 ccttatgtg tagtttcat cagttattt gttctattt attgtctct tccactttaa 4620
 gaaaattta taacaaaag aattttgtc attatctat tcttggcag cgttgggttg 4680
 tattgaatc actgaataa cttttttat gggatacaaa atttgaata tttctatta 4740
 tgcctttgt atcatatc caatctcc acttctagt tccctgatt tcttcaaaa 4800
 gattctgg aagaatgtac gaaaaatgt gaaacctat aatccatggg ataggcttct 4860
 agcagctgt atatcgctt acctgctg ttactgttg atttctct tacaacta 4920
 tgaagaaaa tatggaggca gatcaatag aagaatccc tttttcagaa accttcaac 4980
 gaagtctaaa aataggaaag tccagaacc accagacaat gggatgaag atgaagatgt 5040
 caaagctgaa agactaaag tcaagagct gatgggttc cagttgttg aggaagacc 5100
 alcaattatg gtacagaatt tgcataaga atatgatgc aaganaaat tcttcttct 5160
 aagaaaaga aagaatggg caactaata catctcttc ttgtgaaa aaggagatgt 5220
 cttagacta ttgggtccaa atgtgtgtg caaaagcaca atttataa ttctgttgg 5280
 tgatatgaa ccaactcag gccagttat tttagagat tattttcag agacaagtga 5340
 agatgatct tcaatgaat gtaggttga cgtctcag ataaacctt ttgtgcaga 5400
 tactaatg gtaaacatt ttgaattta tggagctgtc aaagaatga tgcacagga 5460
 catgaagaa cctaatgct gaatacaaa tgcactgtc ttaaaaag accttcaaga 5520
 gactttaaag aactactctc caggatacaa acgaagtgc ttgttctc taagtatgt 5580
 aggaatct catatctt tctatagta accatctaca gttatgctc ccaagacaa 5640
 acagcatg ttggcagaa ttgcactgc atttaaac agaaaagggg tctctattct 5700
 gacctctac tatagtggg aggcagggc ttctgtgt cagtagcta tcatgtgtc 5760
 ttggcagtt agatgtatg gaacagtaca acatctaaag agtaatttg gaaaggtga 5820
 ctttttggaa atcaattga aggttgat agaaaacta gaagttagcc gcttcaag 5880
 agaaattcag tatatttcc caatgcaag cgtcagaa agttttctt ctatttggc 5940
 ttataaatt cctcaggaag atgtctcgc cttttcaga tctttttta agcttggaa 6000
 agctaaat gcttttgcra tgaagata tagcttttct caagcaact tggacaggt 6060
 ttgttagaa cttactaag aacaagaga gaagataat agtttggaa ctttaacag 6120
 cacacttgg ttggnaacga cacagaaga tagatagta ttttgaatt ttattgttgc 6180
 gctgtgtac tggactct tcttttca cttatttta actttgttt aaaaagttt 6240
 ttatttgaat ggtcaactgga gaaccaagaa ctttttcaa gctcttaatt 6300

lgaatgctg tgggtgctg tttgtcttc cttaataaa aacgcatgta taattaaaa 6360
 aaaaaaa
 <210> 34
 <211> 2558
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> misc_feature
 <223> Incyte ID No: 3105257CBI
 <220>
 <221> unanure
 <222> 2523
 <223> a, t, c, g, or other
 <400> 34
 atggggcgcg gggggcgcgc tgcctctggg cgttggagcc gggggcgcg ggaaggatcg 60
 ctggcggggc gggggctctg ggggtctggg gggccacgcg gggctccggg gggctccggg 120
 gctgggggct tggggcgcgc agctggcggg gggggggcct gggggcggcg gggggcggcg 180
 ccttcgcggg agggaccacgc gctaacgacc atggggcgcg gggctctgga gggcgggaag 240
 ccccgggggc tgggttaacc tggggcgcag agggcggcgc cggggctggt atctctacag 300
 atggctgggt ttatctccat gctggccct tctctctg ggaagatcat cgtatcgatc 360
 taacaaccaa ccaactgtga ctacggcgac aacctggacc gggcttgcct agggctagct 420
 gggggcttc tctgtggctg tggcgccaat tgcctctggt tctaccatc gcaacttcaa 480
 ggtccgggca ttgtatagtg gctggagact tcatatct cctccatct gaggcggag 540
 gtgtctctc ttgaacagac tggcagagga gaattgat accggctctc atcagacact 600
 gaaccctcgg ggggtcaggt gactggaaac ccttcagatg ggtccagggc cggggcgcag 660
 gcttcgggag gaaatcagat gatgtttt gcttaccata atcgggcgc ctgtgttgg 720
 aggtgggggc ccccggtctc aatcatgct gtaatttatg ggggatact agggaaactg 780
 accaaagaaa cctggcgtac cctggcaca ggcactcgc tggcttggga acgatctgga 840
 aatgggggaa cgtctgcgcg ttctgggaaa gaatggagct aatcggagaa atatggcag 900
 aagtgagcag ggtctccgag aacctggctc gggcttctg tctgtgtaaa agggggctg 1020
 ctggatgggaa gttggccaat gacgltgggt gactctctt ccttctcaat gtagcttcc 1080
 tgggttggaa taagatctg aggtctggc tcttctact cggggcgcg gaaggacgg 1140
 ggttggaggg ggggcctctg gggctcccg gggagagagc ccaagctggc tttaacagag 1200
 ggggttccat taatggagaa aggtctcag ggtgttgg agtttaagaa cgttcatctt 1260
 ggttccctcg ctggcgcgga ggtgcacata ttccagagat tcaaggcttc catctcgta 1320
 ggaatctgca cggcactggt tggcgcaagt ggttctggca aatcaaat gcttccacc 1380
 ctggcgaggt ttatggagcc tggcttggga actatagtc ttgtatggca ttgacatcgg 1440
 cggctaaacc cagtgtggct gaaatccaaa atctggagcg ttggctgaga accaatcttg 1500
 tcttctgct ccatgtctga gaaatctgct tatgtgtctg atgacccctc ctctgtgac 1560
 gctgggggaa tccagagagt ggttgaagtg gggtaacag ttggctcagc cgggaatttc 1620
 ccccaagggt tcaacacgtt ggttgggaaa aagggttgtt- tctctcaag ttggcgcaaaa 1680
 caggcgatg cgttggcgcg tggcttggga aagatccca aattctctc cctgatgaa 1740
 gaaacagtg cgttggatgc cgaataatg tatcctgttc aagaagctc agatcgatct 1800
 atgggtggaa gaacgtgctt agttatgc catgtctgt ccaactttaa gaatgtaact 1860
 atgggtctg ttcttgacca aggaataat actgaaatg gaaacatgaa atggatctat 1920
 tcaaaaccaa atgggataa cagaataat atgaaacaa aagatttlat ttccggctaaa 1980
 ggaagcaat actgtttaa aatattgac tttaatgaa aacgtgttg cagaataaaa 2040
 atccagagac tatgataac ataaacata tatcagatc ttgaaataa accaatctt 2100
 ctccaaagtg tttaaaagat tgtttggaaa cgttaagct ctccagact tttaattcg 2160
 agtttcaata attgaaact tttaatgtc tttagacatg aagttatct caggtttct 2220
 attctctt ctgtggagat atttcaat atatagatg atgtcaatc ttcttggc 2280
 tgcgtttaa gtttgagct atgtcaat gaaactta aagaagcaat tataataaa 2340
 aagcctgctt atcttaagc agtttaccga taactgtgta atcttcttgg tagatctaa 2400
 cctaaactta atcgatctt taccgatcga ataaagcgt atctgaattt ataatcttat 2460
 tgggtaaact ctggaggaac cctcttctt aacacggcg aggaaggaggt tttttggcc 2520
 cggcggtt aagggggac aggaataatc cctccgaa 2558
 <210> 35

<211> 5065
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> misc_feature
 <223> Incyte ID No: 3200979CBI

<400> 35
 atgttataaa aagaataag cgttgcata caaatcag ctctctgta caagaattt 60
 cttaaaaaat ggaataaaa aagaagttt atgggcatc atttggcatc ctcttgaaa 120
 cacttcagag ctacccgtt tcttgacaa cctcccaag tcttggagag cgtggatag 180
 tttaatgact ctggcctggt agttgcat acacggatca gtaactaac acaagaata 240
 atgaataaga tggccttgg ttcccttaag aagaagaaa cactcttgg gacacagat 300
 gaagagaca tgaataaac acttcaaaa aatatcaaaa aatgggttgg agttatatt 360
 aggttaact tctatatac cttgaagtt aattggggat atgaatccc agttatag 420
 gggacactcg aatacaaaa acactgttg gcaatggag gtaaatct ttgtacttg 480
 gaaagtact ggttaaaag gtttgagct ttccagcgc caattatgc ttgaatata 540
 gaagtacaaa caaatcatlc tgaatggag gagtggac cagtatctg aatlaatat 600
 aagatcacac ctctcatlc taaaggagaa atlatgaag aatgttatc tttaacttc 660
 ttagtctct tctctctt tatatact ccatcttca atgttcag ggaagagaa 720
 aatttataa aactgatgac agttgaggt ctccggagat cagcatctg ggtcccggt 780
 gaattgcact acatttgcct catcttcat atgtccatc ttatgtctc taqtaataa 840
 tcaactccaa ttgtatttca taactgttc atgtgatat tcaactata taqtaataa 900
 ggtcttctt tgaatagact ggtcttccc atgaagttt taatlaagaa accatggc 960
 ggtgttttg ctggatct ctacactgta ttgtgggat ttctggggt cagtgtgta 1020
 tatgaacaa ttctttatc ttggagtg gttatagc tctctagcc ttgtgcttc 1080
 actgttggaa tggcagatc taacacccg gataatctac taagtgtgtc tatcttcc 1140
 gatccctctt ggtatttca caaaatgaa ggcacttctc taagtgttc atttgatat 1200
 ctcttcatc ttatattac attatatt gaggagttc taactgtata agtatagat 1260
 ggggactctc catatttct cttaaagtc tcaatgctc tcaattggt caaacatca aattctcat 1320
 ctggaaact ttggagtg aataatct gggactctc ctgatatc ttltgaacg 1380
 gtgtcccgag aatctcag aaaaagagc ataaagata gaattgtat aaagaatat 1440
 aattggagaa ctggaaaagt agaaagctc caagagatc ttlttgatc atgtgaaga 1500
 cggatcacac caatcttgg gcaatagga gctgggaac caacactctc aactcttct 1560
 agtgatctgt cgttttcc agaaagatca ggcactatct ataatccca acccttgaa 1620
 ataatgtac ttgaagaa tagaagaa atgtgattc gttccagct caatttcaa 1680
 ttggacttc tcaatggag agaaactc aggtatttg ctataataa aggtatctag 1740
 ccaaaagaa ttgaacaaag gtttttgg ctgaatgaac caactctgg atgtatccc 1800
 ttlttaagag accgaagtg gggcctcct aaggaaagc aatlaagaa acttaactc 1860
 ttcaatgacc aatcatgaa ttgaagttac ggcactatct ataatccca acccttgaa 1920
 aatggagat ttgaattgc aggtatctc ttgttctga agcgaagtg ggttatgaa 1980
 taactctaa gtttaccag gaattgaag ttggagagtg ttggagagc atctctat 2040
 aagcagaca ttctgtatg caagttaaca acgaagag ttgaataatc atccctatc 2100
 ttggctttg aaaaaagaa caatttcca gatctttaa gttacttga taagtgtct 2160
 gacggggca taaggaaatc tggctttaa gttacatlc ttgaatagt atcttgac 2220
 ctgaagagaa aatcagaat ttatgaaca gatlttgaat ttgggaaca agagaataa 2280
 cagtggaca gaataatct agttgagct gaattggagc aggtatctt ttctctctc 2340
 gaaataagaa aggtatctg tatggagct ctctggagc acaatctta ttgattgca 2400
 aactctgct accgtatct aaggtctga aggaagctc ttltgttct gttacatga 2460
 ctggaaatg ctlttaacc catcatctta ggaagatata ttgaatagt aactctgaa 2520
 actcattgt ggggttcc acccgatg tatctctc ctctggaaca aatcccgag 2580
 aggcctctta agcctggtt aatgttlaat aatcagagt caaatatga agcctctg 2640
 catctatga agtctcagc tatgtttg gaattagat aatlaagaa caaaatgct 2700
 tcaagatc cctccacaa ttggggcact atagtgtc gttacagaa ggttatagaa 2760
 tttaagct catgtaac caagaatg aatgttctc ctgtctat ctgtctat 2820
 agcaatcct ttatltgaat tttaactc acagatctta ttcaattgga gttacatct 2880
 ttlttttgg atgcaatg tcttcaat ggttatagc atgttctctg atttgttg 2940
 ttgaatgaa actgatctc tcttcaat ttgttatag gttatagga gttatagat 3000
 ccttccctca tctcttcat ctgttgag aaaaagtc aatccaatc atgtatctta 3060
 gggcctcggc ctctagaaa ctgtgtgga caggtctcg ttgaactct ataacctc 3120
 ttgatctcc ttcaataaa cttaactac tttaactac tttaagatc caggtctca 3180
 ttggaaacta tgtttgtt ttgtgtatg ataatgtt ttggagctc tcttatatc 3240

ctcaagtacg tgccttctat catcttcttg aagtgagagaa aaaaataatgg cttttgggtct 3300
 ttgtgctttt tcatgtcttt aatatgtgta tccacaattc tggatcaaac caagtatgaa 3360
 aaacccaggt taatttttgt tatgatatac taacctctct ttactttcct caatagtatcg 3420
 ttaatgtacc agctcaactt tatgatgatg agaaacttgg acagcttgga caatagaata 3480
 aatgnagcca ataaacacat ttttttaaca aacttaatac cataccttca gagtgattat 3540
 ttcctttttg tcaatagttg tctgaaatg agtatggaa atgaaaacat gaataaagac 3600
 cagcttttca gaattctccc agcaagtaga gaactcata ccaatccaga agagcttgaa 3660
 gaagatgttc aagctgaag agtccaagca gcaaatgcac tcaactatcc aaacttgagg 3720
 gaggaacacg tcaatactgc agctgttta caaagaatc attatgagac aaagaagaat 3780
 tgccttttca caacaaagaa gaagcagcc atcagaatg ttttgttttg tgttaaaaaa 3840
 ggtgaagttt tgggtattact aggcacaaat agagtctgga aagatcttc cattaaaatg 3900
 ataactgggt gaacagtgc aactcagga ttggtgtgtg tacaaggcaa cagagcatca 3960
 tgaaggcaac agcgtgaca cagctcaag ttcttgggtg actgccctca gagaactca 4020
 ctatggccca agcttcaaat gaagcagac ttggagtgtg atgcagctgt gaagagctg 4080
 ggcagaagag atgctgctct cagttattca ctgtgtgttg aagctcttaa gctccaggaa 4140
 caactaagg cctcctgtga aactatata caggggaata agagaaagct gtgctttgtg 4200
 ctgagatccc tggggaaacc atcagtggtg ctcttagatg agcgtttcac cgggtaggac 4260
 cccgaggggc agcagcaaat gtggcagata ctctaggcta ccatataaaa ccaggagagg 4320
 ggcacctctg tgcacaccca taaatgtca gaggctaaat ccttgtgtga ccgtgtggcc 4380
 atcaaggatt attacttga aataaaatg agcagagct gcttggcagg aaagatatc ctctttaa 4440
 ggttaagatt tgaagctttt cccacaggtg gcttggcagg aaagatatc ctctttaa 4500
 acagagatt ttactatca tctcagagct gcttggcagg aaagatatc ctctttaa 4560
 ggtataagt taccctgtga gtagctcac cctctatctc ggcctttt caagttagag 4620
 gcatgaac agactttca catgagaa taaagctctc taaagctctc cttagagacg 4680
 gtaattctga acactgtga agcagagag ctgggaatg ttgatgata atgtgataca 4740
 acagttgaat gaaactctct cccagagga gaccttaaa atgaagaact tccataatt 4800
 caatttttag tctactata tcttagtttt ccaataattc aaagaagct tctctttac 4860
 tbaagttaac aaaaagaaat attcaaatg ttaaacatgc aaatagatt acagtttca 4920
 tttttaaaa tttagatgta agcaaaaagg aaatatagg aaagtagta gcaaaaatta 4980
 acaaaatagg acatgtatt atcccacaa tgggtctatt ttgctgttag gggatccat 5040
 agttagaac gcggagccgc gtgtg

<210> 36
 <211> 1677
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 6754139CBI

<400> 36
 gtggaacag gtttgagaga tactgagagg ggcagagcag tgggttttag aatccctggg 60
 tgaagcttg gactctgtg gctatttgg gccctcttag tcttggtga gaggcaggca 120
 gactccaggt ccttganaag gggaggttgg aggaanaatt tgcagctcg ggcacagaag 180
 atagtaccag ttaactcat ggcctttacc tatgtgtcc ctgcagcag gccagggagg 240
 aactagacc acagctagag caagagaagg cagacacacg gagacactc ataggagacg 300
 ggcacagcc ctggaggtgg aggtgttag cagagccct gggactaggg cctggatagg 360
 acaaactccc ttaacttgc tcaagatgc ctggagctg agggccggct ggccttcagg 420
 ctgttctgtg aacttttgc caactacaca agtgccttga gactcttggc agacacacg 480
 cagactctga atgtacctt gtaggtgaca ctgtcccca tcaatgacat gattgaacgg 540
 aacaggttgc tgaacctga tctgttgata cggagagagt gpacacatgc ctactacga 600
 tgggaaccca atgcetata tggcttgat gcatccgca tcccacagag tcttgttgg 660
 ggcgcagaca tgaacttca taaacaaagg cagcgcagc cctcagatc ccagacacc 720
 aacgttggtc tggccagca tggcgcctgt cgtgtggag cgcctggcat cagcgcagc 780
 tctgtccgag tgaatgtag agccttccc ttgcagccc agcactcggc ctgagcttc 840
 ggcctctga ctacggggc gaccaaagt cagtgtgcgc cgcgcgggc tgcagcagc 900
 ctggcggact tctgtagaa ctgtagatgg gctgtgtcgg cagtcgcggc gggcgggcg 960
 gttctaaact aggtgtgtgt tctcagacc taccggagc tcaactcac gctgtctgt 1020
 gcgcgcggc cagcgcgcta cgtgtgaac ctctgtctg cctgtgtcgc catctcgtc 1080
 ctggcgcgc tgccttca ctgctccc gactcagcg aagagtgct cgtggcgctc 1140
 acagtgctgc tggcgtcac ggtcttccag atgttctgag tgaagacat gccacggcc 1200
 gagagcgtgc cgtctatcgg gaagtactac atggccacta tgaactcgt cacattctca 1260

acagacacta caatccttat catgaacctg cattaactgt gtcccagtgt ccgcccagtg 1320
 ccagctctgg ctagggccct cctgtctgga cactctggac gggcctgtg cgtcgggaa 1380
 agagggagg cctgtgggca gtccaggcca cctgagttat cctctagccc ccagctcct 1440
 ctgagggagg ctggccccc agcggccct tgcacagc cactgtct ctgctggcag 1500
 gaagccctac tgcacacagt agccacctt gcaatacct tccgacgcca ccagagtcgc 1560
 cagctctg atgagagctg gaagcgcctg cccctgtgga tggaccgctt cctctggcc 1620
 atctcttct catggccct ggtcatgagc gtccctgtgc tggcagagg cctgtga 1677

<210> 37
 <211> 3714
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 6996659CBI

<400> 37
 atgagagac tgaatttttg gtggctgtg agcagggtct gtctctgtt gccgcgcgc 60
 tggcacttg tctggcggg ggtgccagc tctctcgc accgcagcc ctgcagatc 120
 ctaagagca tgggcagcg gctgaggatg ggcgggtgc acttcagcc ctgaccacc 180
 gcccccgc cggcagcgc cgtccggag cactccgag caggagcca gaggatgag 240
 ccgagacag ggaactagcg gtcccgcg cctccgcgg gcgcacgtg gtggggagc 300
 acctgtcg gccggggccc gccggctcc cgtaaagcc ggaaggcgc cagggcagag 360
 gctcttgc cagggagcg cctctattt gcgcggcca acctgaacc cgtggaaagg 420
 gtctacctt acaactctc ttbgaaata gtgagggca tggagcagg cctggggcag 480
 ctgcacttt tgcctcttc tctccatgt tgcctatga cagtgagcc tttctcttc 540
 ctgcgaagtg tggccatcac cgtgtgtgtg caagggtgt cggcgtgct cgcctcccc 600
 cagagccag tggcaaatgat ggaactcag ttggctcagt tagctctga cttccagtg 660
 atcagatcg tggcagagc gtttcgcgg gtagctaga accctctca cctacaactg 720
 agttgaaa attcattag ttctgatgt gtgtgtgtg caggagact ggaacatcac 780
 aactgtaca attttagt gtgtgtgtg caggagact ggaacatcac cgtattctc 840
 cctcttccc agaatattc caagtccac ctgtgtcta tcatcaat cagcgtcaac 900
 ctccctccc cccagacct cttagcttc ctacagatcc agttgagag tattaaaga 960
 agcacacca cagtgtgat gtttggcgc gactcggaa gttccggcg gatttcgaa 1020
 attacaccc agtttgggt cagccccct gaactctgt ggtgtctggg agattcccc 1080
 aattggagg aactgaggag agaggtctg cctttaggac tcatgtctca tggaaaaa 1140
 acacagcttg tctttgagca ctactacaa gatgtatgg agctgtctgc aagagctga 1200
 gccacagca ccatgatcca accgaactt gctctcttc ccagcagct gaactgact 1260
 gaggtgaaa ctacaaatt cacttcagg tatatttat caagtttct agccaatcac 1320
 actttcag gctcagtg ttccatcaga gtaaaagt cccactgt cagctcagaa 1380
 acaacttt tcatctggaa tctcaaat gacccatgg gaaagccaat gtggaccgcg 1440
 ttgggcagt gccagggggg aaagatttc atggactat gaatatgcc agacagccc 1500
 cagagacaca aaacacctt ccaactcca agtaagctac acttgaggt ggttacctg 1560
 attgagcac cttttgtt ccaagggag tagatgatg aaggttgtg cctgtctggc 1620
 caactctgc tagaccctc gactaatgac tctcccat tggcagct ttttagcagc 1680
 ctccataga gtaatgatac agtgccatt aaattcaaga agtctgcta tggatatgc 1740
 attgatctgc tggaaaagt agcagaac atgaaacttg acttcagct ctatatgta 1800
 ggggaaggaa agtatgggc atggaaaat gggcacttgg ctggctagt ggttgatctc 1860
 ctggagagca ctgcacat ggcactcat tcttttaga tcaatactg accgagccag 1920
 aactctgtt atgcctctt gtttgcaga acagtggcca tcaaacctc aaattgttg 1980
 gtagacag ctccactgg agccttcag tggccactc actggacaat gtggctggg 2040
 attttgttg cttgcaat cactgcctc tctctactc tgtatgaat gaaggtcca 2100
 ttgttttga ctcccaagg ggaataaga agtaagctt tctcttctc ttcagctct 2160
 aactctgt atgcctctt gtttgcaga acagtggcca tcaaacctc aaattgttg 2220
 actggaagt tctatgaa ccttgggc attttgtga tgtttgtc tccacata 2280
 accgaaact ttgtctgt gatgtagt gagaagatc attgaagct tctggaata 2340
 catgaccca agttactca tcttccaa ggaactcgt tggaaactgt ccggaagac 2400
 agtgcctag atlatgtag acaagtttc cagagatgc atgaatat ggaaggtac 2460
 aatgttccag ccaacctga tggagtggag tatcagaaga atgatccga gaactagac 2520
 gcttcatca tggagaagc cctctggat tatgaagtg caatagatc tgaactgac 2580
 ccttcatg tggggagac atttgccata gaagttacg gaattggcc cccacctac 2640
 tctctactg tggggagac atttgccata atcagttacg acaagtcga tgggtttatg 2700

gatalgtctc atgacagatg gtacaggggtg gtccctgttg gcaaggagag ttctgtctgc 2760
 acgagagctc tgcacaggg catcaaacac ttctctgggc tcttttgtgt gctgtgtcat 2820
 ggaattgtgc tgcctatctt gaccacatc ggttagacaca tagatagag gctgtgtcat 2880
 ccaagatctc aaacacatc cagatctgac tacgtctcc acaccagac ggaattacac 2940
 agagagatca ataacatgc talagagagaa aagacagagc attcaagcc caaacgttgg 3000
 gaaaagaggtc cttaattgtgt acccgtctag ctacagctat ggaatactgc caatctgagt 3060
 catgacacac gacgagaaata cttctttagt gattagagag gacaaacaa gctcagacac 3120
 cggatcacac aggaacatccc cctcccctca agggagagag agtccctcgc ctgtggagac 3180
 accaatttgc aggcagatct cctaaatgtta tctggagact cagttagatc ggaattctca 3240
 agacgtcaga agcagatctca ggttagatcgt cagagctcgc agctgtgtct gaaagagaaa 3300
 accggagcttg agcagatctca aagggacaggt cggagcttggt agtctcaggt gaccacatcg 3360
 ctccctcttc tcaagtctcg acctctctc ggtgccttga gacctttgt aatgtctctt 3420
 gtaacatctc gacaaaggtg tggggagagc ggggtccagg tctctttaa ggtcagagct 3480
 tctcccctcc gcttaaaagt cagcagagc tctctcagc tctactctc agtctctct 3540
 ggttagaggtg ttctctcagc aggaatctta gtcaagata agctctgtgc gaaagatctg 3600
 tgaataacaa gtttccctcc agtgccttta acctcttacc caggtctcgc agtaaatctt 3660
 ctgtgtttta gacctctc tgcattacca acaaagata aattgttacc tcaac 3714

<210> 38
 <211> 1009
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc.feature
 <223> Incyte ID No: 7472747CB1

<400> 38
 cgaaggagag gggacgggag gggcagagct gggcacagcg cccatgagtc ttaactctca 60
 cctgtctccc accctctccc catagcagc ccccaaggcc tggcaggttc ctgtccacag 120
 catgtccacgt gcttgagctct ggaatctcgt ggggtggcccg gttgttcctg tctgtctgac 180
 ctaagtctgc taccgtctgc tgggttgacac tatctctcag ctgtctaaaga gggcagcgaga 240
 ggtctagacc agggacacgtg tctaatgttga gaaagtctgc ttcctgtaga actacacacg 300
 ccggagacac tgggacagagc agcaatttgc gaaagtctac atcgatggagc ggggtgaag 360
 tctgaaaccc aagggacact ctacacaccc cagacatctgg gacttttcca gaaattctct 420
 ctttgacagc aagatctgca ctacacatg ttatggagac ctggcaccca gmacaaggcg 480
 aagttcacag tctgtgtctc tctatgacct gtttgagacc cccgtttaaag tgaatctctc 540
 caacacatct ggcacagagcg tgggtggccca tctgtgcgcg attgaaagat gggagagacg 600
 tcccaaggcg tcccaaggagc tatctcaagt cctgtgacct gctctgttcc tgaaccttgg 660
 gacgttgatc attctacatc tccacacacat ggtcttcagc cagtgtgagag ctgtgtacgt 720
 ccggcgtgggc tctctcttct cttctatcac atttgacac atgtgcttg gggcctatgt 780
 tgcagggacaa gaccacagca agaatatbat ctcaagtatc cggagcttgg cagcacatctg 840
 gatctctctg ggcctgtgct ggtctgtgct gatctctcca ctgggcccgc tggctctgca 900
 cgaatctctg cagctctgct tgcctagtag gggcctctgcg gtcaaggatg gggcagcctc 960
 tgaacctcgt gggctcccca gggctcagaa gattcccatc tctgcatga 1009

<210> 39
 <211> 1155
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc.feature
 <223> Incyte ID No: 7474121CB1

<400> 39
 atggagagct cggggacacc ccaaggcaggg agatgtccc cagagagcct gggaagatc 60
 tcccacagcc tgggtctctc ctgtcttctg gtacacatag cccgtgtggg tggctgtgct 120
 ttctctgcca ttggagagag ccaagtctcg gggcagcagc aatgtggagg ttcttgaag 180
 ttctctggag agctctcagc aatcttgacc tgcacatgaaa cagtgtgtgga agaaagaaa 240
 cagatctccc agggacatct gaaagaggtg aagctcaggt ggtttacac gacacacac 300
 tggctctccc tgaagctcgt ctcttctgac tgaacgtgtg tcaagacgt gggctatggc 360
 taactatcac ccgtacacag gctttgcagc taattgtgca tgcctatgg tctcttggc 420

atcccctga tgttctctct tctcacagac aagagcgaaa tctctgcaac catcttatct 480
 aactatata atcggtctccg aaattccctc ttcttaccg gcccctctct ctccacatg 540
 tggcccaatc ctctcttcca gaaaacacg gaccacacg cccgcagatga agcttctct 600
 cagttctatc tcaattgttga agagcttcca gggcccaaac ttggacatga tctctcagc 660
 caactctgca atagctcagc gtttagagaa tctctagatc ttgaataaa gaaacatga 720
 ccaatggccc cacaagacat ggaagagatc aactctgtc ccaacttgtt gttggagaaa 780
 ctctacatcc ccaatctcag caactcgtgt gaagttagac agaaagtga gaggtttagc 840
 atcccctcc ccaatctcag ccttatgtct ttgtcctaaa tctctctggc agtctcacc 900
 ctcccctctc gggagacaaa gtttgatctc gagaatggct tcaatttgc ctltgtacca 960
 ctccacacaa ttgggttttg ggaatctgtc ttagaacac ctactctctc cgttctctc 1020
 tccatttcaa tcaatgttgg aatggagatc gttgtatg ctctcaagt gttggcaaac 1080
 aggtttagtg aatataaaa aatgttatg ctatctcttg caaaggaaa gttttacac 1140
 ctgttataa agtga 1155

<210> 40
 <211> 2733
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc.feature
 <223> Incyte ID No: 7475615CB1

<400> 40
 cccctcatct agtctcttca ccaagcacag gaaagcacg ctgatagag cgaagatgag 60
 cgaagcagat ttgglttgag ggaagttttt gaaatgtca atgaagttca agacagatga 120
 cctgtggcct gtttagagag ggaatgtcac tccgaagatg tacttagaca cggaaatcac 180
 gaacacatc cgtctgtccg ctcccacag acatctgcag ccttgccagc cggatcttg 240
 taccacatc ctlttaaaag ccccagatat gaggcacagc aggcctccct acgttgttga 300
 cgaagacgca taccatctca cccctcttca gaaatgtctc ttgaagaaaga ccaagacata 360
 ccaatgttga gaaacatctc gcaatctctc caatgttct tccggcagaa tccaaacatg 420
 ggttttggg ctgtctgcgg tgcctctcgt gctcccacag taagatata aagacatat 480
 catctccagc cctgtcgggt gactcagcg ggaatccalc caggttccac aaggtctgac 540
 attgtctctc ttggccacag tctctcagc caatggctc tactctctc tctcccctc 600
 cctggacata tctctcttgg ggggtgttga ccaagatggc ccaagatgct ctccgttat 660
 cagatctctg gttgttataa tctgttctga gcttggccca gattcgaat tcaagttct 720
 caacacatgc acaaataga gctatgttga caacagagc atggagcttg agaggttga 780
 cgtgtcagct acgtttagct gactcacgc catcacacgc atggatctga tgccttga 840
 gtttgttct gttggcatct accctctcga gtaacatctc ggggtcttca tgcctgcgc 900
 cggcctcaga atccatgatt cgtgttctca gtaacatctc gaaatgaa tcccctccta 960
 caagacgca ggtccatctg ctctacatc catgtacat tgaataaac tcccacacac 1020
 caaatcgcc tgcctatct tgcctctat cagcgttgc ttccgttgc tggtagagga 1080
 gttcaatgct cgtacatctc acaagatctg ctctccac cctcaagaa tgaattgtgt 1140
 gttgttga acaatctct cgggtggctg taagatggcc aaaaagatc acatgtagat 1200
 cgttggagaa atccaaagcg gtttcccaac cccgtgtcgg cctgtgtctc cagagtgaat 1260
 ggaatgata gggacagctc tctctcagc calogtagc taagttatca acctgtctat 1320
 gggccgagac ctggcacaa agcaggtga cgaagtggat tgaacacag agatgatcgc 1380
 tctgtctcgc agcaactct ttgtctctc ctlttaaatc caatgtctg tctgtgtcgc 1440
 tctgtctac ctgtctgttg atggatctg agaaatctc caatgtctg taaggagcct 1500
 gatctgtc aatctcagc acctctcaa gcaactcac gactctcgt acctgtggag 1560
 gaagagcag ctggaatct gacttgggt ctctccagc ctctctctc tctctctag 1620
 ccggcccat ggtgttgagc tgggtgtcgc ctctccgc ctgtctgtg tcttccagc 1680
 tcaatttga aatgtctat cactgtgac gttcaatgag atgtacatc attgtagatc 1740
 ccaagctcat aatagggcc agaatacca gggatataa atcatagct atgtgacccc 1800
 tctctactt gccaactcag agatctttag gcaaaagtc atggcagaa ctgtctcct 1860
 gaaagatc cagcagatc ttgaatgtc gcccctcac gaccacaaa acacacagac 1920
 cccgtctaac gttccagcgc tgtctatct caactctac ccttgaagc cctcacctc 1980
 cagagtttag ccaacacgc cccgttgagcg ccccgagcg ccaagtgaaa tggatctcgc 2040
 cgtcccacac ttcttctac tccaaacct calctctgag ataaagttag tcaagttctg 2100
 gactttag gttcatcag cccgtgcaca gctgaatcc acctaatgaa agatcggct 2160
 gaagttctc ttgtttagac tcaatgcca gtttgttacc gaacttagc atggagatc 2220
 ctlttagag ggtgttagc aatgcaagc gtttctctc agatctagc atcgatctct 2280
 ctlttccag gaaatgta gaaagtga cccagagcac aacttcaag gggctcagc 2340

ggatgctgag ctctctctgt acgactcaga ggaagacatt cgcagctact gggacttaga 2400
gcaggagatg ttccggagca tgtttcacgc agagaccctg acgcctctgt ggggctcag 2460
ccagttctca tgttgctctac agagtgcctg gcacttgaga ctctcataaa gtagagacct 2520
gggggtcagc ggggtgtctg gggagggaaa gtcatccccc cagagctctt gttctctct 2580
ctctctcccc tctctctccc ctctctctcc tcccgcgac tccagagaga gctctcagc 2640
ggcctggggg tgtacacctt ccagagatga gagtctggtg agcccaacct tcaaccgtca 2700
ggcctggggg caatggacaa gacctctgtc cac 2733

<210> 41

<211> 3457

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7475656CB1

<400> 41

cgagctcaga gcccgcgcg tgcgcggcgc cgtctccgg gcatggaagg aggggcaag 60
cccaactctt cgtctaacag cggggcagat ggcacacgcg tcttcccgc caaggctcc 120
ggccggggcg cggggccggc cgggcgcag aegcctctgg gaccccccgc ggggggcggc 180
ggggccggcg cgaaggagca cggcaactcc gtgtcttcca aggtggacgg cgttggcggc 240
gaggagcggt cggggggctt cgaagacgcc gaggggcccc ggcgcagta cggcttctat 300
cagagcgagt taactctcat gctgagccc ggggtcaaca aattctccct cgcgatggtt 360
ggagccaga aggggttggg aaaggcgag gaaggggta aaactgcag cttctggatt 420
atcccacctt acagtgtatt caggttttac agattttta taagtcttat aatgatggtt 480
ggaaacttag tcatataccc agttggaatc acattcttca cagagcaaac aacacacaa 540
tggatlattt tcaatgtgac acagatata gtttctcat tggacctgat catgatatt 600
agmacctgga cgtcaatgca agacagctc gaactatcc tggaccacca agtgaatca 660
atgaatatt taanaatgc gtttgggtt gacttcatct catctatccc agtggattat 720
atcttcttta tgtttgaaaa aggaatggat tctgaagttt acaagacgc caggcgctt 780
cgcatctgta gttttacaa aattctcagt ctcttgcgtt tattacagc ttcaaggta 840
atlagatac tacataatg ggaagagata ttccacatga catatgact cgcagtgca 900
gtggtgagaa ttlttaactt catcgccatg atctgtccc ttgtccactg ggaatggtt 960
ctcagttct ttatgacctt acctcgagac tcccaccag attgttgggt gcttttaaat 1020
gaaatggtta atgatcttct gggaaagcag tattatcag cactttcca agctatgatt 1080
cacatgtgtt gcaatgggta ttgagcccaa gccccagtca gcatgtctga cctctggatt 1140
accatgctga gcaatgct cggggccacc tgcatagca ttgttgcgg caatgcacc 1200
gctttaatcc agtctctgga ttcttcgagg cgcagatgac agagaagata taagcaagt 1260
gaaacatac tgtcatctca taagtacaa gctgatatgc gtcagaagat acatgattac 1320
tatgaacaa galaccacag caaaactttt gatgggaaa atattctcaa tgaactcaat 1380
gatcctctga gagagagat agtcaacttc aactgtcga aactgttgc tacaatgctt 1440
ttatttgta atcggtacc taattttgt actgcactg agtgcactg tgaacaagt 1500
gtgttcaac ctggagatta tatcataga gaaggagcgc tgggtcaaaa aatgatttc 1560
attcaacag gtgttgcctg tgtcatata aatccagta aagaaatga gctgacagat 1620
ggctcttact ttggagagat ttgctgtgtt accaaggac gtctactgc cagtgttcca 1680
gtcgaacat attgtctgtt ttactactt tccgtggaca atttcaacg ggtcttgag 1740
gaataccaa tgaatgagag agcctttgag acagttgca ttgaccgact agatcgaat 1800
ggaaagaaa attcaattt tctgcaaaag ttccagaagg attgaaac ttggtgtttc 1860
aaacaacag agaacgaaat cctcaagaa attgtgaaac atgacagga gtaggttcag 1920
gaaatgcctt ccaatcaata tctcaaatg acaaacttga attccacat gtctactacg 1980
acccgactt cccgactag gacaaattt ccaacgggtt acacagcac cagctctct 2040
cacagaacc tgaactccc cagtcccag acacagacc ccacagcat accaactct 2100
tcaactctgt cctcacactc cgggtcttgc agctctctgt tccagagcc ttggcgctt 2160
cgaaactttc actatcttc cccacccgc tcccagctgt cactatgca cagcagcct 2220
cagagagcag tagagatgt cccagcctg cagactcag cagacagcc gtcccgcag 2280
ccacagacac cttggagctc caccgcgaaa atgaaatgc acaagtcag caggcgctt 2340
cacacacaa acctgaccc ggaagtccag ccaactcag cctcagacc ctcgtctccc 2400
caagaggtt ccaactgat tcccagact catctcactt tgggcgact caggacctcc 2460
atcccctaac cgttgcagg ggtcccagga acgggccttc aggcaggggg cagagacct 2520
gtcccgcag cgttcactgt ctccagactt ctccagacg gtgtgtcgg gaagcatccc ccccaacaa 2580
ggagtctctc cagcacccc tccacagca gctgtcttc caagagact tctctcagc 2640
ttaaacacag acccagacg agaaagacca cgaatttgtt caaatttatg atccctgtg 2700

attgtcaag cagaagaaa tacttctca aactggact ataactagat ctattttat 2760
tctatctct gatagatccc tctagctac tatgaagaga tattttagac agctgtggcc 2820
tcccgtaga atgcataaat atatacat atactataa atactatct aaattcccaa 2880
ggaggggcga aaagcctgt ttgacttca gtgtatata tctctcttc ttaaatct 2940
taagagattt aaatagcgt tgaagatta ttatttcta acctacttt acttaagtc 3000
ttgatagt atattctct atttatgaa gactcttgg attcaatgga acaaaaactg 3060
atttataaaa ggaactcaa atgactagt aaatagccc aatcaaaact tctttctat 3120
agctgtgtct ctgcactaa atgttaatc attaatggt gagaattaa taacaaatcc 3180
catttttag atctaaatg tatttgggt ctttcaatt caaataggt taagaatgc 3240
actactgct tggccacctt agagactag catggcaact gtttgttaag atataccta 3300
actcaaaa ctttcatga tctttcagaa agctgggga aaattaatat tgtttctat 3360
gtgtatcgg actttacca agactgac atgttgtt graaataact ttttcaacc 3420
aaataaaat agctattctg tgttgaaaa aaaaaa 3457

<210> 42

<211> 5622

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7480612CB1

<400> 42

ctggacaagg agaaaaact agggaaaaaa ccaacagaat ttgttggcat gttctacaca 60
cagacatggt ctttcagaa gccacgtga taanaacag ttttaaaaga ggcacacctt 120
tgtagaggag tcttggaag attctcatt gttttctgg acaaaaagag accagtggat 180
ccaagtgtt caaatacttc tcttattt tttaactct attgtctgc aatatttact 240
ttacccgtt aatgaacag acaaatggt taanaagag ataagcgtgc gtcacaaat 300
tcaggctctt ctgcacaaga atttcttca aaatgggga taanaagag agttctgga 360
ggaatggaca ataactgt tctaggctt atatttgc atcttttgg aacacttcc 420
agctaccgt ttcttgaac aactcttaa agcttggga agcttggatc agttaaaga 480
ctctggcgtt gtagtggcat atacacagt cagtaacata acacaaagga taatgaata 540
gatggcctg gctctctta tgaagaag aacagtcat gggacaccag atgaagagac 600
catgacata aacttccaa aaaaaccca tgaattcga tgaattatct tttagtatac 660
ttttctatg cgtctaat ttatgtggg atgtgaatc ccagttaa atgaagacac 720
tgaatacaca ggtcactgtt ggccatgca tgtgaaat ttgtttact tggcaagta 780
ctggctaaa aggttgttag ctttcaagc tgcgaatcat gctgaatta tagaagtaac 840
aacaaatcat ctgttaatgg aggaattgac acagtctat ggaataata tgaagatacc 900
acctttcat ttatagggg aaattatga atattatgaa tcatttact tttagtttc 960
ttttctctt ttataact ttgcatcatt aaatgtgca agggaaagag gaaaatttaa 1020
gaaactgat acagtatgg gtcctcgaga gtacgattc tggctctctt ggggattgac 1080
atacatctgc tcatctca ttatgcat ttttatgct ctggctata catcaatccc 1140
aatgtatct cactctggtt tcatgggat attcaactc tatagcttat atggccttct 1200
tttgggtgca ttggcttcc tcatggagt tttaacaag aaacctatgc ttgctgttt 1260
ggctggattt ccttcaactg tattttggg atgtctggga ttcactgtgt tatagagca 1320
aactctctta tcttgggat gggattaaag tcttcttagc ccttttgcct taactgtgg 1380
taaggcccg atcacacc tgggaatta tttaattgtt gtaatttttc ctgacccc 1440
tgggattcca tacaataga tggccacttt ttcaatttg gaatttgata ctctttcta 1500
tttgattatc caataatgt ttgacgagt tttaactggt aaggatggcc atggggattc 1560
tccatttat ccttcaatgt ggttataag cctcatttg gtccaaact caaaatctc atcagaat 1620
ctttgaaat gaataaact ctgacatc ctctgatgt tctttgac tcttttgcac cgtgtctcc 1680
agaattccat ggaaagaaag ccaataagat cagaatggt ataaagaat aaatggaaa 1740
tgcaggaaa gtagagcatc tgcagagat atttttgac atatatgaag gacagatcac 1800
tgcaactttt gggcataag gagctggtaa atcaacactg ctaaacatc ttagtggatt 1860
gtctgttct acagagagt cagccactat tttaactact caactctgt aaataactga 1920
ttcagagaa atagaaagc atattggat ttgtccagc ttcaatttc aatttgactt 1980
ctctactgt agaaaaacc tcaaggtatt tgcataata aaaggatc agccaaaaga 2040
agtggaaca paggtatctc tgcagatga accactgct cgaattgact cctttcaag 2100
acacagagt tggagcttc tgaagagca taaagtacag cgaattatc tcttcaglac 2160
cgaattatg gatgagct acatctgc cacttggc tgaagagaa gtattctgt ctaattggaa 2220
gtgaatag cagagatcat ctgtgttc gaagcgaa tgggttatg gatcatctt 2280
aaagttacac aggaatgaaa tgtgtgacac agaaaaatc acatccccta ttaagcagca 2340

gacccgacgac gacacagatgaa caacagagag tggagagaaaa ctgtatataa gtttgcctt 2400
ggaataaaag aacaaatttc cagatcttta cctgagacct gataagatgt ctgacagagg 2460
caataagaaat taagtcggtt cagtgacatc tctgatacga gatactctga acctcaagg 2520
aaatctcgca atltatgaaac cagatcttga catltgggaa caagagaaaa taacatgtgac 2580
aagaataatc ggaatgtgagc ctgaatacga acaatggctt tgtctcttc ctgaaacaa 2640
aaagagcgagc agttagtgag cttcttgag aagcaaatc taagcagtg caaaccttg 2700
cttcttaag taagagctgt aagtgagagc tcttcttgt tcttctagc taacttgaaat 2760
tgctttatc cccatcttc tagaagaat aatgtatataa gtaacttgat aactcttgat 2820
ctggagatct taacacagta tgaattctc ttctctggaa caaatctgag aagagctct 2880
baacagcgctg taactctga ataatacag atcaaatat gaagactcag tgaactctat 2940
ggaatgtcag gttatagttc tggaaataga tgaactttag aacagaaatg gctcagatga 3000
tccctctcac aatggagacaa tcaatgtgc tgggtgacag aagatataaa gattctctgt 3060
tgctgttaac acctagaaat tgaattgtt tccgtgtct atggaaatg gagaagacct 3120
ctctttggga atttttaac taacagagc taatcaaatg gagaagacct caatttttt 3180
ttcaataaac aaaaatctc ttcaactaa gttctcttc tctaacct ctattcttg 3240
tcagaaaaat gttaacacc agtatagat ttacagcttc tggctctag caaacctgtg 3300
tggaagagct cctgggtgaa ttccatata ctcttgat ctcttttca taactttaat 3360
ttacaaatc atattctggt gattctagct ttcaatggaa ctatgtgtg ttctgtgtgt 3420
atgcataat gtttggagc ttctctat atctctcaa tatgtcagc taatctctc 3480
tcagaaatct aaaaataata atggtcttg gttcttgagc ttctttatg taagtataa 3540
taacagatct agcttcat aaaaatgtc taagtgtat ttctatata tcttatttt 3600
tgtaagtcta ttatgtgc atcaatttc ttctgttct ccaatactc agaatgtat 3660
ttctctttt gtaataaggt gttctgaat gaaatagaa atgaataaa tgaataaag 3720
ccagatctc aaaaatctc cagagagag aagaatcat ccaatctcg aagagcgaa 3780
aagagagat gaaatgttc aagctgag agtcaagaa gcaaatgac tcaatctc 3840
aaacttgag gaaagacaa tcaaatctc aagctgttta caaaagaa ataatagac 3900
aaagaaatgt tgccttcaa caagaaagaa gaaataatg atcagaatg taactgttg 3960
tgtaaaaaa gtttgaatc ttggatattc agaaacaaat ggaacatga aaatctctc 4020
catataaaat aaactgtgt gaaacaaagc aactcagaa ttgtgttgt taacaaatg 4080
cagagatca gtaagcgaa agatctgaa aactcagag ttcttgagc aacgacctaa 4140
gaaagactca ctgttgcca agcttcaat gaaagagaa ttggatgtt atgaaatgt 4200
gaaagagaa ctgttgaaa atgtctgtc cagatctca gaaatgttg aagactctaa 4260
gtccagaaa caacttaag atctctgaa aactctca gaaatgttg aagaaagat 4320
gtgcttggt ctggagctc ttggagacc atcagtggt cttaagatg agcgtttac 4380
caggaatgac cctggagctc agcgaatc gttgagata cttaagata cgtttaaaa 4440
caggaatgag gttccctctc tgaacacaa ttacatga gaaatgttg ctgtgtgta 4500
cgttatgtgc atgtatgtgt caggaagatc aagtgatc gttctcaatc aactctgaa 4560
aaacaaatc gttatagatc atctcagaa aataaaatg aaagaacaa cccaggttga 4620
agctctcac aagaagatc tgaagcttc ccaagatc gcttgagag aaagatctc 4680
ccctttatg ggtatagat taactgtga gaaatcag cctttatc gttgtcttt 4740
caagatgag ggaatgaac aagcttcaa ccttgagaa taacagctc ctgagctac 4800
cttgagagag gttatcttg aactctga aagagagag ctggaaatg ttgaataaa 4860
aatgtgaaac aactgtgat gaaatctc cccagagaa gaccttaaa atgaagaaac 4920
tccaaatc caatttag tccatgaac ttgtatgtt ccaatctc acaagatgt 4980
ttctcttaac ttacgttaac aaaaagaa atcttaaaa catcaataa tgaatcag 5040
ttcaatctc aaaaatcag gttatctc ccaacatg gttatgtg tgaatataa 5100
aaatataaa atcaataat attagtgt ttacgtga ttatcaata agcaaacat 5160
ggaagatct taataaaa aatataggt tttagagc aactcaat gatacagaa 5220
tttttaaaa ttatctaac ggaatcaat tttaatgaa atcaatata aagaatctg 5280
ataaacagat gaaatgtgac agcataaat ttatataa tgaatctgt atataaat 5340
aaatgtgat tgaatctg taatataa agtatag agtatag aaagaatag caataatc 5400
taagaagaa cctgagatc atattctc taagcaatc atataatgt attttatg 5460
atcttttca aggaatgac agtatagc taatctgt cttaacta cgtcagaa 5520
tcaatgtct tgcctgtct atgtgtatgt aggtgtgac ca 5580

<210> 43
<211> 2600
<212> DNA
<213> Homo sapiens
<220>
<221> misc_feature

<223> Incyte ID No: 6952742CB1
<400> 43
ccctctcaatc ggaacccctc acgatgagt ctgatacag caagggccag cgtcatca 60
taacggagac ggggcctggg ccggcgctc cagagagag ccaagacaa atcccgaa 120
ctgggttgag gaaaggtccg ccccaaggg ccggttgct cggggaaat tgccttgg 180
ccgtgaaac ggggtctctc gtaacatgc cactctctc gacgaagatc ggcacagaa 240
cttcaacaa cagatgagac agttccctg agctctctg gacggagagc gggccagag 300
cgttcagag caagagccca gaaacccggg gttctgtga gatgcttgaag gaaacaggt 360
ggttcagag cttctcaat gttctgtg ccggcgctc gttgaagaa cgtcccgag 420
ccaaagcgtc gttctcag taacggcgg ggaatgaat gttgaagac gtaatctg 480
ggcgtgacat cggaaatac ctggcatag cctaaatc gttgcacgg gttgcagaa 540
tctcaacat ctataagctc ttcttgcaa acctaatc atctcaat gttgcagag 600
ggcgtgac gttggagac ttacagctc ttggagcat gttggagag gttgtgaa 660
ggagagctca caaccagc tttagacct ccaagagaa cctgaagac gtagcaaa 720
gaaagaaat caaagctgc gttcagag tgaacaggg ggtcagagc taagcaaac 780
gtgtgcagc cgttcacag ctatagac gttatgac gttctcag gttcagag 840
ggcgtgctc gtttcagac taactcac agcaatct gaaatgagc gttcagag 900
cctcgtgac catctgac ttgaagac taagatct ggtgttgag atccggcg 960
aagaagag cagatgag gttcaat agctgaat gttgcagc gctgcagag 1020
ccaaagtg cgaatgtgc acaagaa ttgtctgag gttgtctga gctcgaaag 1080
agctctga cactacag cactgctga ggttgagct gccaagag ctgtcgaa 1140
tcgtgtgag cactacag ttcaactg gtagctca caagctct ggtcgagag 1200
tggttgagc ttgagatc gttcagc cctctagat cccaagac aggtatag 1260
aaggttgag ttgagatc gttcagc cctctgag tgcagatc tcaatcag 1320
tggtgagat gttgcagc agtcaagat acctgtgag ttgcacaa aggtctgt 1380
ctgtgtgct ctgaacag ctacagctc tctcaatc ctgcagac agcgagag 1440
ctgtgtgag cttgtgag aagcaatc gttcgagac acctgtgac aggtgtga 1500
ggcgaacat gttctgag atgtctgag cgtctgag gttctcaac gactcaag 1560
gaaatgtc gttctgag atgtctgag gcttgagag gttctgag aaggtgtg 1620
acctcagc gttgtgag atgaagc ctgaagc gttctgag gttctgag 1680
cactatct gttgtcag aagaagc gttctgag ttgtctgag ctctgag 1740
tcaagctc cgtgtcag aagaagc gttctgag gttctgag atcccgag 1800
cgtctcag cagatgag acagatc aggtctc cctgaagc ggtgtgag 1860
gtgtcagc ttgtgagc cttaatag ctcaaaag ctctctag cagttatc 1920
aagatgag cgtgtgag gttctgag ttgtctgag ttgtctgag gttgtgag 1980
gggtgtgag gttgtgag gttctgag ttgtctgag ttgtctgag gttgtgag 2040
ttgtctc agatgagc gttgtgag cgttgagaa ctttgagc gttctgag 2100
ccctgagc cttgagc gttctgag gttctgag ttgtctgag atctgag 2160
gaaatgtc cttgagc gttctgag gttctgag ttgtctgag atctgag 2220
gtgtgagc ttgtgagc atgaagc cctgagc gttgtgag cttctgag 2280
ccctctgag gaaagcag gttgtgag cctgagc gttgtgag gttgtgag 2340
cagatgagc cttgagc gttgtgag gttgtgag gttgtgag gttgtgag 2400
tctgtgag cagttcag gttgtgag gttgtgag gttgtgag gttgtgag 2460
gatacagc aggtcagc gttgtgag gttgtgag gttgtgag gttgtgag 2520
cccaagtc ttgtgagc gttgtgag gttgtgag gttgtgag gttgtgag 2580

<210> 44
<211> 2917
<212> DNA
<213> Homo sapiens
<220>
<221> misc_feature
<223> Incyte ID No: 7478795CB1

<400> 44
gaaagagaa gaaagagag aagaagcgt cggagagac cttaatag cagagagaa 60
actgaacac agacagag ctgtgtgag ccccttcaa gaaagacat gttgtcag 120
aagagcgt gctcagc gttgtgag atgtgtgag atgtgtgag ttcaatcag 180
cttcagag cccagacat ccaacagaa gaaatgag gttgtgag gttgtgag 240
cttcagcct catgagtg gaaatgag ttcaatag catcaatc ttcaatcag 300

tggaccgacg cctctctggag gacatcgcc acttcaacat ctttgactcg gtctgggac 360
 tctgggcagc ctgcctgtac cgcagctgac tgcctgtggg agccacattt ggtgtggcca 420
 agaacagctg gctggggccc cggcgctgcg gggcctctgt gcttggccatc acctctgtt 480
 gacctcttgt ggtcactctat gccatgggtga agctgtgtgt cttctcagag gtccggagcc 540
 ccatctggga cccctgggltt tgggcccctgt tctgttggac gtacatttca cttggcgat 600
 cctctcagct ctggtgtgtg ctgtccacag tgcggccagc caccagggc ctggagccag 660
 gggcgccccc cgaaggtctgag ggtttccctg ggaagcgccc gcaacggccc gagagggcgt 720
 ctggggccac cttctcagag ctgtctctct acaccagcc cgcagctggc tctctctgg 780
 cggctccctt gttccatc gtggagctc tgggagagac ctctccccc tactacagg 840
 gccgcgccat tgaTggccatc gtatccaga aaagcatgga tcaagtccag accgctctgt 900
 tcaatgtgtg cttgtctggcc atgtgcagct catttgcgc agtatctgg ggcgcgatt 960
 ttacccctat cttgtccccc ctgaacatc gcttggaaa ctgtctctt cgtctctgg 1020
 tgtccacaga gaaacagctc tttagtgaga accgcacagg ggaacctcact tcccgctga 1080
 ccttggacac caacatgttc agcagctgtg tctccacaga catcaatgt tcttgcgga 1140
 acacagtcna ggttcagggc gtgtgtgtct tcatgttcag cctctatgg cagctctact 1200
 tggctacatt catgtggcttc ccatatcata tgaTgtgttc caacatctac ggaagctact 1260
 acaagagggt ctccaaagg gtccagaatg ccttggccag agcgagcaac accggggagg 1320
 agaccatcag tgcctagaag actgtccgga gcttgcgaa tgaaggaggag gaggcagagg 1380
 tgtacctggc gaagctgcag caggtgtaca agctgaacag gaaggaggca gtgctctaca 1440
 tgtactactg ctggggcagc gggctcaac tgcctgtgtt ccaggctcag atoctact 1500
 accggggcca cctgtctatc ctaggccaga tgaccagcgg caacctcact gcttctaca 1560
 tctacagatt tgtctctggga gattgtatgg agtccgtggg ctccgtctac agtggcctga 1620
 tgcaggaggt gggggctgtg gagaaggtgt tgcagttcat cgaccggcag ccgacctagg 1680
 tgcacgtgg cagcttgccc ccgcacaccc tggagggccg ggttgacttt gagaatgtga 1740
 ccttcaacta cgcacactgg ccccaacccc aggtctctga gaatgtctcc ttacgctctg 1800
 cccccggca ggttgaagccc ctggtggggc cctcggggcag tgggaagagc tctgtctga 1860
 acatctctga gaactctctac cccctggagg gggggccgggt gctgtgggac gccaagcccc 1920
 tcaagcccta gaccacaag tactgtacc gttgcgaa cttgtgtgag caggagcccc 1980
 tgtgttgc cgctccact acggtatca tctctacgg cctggccact gtctcttg 2040
 aga tgggtgtt ggaagcccca cagaagcca atgcacagg ttacatcag gaactccagg 2100
 accgtctagg cacagagca tgggagagag ggcctcagct gtacatctgc cagaagcagc 2160
 ggggtggccat gggcgggct ctgtgtcggga accctcaggt cctcatctgc gatgaagcca 2220
 ccagcgctt ggaTgcagag agcagatc tgaTccagca ggcatactat ggcacactgc 2280
 aagaacacac ggtactcact atcgccacc ggtcagacc cgtggagacc ggcacactc 2340
 ttgtgtgtt ggaacagggc cgcgtgtgtc agcaggggac cccacggcag ctctggccc 2400
 agggccagct ctacgcagc atgtgtcagc ggcagatgtt gggcgcttcag ccccgccagc 2460
 acttccagc tggccacaac gagctctgtg ccaagggcag tcaacaggc tgaTggggg 2520
 cccctgttc tcccggtggg cagagagacc cgggtgcgc ctggcagatg tggccacgga 2580
 gggcccccac tgcctctcga gccagggctt gcagactgga aagcagact gcaatgtccc 2640
 atgggcaacc gcttctctga ctttgcacct ggtccctgg ccaatccag gcaactcct 2700
 accctgtctg gctctgagcca accgtctcag ggaactccct agctctctaa gcaagggtag 2760
 agctgcttt ttaaacatag gtcttaccag ggttttact gttgtgttg aggcaccca 2820
 gtaactctt agattctaaa aaccttttc taattgggag taatggcgg cactttcac 2880
 aagatgttt agaaactttt gaggccaggag tgaatgg 2917

<210> 45
 <211> 1474
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> misc_feature
 <223> Incyte ID No: 656293CB1

<400> 45
 atggggctcc ggaagccacca cctcagcctg ggcctttgc ttctgtttt actcctcga 60
 gagTgctcgg gagctgaagg cggctggct ctaagctgt tccgtacct cttgcgcaac 120
 taacaaagt cctcagacc tctggagcc acagaccaga ctctgaatg gacctggag 180
 gtgcactgt cccagatcat cgaatggat gaaaggaaac aggtgtctgac cctgtatctg 240
 tggatacggc agaggtggac agatgctac ctacgtatgg acccaatgc ctatgtgtgc 300
 ctggatgcca tccgataccc cagcagctct gttggggcgc cagacatgt actctatac 360
 aaagcgagc cggcgctcc aggtctcgc agcaacacg tggctctgg ccagatggc 420
 gctgtgctt gggacgcgccc ggttcatacgc cgcagctctgt gccgctgtga tgtagcagcc 480

ttcccttgc acgccacga ctgcggcctg acgtttcgct cctggactca cggcgggcac 540
 caactggatg tgcggcccg cgcactcga cgcagctcgg cggacttcgt ggaagacgtg 600
 gagtggcgct tcttgggact gccggcgagg cggcgctggc tcaactcagg tgcctgtctc 660
 tgacctacc ctagctctac cttcacgctg ctgcggcgcg tcccgctgag gtctgtctc 720
 tgaacctcga tctgtccctct gctgtctatc tgcgtctctt cgcgcctgc cttcacactg 780
 cctctcagct caggtcagaa ggtgtcgtc gggctcaacc tgcctcggc cttcacgctc 840
 ttcaatgtc tgtggcga gacatgcca cggcgacga ggtgcgct catcgggaag 900
 tactacatg acctatgac atgttccca ttctcaacg cactcacat cctctactg 960
 aacctcatt acgtgtgttc cagTgtccc cagTggcag catTggtag ggcctcctg 1020
 ctggacac ttgcacgggg cctgtgctg cgggaagag gggagccctg tggcgactg 1080
 aggcacact gttatcttc tagccccag tgcctcgaag gaggggctgg cccccagc 1140
 ggccttggc agagccacg atgtctgtc gccagggaag cctactgca ccaatgagc 1200
 acctatgca atactctgc cagccacga gtgtccacg gtgtccacg gactctgga 1260
 cgcctggcca atgtgtgga cgccttctc ctggccact tcttccat ggcctgtgac 1320
 atgcacccc tgtgtcgtgt gcaagccctg tgaaggctgg gactaagtc cagggactg 1380
 ctgcagccac agctctctca gcaagggcca gcaagggca agtggTgct ggtctttggg 1440
 ccagccagtc tctcccact gactctaaga tctt

<210> 46
 <211> 1742
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> misc_feature
 <223> Incyte ID No: 747395CB1

<400> 46
 tgggctgcg tccctgcctc cgcgtgcgag ccccgccgt agcgccctc gagcccgcg 60
 ccacatctct tggagcagag atggctgtgc caccacgta tgcgatctt ggcanaactg 120
 ccaggaggt gcttccagc ggcataagg ttacaagct caggctcagc caacacgag 180
 accacaaag tgcggcgag tctggaacc agtagacag ggaactagta cggctgtgac 240
 ttacagaga atTggaatc cgcacatca ctagcaccg agattactgt ggaagatcag 300
 ctgcagcg gactgaagct gaacttcgat tcaactctt cactcaacac tgggaaaaa 360
 aatgtaaaa tcaagacagc gtacacatta accTggcta cctgggctg gcacatggat 420
 ttgcacatg ctggcgcttc catccgggtt gctctgtgc taggtttacga gggctggctg 480
 ctgcgtacc agatgaatt tgaactgca aaatccgag tgaccacag caactttgca 540
 gtTgttaca agactgatga attccagctt cacactaatg tgaatgacgg gacagagttt 600
 gggcgctca ttacacaga agTgaacaag aagtTggaga cgcgtgcca tcttgcctgg 660
 acagcgagg acgTaaacac ggcTtcgga atagcagcca agtaacagat tgaacctgac 720
 gctgtctt cggTaaagt gaacaactcc agctgtatg gtttaggata cactcagact 780
 ctaaaagcag gtaltaaact gacactgtca gctctctgg atTgcaagaa cgtcaatgct 840
 ggtTggcaca agctTgtctt agactggaa ttTcaagcat aaTgaatc tctcaaatg 900
 tttaatttta aactatttg cagcatagct acctcagaa tttagtgtat cttttaatgt 960
 tgaTgtctg ggtTgaagt attgttaatt atgttagccc tccaggttaa agTgattca 1020
 gctttaagt gttaccttc cagaggtaca gaagaaact atttcaaaa aggtTccttt 1080
 cagTgttga cTcggggaga actTgtggc cctTtgaga tgcaggttt ctttttctc 1140
 taGaattggc tGaagtgga agcgataat atgtagcac ttgttaatt catattagat 1200
 aaTgaatga aattgtgatt tctgtgaat cgaaccttgg ttccctaac ctaattgatg 1260
 agaggtcgc tctgtgatgg tgtTaaaaa ctaacctgaa tgggacttt ttgacagat 1320
 ctctacacc tgttccccc ccaTtctat atcatctct ttacaccaa aggtTcagc 1380
 ggtTgtgtaa gTttctctt tgtTgcatt ttggggtggag aggtTggag tgaTgaagc 1440
 aaaaatcag gactattcc ttctgtgtt ggtTtttt ttggccttg caccagata 1500
 tgaatatagct tccagagct ccaagctaa gctTggaagt gctgtgtga ttgtaactac 1560
 atggTgaaca cactcagaa ctaaatTgga cttctgtgtt attctcaca ctaaatTgt 1620
 ttTttagag taaTgggla catTtagag tctccattt gttggaatta gatctctct 1680
 tcaagggtcg gattacacac ttaaaactg attaatattg acttaaaaaa aaggggcgca 1740
 ta

<210> 47
 <211> 2312
 <212> DNA
 <213> Homo sapiens

<220> miac_feature
<221> Incyte ID No: 7474111CB1
<400> 47
ccaagagaaag gctgtttttt gscgttgaca aggcgtttctt cccagagact tcaatccact 60
ctaagggcag gactctgaga ggttcacgcg gctgttaacg tttagggctt tcccagagct 120
cgagcccccag aacgtgaggg cggggggcgg gtcgggggac gcttcccttc gtcgggttcg 180
agggggagac gaggcttgag gattggggctt gtaagcgcctc tctgaagcag cygttaagttg 240
gtcgcgcccg ggggggggag gaagagcaaa gtcgggacga taagccggtt cgtgggtttgc 300
ttccctgaga aatcttagg ggcgtttccg gaaccgttgc tgggttcggg acaattttcg 360
gaanaagct tccaaagcc gtcgttcccg gacccggagc ccaacattgg ctttgagaca 420
ggcgttttaa aaggggccct gggaacagcc caattaccc cgaatggaca ggagagagag 480
cggaagagag calcccgcg cttgggttccc gttcctccc cccgtggggc cggatattcg 540
caatacagag cggagcggag gggagagagc gcttcaacag gggggagagc cgggttcagc 600
ggggggggcg cgggggagcc cgggcggcgc aagccagagc gcttcaagc gttcggggccc 660
ggcgtttggg agggggagac gggcccgctg tgttcgagc ccccgggcgg gagggttcggc 720
gagggagcac agccggcttc cccgcgggtt ggtgttcagc gtcggcgccg cgaagttacc 780
cgagccccc agcgtggggc ggtctcgggt gttctgagag tggggcgccg cggatattcg 840
ctgcggcgag agcgttgag gacttccc ctggcgcccg tgaagccggt gacggattcg 900
cggtcccgag ggcagtgag gacttccc ctggcgcccg tgaagccggt gacggattcg 960
ttcttcagac ggcactcgga ggccttcagg ttacatccc tcaagcgag tccctccag 1020
cgctgtcttg agcagctcg gctgaacctt gaggagccca cgttcgtcgt ggcgcggcag 1080
atccttgcat gcgtgtcggt ggtgttcgtt atcgtgtcca tgggttcgtt gtcggcgag 1140
agctgtcgag actggcgcaa cgcagccgcg gacaaacgca gcttggatga cggagacag 1200
abaatttag ctatctgcat agtttgttc actgcagat gcatcgtgag gttcatgtc 1260
tccaaaaa agtttgtt ttgtcaaga cccctgaca tcatgtatt actggcaac 1320
accagatatt acaatctgt ttgtataga gttgttacg gcgaagacac tcaatccag 1380
agggttgcat gttactttag gttactttag atgtataga ttltttgggt gatttaggt 1440
ggcgttgac tcaattggct tcaagacatc ggtttgacac tcaaacgttg ctacggatcg 1500
atggtttatg tactttctt catttgttc gcaattggca ccttttagtg acttttcag 1560
ctcttgaaac atggggctga ccttgaaac tccaaacag actttacag actttccag 1620
gctcgtcggt ggtgttgcat cctcatgac acaattggct atggagatc gtaatccatc 1680
acagtgtctt gaagatatt tgaagaggtt tgtttgttca gttgaattgt tcaattgag 1740
ttacacttcc cttttatcca ccaagattt gttcgaggtt atcaagatc caagtttga 1800
tccttgaggt atagtagag gcttccgac gaaatccctga ataaatgac tgaatgac 1860
ttcttgata tcaatctag aagaatttg atgtcttcc ataatatc gttttctgt 1920
gggttgagac tgcacttga ttgtatcat cttgttagag taanaattat ccttccagc 1980
cgaggtgata acaagtttca cttgtatag agtaataaga attgagacg caaagagaa 2040
ataatgacac ctgaagtaaa ctttagagc cggtttatt tagatttggt ttccgtttc 2100
cttgaatgat tacaatttt taanaatag atatttgaac caatttcaa caaagagaa 2160
ctattttccc aatgtttct cacttttga attcagagag agctttgagc ttatagttt 2220
tttgttttga gaggtaacat ttcatcttca aatgttttat aattttcat atcaattgta 2280
gaatgtacct ggaataacat gtcaacagcg ag

<210> 48
<211> 2320
<212> DNA
<213> Homo sapiens
<220> miac_feature
<221> Incyte ID No: 7480826CB1
<400> 48
ccgcgcgttc gtaggtctgc tccacagcgg cggagacgg ccgctttaga agccttttc 60
agtaactgta gaagagagcc gacacagctt taccagacgg tccggccttt ccccttaccac 120
agtgagagcgg gtaggattgg gtagggcaatt tgcattttcc ttatttatla aatatatttt 180
tccgtccgctt tcttgatttt ggggaaattt ttgttttgtt ctttatgatt tgaattttag 240
tgaagagaaac ccaactgaga cgtctcggtt agaatagaga ccaacggagc cgaactcgcg 300
ggcgttgacac caacggcaca ggaacaaagg aatccacagc taccagagcg acccgatctc 360
tcccagttgt tagcatgagc aagggccgaa ttggagcatt cagattttcc cgggattgag 420
aagagcagag ctacagttcc aacagagact tcaactatct ctaccacac aagcaagctg 480

ctctgaaaag ccatttgcga gatgtagatc ctgaacaaga gaacttttla cttgaatcga 540
atttggggga gaagaaagat gaacagagat ttcaacagag taactattcc tttagaatg 600
cagttattta tctgagcaat ggcatttggtt gaatggagat ccttgggctt tctatgcga 660
tgacttaac ttgaattctt cttttatlaa tctcttgagc attttgtca aatatcttc 720
tgattctgt tcaatccct ttgaagactt ccaatgagag aggggtcttla ttatgtaac 780
aatggagaa ttgaagcatt ggaattagtg gaagcttgc aagacttcgga tcaatttcaa 840
tgaagagct ttgaagcttg tcaagttacc tctctatgt gaaatagag ttgcctttg 900
tgatccagac attaaagagc attgaagata aaacctgatt gttgaatcgt aacggagac 960
aatgttttt gttgggttca ttgttggtca tctttctt gtcggttgtt aagaattttag 1020
gaatttttgg atabaccagc ggccttctct tgtttgttat gttgttctct ctgaatttgg 1080
tcatttggga gaattttcag gttccgttct ctgttgagagc gcttttata attaagagaa 1140
caataaacac caacttaca cagcaaaag cctttgagc tggttttcga ctatagagta 1200
ctgaataaga cctctgcaga ccccaatatt ttattttcaa ctcagagac gttcatcg 1260
tgcnaattct gacttttca ttgtcttgc atccgtcgtt tcttccatc tatgaaagac 1320
tgaagagcg cagcgtttag agaatatga atgtttccaa attttcaatc ttgttatgt 1380
tcttcaaga tctgtctgc ggccttcttg gatacctaac attttcaga catgtttatg 1440
cagatttct tcttaccac tcttctatct ttggagactga tatcttctt cctatttgc 1500
gctcgttgt tttaatggct ttgaacatga cagttacagt agttatttc ccaatccgga 1560
gtttctgtac tcaattgttg ttgtatcaa aagattttcag ttgttggtgt catagtctga 1620
ttaaatgac taacttggtg tttaacaaat taattgtcat cttgttccca actattggg 1680
ataacttttg ttltattgtt gtaactggag ctttcaatgt gatttttat cttcctttcg 1740
ccttctatct caagtttggtt gagaagagc ctatgaaac ttgaataag attgggctt 1800
tgcttctct gttaaatggt ttacttggtga ttgaacggag catggcccttg attgttttgg 1860
atttggtaaa caatggagct gtaggttggtc attaatlygc accacttcaa cttcaacttc 1920
gtccacttga ttgcagtggt gagttaactc taactatlyg aaatttccca taattgttc 1980
agtttactc ccttttgag ttatttttag ttattttag aaactttcat gttatgtga atgtagag 2040
aaacttttg ttgttttg ttatttttag ttattttag aaactttcat gttatgtga atgtagag 2100
aacaacaaa cccagcgctt cgggtttagg aaacttttag ttattttca tcccgagaa 2160
tggagacac cttaacatt attcaagcac atgtcagctt gttatgtgt ttcaatttga 2220
tatttttaa tttaatttga atgttgcaaa ttgaacacat ttgatttga aaactttct 2280
tttaatttcc agcaaatatc ttgaatacaa ttgaacaaa

<210> 49
<211> 1781
<212> DNA
<213> Homo sapiens
<220> miac_feature
<221> Incyte ID No: 6025572CB1
<400> 49
gtcacattct cggcagtagc atgtctgagc gtttttccgg ttcccgctt cccctcaatc 60
tagctccgct actcaatttt agcacattgt ggcggttttt atactctct ccaatlygaa 120
tcgttgagctt gcgaataaga agcagaataa ggcgcttttt gacgctcat ccaatlygaa 180
ggactttctg ggcggcgagg ttgcggcgagc ttgttccagc aaggggttgg cggccatcga 240
gggggtgag -ctgcgtcgc agtgcagagc gttcccgagc caatctacgc ccgaagctg 300
gtacaagagc atgttgagct gcttgggtgc gattctcgc gacgaggttt tctttagtt 360
ttgggttggc aattttgcaa atgtatttg gtaattcgc acacaagctc taaatttgc 420
ttttaagagc aaataaagc agttattct gttctgagtt ataaagaaga aacagtttgc 480
gaagttgtt ttggcaaac ttgcttcttg ttgagcttgc ggggcaaac ccttatgtgt 540
agttatcct ctgaattttg cccgaacggc attaagtttc gaatttggaa aaggttcgga 600
ggagagcaaa ttcaagaggt ttggttgatc ttgattgaa atggaanaat cagatlygag 660
ttgttggttla tacaagaggt ttgttttgc agtaagggc atcaattgtt accgagcct 720
ttatttga gttatgaga cagtttagg ttatrtacca aagcaaga aaactccat 780
tcgtttccc ttltttatly ctaagttgt gactatctgc tctgaaatc ttctttatc 840
atttgacaa gtttgaagc gttatgtat gcaagttgtt gaaacttcaac gtaaatatga 900
aggaacata gactgttttg ttgaagata ccaaatlyga ggaactcagtt ccttttttgc 960
ttgggccttc taactttcc ttgcggtag aggggttctt ttgttggttg ttattttaga 1020
tgaattttaa gaaattcttc atattgact ttgttggttag ttatttggag agtaaatga 1080
gaatracag gatttcaact gttacaaca caaattacac aggtccatc ttgataact 1140
ttgattagt ttatgtctgt attttttaa agttcgagtt ctgcatatga gaattatct 1200
ttcaagat tttaatacaa aaatcaagc aatgttgat ttctcttcca cttagatcca 1260

aacacatttt agtgtgatatt ttaatttatt ataggtagta tattttaatt tgttagttta 1320
 aaattctttt tatgatataa aattaatatt ataatcttag attaatgctg aatatagga 1380
 aatgaagaatg gcgtctttta aattgtatt catttaatt acctgttttc catcttttg 1440
 aagtcataatg gatgcataat ttcttaaaa ttctataat agatgtcaat atagtgtag 1500
 gcgaataaa gcttgttctt atacttctt taagacagtt gtattactg tgaataaat 1560
 ttcaagtatc agcttgttat tatagtatg atcattttaa atttgataat gactttagt 1620
 acattataaa actgaactg gaaaataaaa tggcttatct gctgatgttt atcttataa 1680
 taataataat ctgtctagt tgaataatc ttgaacaaa aggtatcttc tgaataata 1740
 gttgtatat ttgttgaca ataaaggaag cttaactgtt a 1781

<210> 50
 <211> 2433
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> misc_feature
 <223> Incyte ID No: 5686561CB1

<400> 50
 catccgctca caatgcocaa tcaatgatc gagacagtag cctcaactgt tgcacagtc 60
 atggcagat cggctgcgag caggcaggtt ggctgaggg aggtcaactg gctggtgag 120
 ggtcttttgc tgtgtttctg agccggctct ctccaggca cegtgtccat ggggtaagc 180
 ggtctctctg gttgcceact ctgcccgcg gactctga gtttgtctt gctggccat 240
 gaagctcagc ctgctgggag gccacaggga gatcagaggt gggcggcggg tgaagggtc 300
 cagccgggttg gttccggggc ctggagctca gctctgggg tggggaccoc ctggagctt 360
 ggaagcgctg gttctctgtt ctgagctatc cagcagatg atggcaggga caggagaggg ctgactact 420
 tcagaacact cctgtagtct ctgactcc cctgtgtct gctgaccac gccacaacc 480
 ccagatgat gattctctg tattcccaa accggccta tgcactctc tcaatagct 540
 tcaactgtat aggaacctg ttctatga acctgtgac agccatcatc tacagtcagt 600
 tccggggctc ctgagtaaa tctctcaga cctcctgctg tggggagcgg ctggacacc 660
 gggctccttt tgaagctca tctctcatg ttggggaggg aggaagcttc ctccaggag 720
 ttgggttgaa gccccaac tctgtcagg ttcttcagaa gttccagtg gacagctcc 780
 acaaacggc catgatgaa aaggtgcgtt cctacagac tgttctgtg tcaagtggg 840
 agtttcagaa gctcttcac gagcttgaca gaagtgggt taagaagcac ccgcggaggc 900
 ccagatgaca gctcctgt ctgcagagcg accagttct cctggccac tactactttg 960
 actacctggg gaacttggct gttgaccgat tgcacaccc aggttgagg cgggagatgg 1020
 tggagctgtc gtcgtgtggt gactgggca acctgtgtc catttgctg tctcgtgtc 1080
 tggatcaga tgtgtcgtct gctgagcgtg atgaactat cctggggatt ctcaactgc 1140
 tctctattt gactactg ttggagatcg tgcataagt cttgtccttg ggcctgcag 1180
 ggtacactgt ctacccagc aagctgtttg accggctct caccgttgtc ctgctgggtt 1200
 ttgagatctc aaacttgttt agaggcgta ttgtgctt tctggaaac agcagctctg 1260
 tggagctgtc gtcgtgtggt gacatgacc gcatctgaa catgtctac gttctcgtc 1320
 tctgtgat catccagc atgaacgca tggcgtggt ggcagatacc gtcctggggc 1380
 ttgtgcagaa catcgtgtct ttggcgggga tctgtgtgtt ggtctactac gatttgcca 1440
 tcaattggat caacttgttt agaggcgta ttgtgctt tctggaaac agcagctctg 1500
 cccctgccaa tggctcggcg cctgttgga gcttcagca gctggagta tgggccaaca 1560
 acttcgatg ctttgcgggt gctcgtgga gcttcagca ctgtgatg gtaacaaact 1620
 ggcaggtgt tctggatga taccggcgt actcaggct actcaggcc gtgtcccaag atctatttg 1680
 ttgtgtggg tgggtgtgcy tctgtcatg ggtctaaact gttctgtgct ctgattctg 1740
 agaaactct tcaaatgtg gccccgca gccactgca gcccctgtct gggacccag 1800
 aggcacacta ccaagtgact gtgagctcc gtttcaggga tatictgag gaggccggg 1860
 aggaatgct cacagatg ctgagcagc acccgact gtggtgtgc aggttactc 1920
 cgggtctcgc tccacagag ggcggcagga gagagaggtt ggcctacaa ggtggccact 1980
 atggagagg ggtcagatgt gtcggcagcc aggcaggaa agacctctc tctgagac 2040
 caataagctg gggacagaaa caaagctctt tgcgtgtg ccacaacca tctacagac 2100
 agtgggtgt gtttcagga ggcgcgctg cctcgcctt ttcttatag tgttcaagt 2160
 agaatctct cgtcagct acccagac ctccagct ttcagaaaa aatgaagaa gagcggtct 2220
 cccctgtcc ctgcagcttc cgtgtgtct tctgtccgg cagccttg gggacacag 2280
 ctgcagagg gctgcacag gttaacctg agactccgg ggcactcagg tgggtagct 2340
 ggtgggttga catggagaa acctgactg tgtttatta ttctatggt tgaatagtg 2400
 tgaatgggtg tttttctta gggttctgat tgc 2433

<210> 51

<211> 1772
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> misc_feature
 <223> Incyte ID No: 1553725CB1
 <400> 51
 cccacgctc ggaactgctg gcattttcc sggaaccagg tccacagtt tatgtgtgag 60
 caagatggag gctgacctgt ctggctttaa catcgatgcc cccgctggg accagtcac 120
 ctctctggg agagtgaag acttctaaa cctcacggac ccccgactg tctttgtac 180
 tgaagggag ggtgactggg ccaagtgat ggtgaagag agcaggatgg ggtgtgtgc 240
 ccagggacc caagtggag agctgtgta tgcaagaag ctgtatgact cggcttcca 300
 cccgaact ggggagaaga tgaatgcat cggcgcatg tcttccagc tctcgggg 360
 ctgtacatc acgggttcca tgtctcagtt ctacagagc atgcggcg tgatctctg 420
 gcagtgggtg aaccagtc tcaatgctt agtcaact acacaagg atgggctc 480
 cccacatca gtcaggcaga tggcccttc ctacttaca gccacaacca ctgctgtgc 540
 caggctgtg gcatgaaca tgtgacaaa gaaagcgcg ccttggtgg ccgctgggt 600
 gcccttgcg gctgctggct cggctcaact gttcaatc cccatgatg gacgaagga 660
 gctcataag gaaactcgc tgaaggacg gaatgaaat gagattggtc attccggag 720
 agtctggcg atagctaca ccaagtagt talttctcg atcaactgt cagctcgtg 780
 gatgacttg ctgcagta tcatggaag gcttgagaa ttgcactca tgcagaagt 840
 caaggtctg cagcccat tgcagttcat gctgagcgg tcttctcca tctcattgt 900
 gccagtggtg tgtgggtctt tccacagaa atgtgaatt cagtttctc attctgaaac 960
 gaagctcaa gacatctca aggcacagta tggagaactt gaggcttat tctactcaa 1020
 taaggttct taaatgccc acttcaagaa gacacagct attcccat tcaaccag 1080
 cctctagct agtgcacac tigtgtctc ctcccttt gccacaagg cctgaagcc 1140
 aggttagat ggggggtggg acaatgaat cctacactt acacctgtt actggttat 1200
 tggacatcg gggaaaaag tgaaaaagg tagcaaggc caatgtctc tagctgttc 1260
 caatacctt gctccttga gaccagac tgggcccc tccagggaga gacatcaag 1320
 caactattt gaaagctta gaaaccttt agaatcttg tccagccag ggttgagaa 1380
 aagacotgg atcaaaaga agcttctata cctcttctt ctgtctcc tctctcca 1440
 gaaatgaa acttttacc atgtaatct agtgaact agtgaactc agggaaaaa 1500
 gaattgtcc ctttgggtt catcactt ccaactct cctctgtt cccctgttc 1560
 agtgcgttc tttttttt ttcaagatg agtattgtc tptgcscag gctggaatg 1620
 agtggcgga tctcgtctca ctgcaactc tgcctctg atcaagcga tctctgtgc 1680
 tgaagcttc agtgcgttg gattacagg cactgtccac caccgtctg caattttgt 1740
 atttagtgg agacgggtt tcaactgct gg 1772

<210> 52
 <211> 1874
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> misc_feature
 <223> Incyte ID No: 1695770CB1

<400> 52
 atctttcca gctccagct ggcaggctaa ggcctccgg agcccaaggc gagcccaagc 60
 agaaccagt ggtgttact ggtcaggat catttccagg ggaatgttc tggccctgg 120
 caggtaaga tggccagag gagaaggagc agaaagcagg aagaatgtc ctcttttgcg 180
 agacgccag gttgagaaa ggtctgtac ttittggctt cctgaggga tatccctct 240
 cacattgga gccagctga gaaagctct caagatccc gcagaatgac aactctgtt 300
 cctgcaacc tctcttctt ttctcttgg acctgcag ggcaggtct cctcaggggtg 360
 gcttggcaa agaggaagt caaatctga accaagggtt cccagcccat gtcctctct 420
 gattctcag acaactatt gggcgcaaa tctgatat atgcaggat tggcccaat 480
 tttaaggcc caactgtga cgtgacctc aacatctca tcaacgttt cagctcgtc 540
 acaagacca caagctgaa cgggtgaat gctcttgc ggcacagtg gaatgacca 600
 cgcctctct acaggaata tctgtgac tctgtgacc tcatctctc catgtggagc 660
 tctatctga agccagact ctitttctt aatgaaaag gggccaact ccatgaggtg 720
 acccggaca acaagtact ggcattctt aagaatggga atgtgtgta cagcatcagg 780

ctgaccctca ttcttgcctc cctgatggac ctcaagaact tcccacatga cctccagagc 840
 tgcacagctc agcttgagag ctttggttac accatlgagag acctctggtc tggatggctc 900
 ggaagctgct cctgtgtcca agtggctggag gggctgtgct tggcccaggt tcatctggcg 960
 gttggagagc atccatggctc ttgtatccag ggtctgtgct cagggaatct cactctggag 1020
 gggggagagc ttccatccga accgagatga ggtctatct tgaattcggt tcccctggcc 1080
 accatccatca gttccatccc gttctggctc cctctctgga tcaacatgga tctgtgctcc 1140
 ggtccgtgtc gcttggtgat caccacgggt cctccatgaa ccaacagagc cttgtgctcc 1200
 cgtgctcttc tgcctgaagt gttctatggt aaggatctgc acctctggt ggtctgggtc 1260
 gttctgtctc ttgtctgctc cttgtggag ttgtgtgcca taattctgtc tcccctggag 1320
 caataagagc tcatagagct tggatggag cagagtgccc aacctgtgct gaaagcatatc 1380
 accatggagc gttctgtcata ttccgtgtgc ttgtgtcttg ggtctgtgag gttatggatc 1440
 gtttgagagc caattggagc ttctgtgctc tatagctccc aacctccag cctccctctca 1500
 agggagagag aacacagagc gaaatctcac gttgacagag ccaagagatc tgaacacatc 1560
 tcccctgggt tcttcccttc cacttctcc accatccata tcttctatgc gttgtgctct 1620
 aagatggagc gttctggagag taccacagc gttctggag ttgggttgga gttatggatc 1680
 cgtgtgtctc gctccctgct tccctctggag ttgggtcttc cctccatgag gttccatata 1740
 ggggtgtgga agttctcttc ctgactctcc agctggagct tcaactccca gttcccaagc 1800
 tatgtgggct tctatgtcat gttgccaatc gttgctgtac ttataagatc gttatctcac 1860
 ccttaaaaaa aaaa 1874

<210> 53
 <211> 6211
 <212> DNA
 <213> Homo sapiens
 <220> m1sc_feature
 <221> Incyte ID No: 4672222CB1

<400> 53
 ctgacggagagc ggtgctgggc ggtctccctc ggtcagatgac cggagagagc ttgtcggaca 60
 attatggagagc tgcctgcttc tgcctgtgac gctggagagct agttatgttc ttgaagagtt 120
 cctgtgtggt tcttgtgctc gctgctctc gctggagctc cggagagctc taccagatgct 180
 tggcagagagc cctggagagc ggtggagagct gatttgtgtc ccttcggagc taccgtccctc 240
 gggagagagc cctggagagc cctggatgga aagcatttc accaagaggg aatgtgata 300
 tcttatcaca agttctgctc gctgctgtc caggatctac ggtatgtaaa ttgtcggaca 360
 acctgctgagc ttgtctgtc gctgctgtc caggatctac ggtatgtaaa ttgtcggaca 420
 tgcctatgga tctcatagct tgaattgttg tgaacatttc aatcagagca ttgaagagtt 480
 gttctgtgga aagcatagag aacagagctc aagatgtct ttgtgtgca taaattctca 540
 aggggttctc cctctccaca ggtctaaatc ttgtgtgca tcatatgaca ccaaatctga 600
 aactcttcg cactctgca ttgaagagct tgaagatggag ttaccacagc ttgtatctca 660
 ttgacatctg ggtcagagca aatttgagct taccacagag atccagagc ttgtctgaaa 720
 aggtctatc aagctctgag ttacacagct agctggatc ttactggag ttgttaaac 780
 aggtgtgca aacatgttc ggtatgctc caaagaactc gcttcagat atctctgaa 840
 gatttgcct atcggaatag ctccatgggg agttatgga aagaagatc atctgttg 900
 ggaagatctg gttgtctct atcaaatct attpagagc ctgagagcc ttgaatttt 960
 ggaatctc cactctat tcatatgct ggtatgtgc actgttgaa agttgtggc 1020
 ggaatctc cttggagag aacttgaaa aactatcat cagaagaa ttcatctg 1080
 gattgtgca ggtgtctc ttgtgtgac tcatatgct gttgtggca atgtatctc 1140
 cagatctct gattctctc aggaagagc cctgttcca ttgtgtgtc ttgaagagc 1200
 agggagagct ggaatctgc ttgtgtatc tcatatgca aagaagagc ggtggagct 1260
 ttctgtgca gttggagc atatatctc cactatcaa aaaaattca actttgca 1320
 gatttgagca cttcatctc ttcaacctc gattggagc atgaagagc aggtgtctc 1380
 cactgttctc caatgtgtc cagatgaa ttcaagatata gatttgaca tacttactg 1440
 accatgaaa gttatgagc cactgagct ttgaagagct atctctcat ttgtgtgg 1500
 ttgaatgagc atgtggaaa atcatgtat ttgtatgca cagatgtgc ttgtgtgac 1560
 cttggaaaac gttatgtctc atgtctgtc aatgataga gttgacttg taaattctc 1620
 tatggaaa ttgtatgca ttgaatctc ccttactc cctgagagc aggtgtgag 1680
 caacatcaa caagttcca ctatccatc gttgttcat cttgtctg agtcaaaa 1740
 ggaatctc cctcagagc ataatgacg ttgtatgat atagagctc ttatgtgaa 1800
 tctcatggga ggaactaca gttacacta ttactgaaa cgtttgat taaattata 1860
 taactgtct ggaataac ggtgtgtctc cgaataac tccagatcc cctctgact 1920
 gctgaagagc catgacttc ttgtcatagc ggtcatgaaa aagaaaaaa ttgtgtcata 1980

ccattctatc aagacagcac agctctacc accaagatc gatcaagta ttgaagagc 2040
 aaagaagaa agaacagagc atgaatctc agaatgtat gatcaagaa ccaagagc 2100
 tcttatcaca cttatggagc tttaattc gttgtctc atgaagagc aggtcatg 2160
 ccgttttca ttgcaagagc gttgaagac gatggagc ttatgtatc cctgagagc 2220
 cctatgtc atgtgagagc aagaagaa gattgagc ttgtatgta cttcagagc 2280
 actaaagagc ttatcaagc attttgca gttgtgctc gttatgtatc aagatgct 2340
 cagagagagc gaaacagc ctatgaatc gctcaatc gaaatgagc atgtgagaa 2400
 cttaacagc ttgaatgagc atgttctc agatctagc cctttgtgag cttacagct 2460
 ttcaagagc ttgtatgagc atgtgtgctc gttatgtgagc aatgtgagc 2520
 gttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 2580
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 2640
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 2700
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 2760
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 2820
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 2880
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 2940
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3000
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3060
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3120
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3180
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3240
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3300
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3360
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3420
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3480
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3540
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3600
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3660
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3720
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3780
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3840
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3900
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 3960
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4020
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4080
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4140
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4200
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4260
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4320
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4380
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4440
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4500
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4560
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4620
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4680
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4740
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4800
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4860
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4920
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 4980
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 5040
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 5100
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 5160
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 5220
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 5280
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 5340
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 5400
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 5460
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 5520
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 5580
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 5640
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 5700
 ttatgagc atcatagaa ttatgagc ttatgagc ttatgagc agttatgagc 5760

tgagcctta gattgaatc ttacgctgg aaattccacc aaagaatcac aatcaactaa 5820
 ttctgtctgt ctgattgita aatattaata ttactgaatc attgtgtttg ctgacactc 5880
 acgaaatgt tactgtgta cttctctcc ttggttgtaa tatagaagg 5940
 tglatcaag tgaatttcg tgactccag acagtttaaaa ggtcaattt ctttgaact 6000
 gattaatcac tcagaaggc cctatagat agagtcggca gctagaaga tttaaggta 6060
 atgataatt agtatattata actttttaa gggctctttg tatagaagag gttcattt 6120
 gactttgttt tgatggggt agtgcctct ctatgtgtt acaataccat taaccaagg 6180
 taggtgtaca tgcagatttt attggcagct g 6211

<210> 54

<211> 3714

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6176128CB1

<400> 54

atggcgggg ccaagctgc gcctcgccg tccagaggca agcgggggcc gggggggccc 60
 ccagcgggg ccgagcccc cgaggagctc cagggctca gccctgctt gccggccgc 120
 ggcggggggt ccgtggggcg cgactgggc cagagcttc ctgtagaaga tttagcctg 180
 gactctccc tgtctcagtt caaggtggag ttctacgtca acgagaacac ctccaaggag 240
 cgactcaagc tgtctctat caaaacaaa agatcgagtc tgaggatccg gctgtcaac 300
 ttctccctga agctgtctac ctgcctgttc taactgtgc ggcctcgtct gcatgaccg 360
 gccctgggca tcggatggty gggctgcca aggcagaact acctctcaa tgactctcc 420
 tccgagata actgggtctc tattctgtg ttggagagaa agatgacact gggcggtac 480
 caggtcatcg tggccataa agcttctcg gagcagtc ttctatcta cctcaagctac 540
 aaaggcaaca tctggagata gacttccg gtgctctcg tectggagat gatcaactc 600
 ctgcctctta tcaactgact ctcttgggcg ccgtctgga acctgtctat cccgtcttt 660
 ctgaaactgt ggtgtggcca gacgctgtc gaataatga ttaagaatt caacgtgtc 720
 atctctggca actcagctca caggtctca tctctctca tctctcttg caccgtgtg 780
 tgcctctgt tccggggg ctgcggatc cagcaactcg agcggggcgg cgagaacctg 840
 cctctctga cctccctta ctctgcatc gtaacttct ccaactggg ctacggctgac 900
 gtacccctca agatctggcc atcgagctg ctgactgtc tcatgtctg gctggcctc 960
 ttgggtctcc cactcagct cggagggctc gtaactctt ggaatggagc gagaagtca 1020
 ggggggaaat acagcgcca ccgtggcgag acggagaagc acgtgtgctt gttgtcagc 1080
 tccctcaaga tcaacttct catggattc ctgaacgagt tctagccca ccccggtc 1140
 caggactatt acgtgtctat ctgtgccc acggagatgg atgtcaaggt ggcagagtc 1200
 ctgcagatcc ctctgtgtc ctaggggtc attactccc agggctctgc actcaagac 1260
 caggactca tggagcgcaa gatggcaat gtagggatc gtttactct cagcagcagg 1320
 aacgaggtgg accgacggc tgcagaccac cagaccatc tgcgcgctg ggcgtgaag 1380
 gacttgcgc ccaactgccc cctctagtc cagatctca aactgaana caagtttca 1440
 gtcaatttg ctgaccactt ggtgtgtgg gaggatgca agtacgcaat gctggcgctg 1500
 aactgcat gccggcgag ctccacctc atcacctgc tggtagaac gtcocgggc 1560
 caggagggag agagttctcc gtagcagtg cagcctagt atggcgctg ctccggcaac 1620
 gagggtgacc acatccgat ggttgacag aagtcttcc gcgagtaga gggcaagagc 1680
 ttaactacg cggccttca cgccecaag agtatggcg tgtcctcat cgggtggaag 1740
 cggggggaca taagaactt cctgtgaac ccggggccc ggcactctt gccgcctct 1800
 gacactgtt tctactca catcaacga gaggagaact cggcttcat ctcaagag 1860
 caggagggag agagttctcc gtagcagtg cagcctagt tggcgctg ctccggcaac 1920
 ctgcctgtg acagactat gcctctca ggaacagtg ccatgagct gaggcgca 1980
 gagcccgcc ctacgcagag cggcgtggg ggcgggggca gcaactggc actgcccacg 2040
 gagaacggct cgggcgggca gcgcccgag atcgcccg tccgggaact ggcgcctct 2100
 ttagccctg tgcctctcc cctgctgag gaccagtgg aggaagggt ggcgcgtcg 2160
 gaggcagg ggtctccgt gtlagagtat gtagaggtt acctctcaa cctgcctac 2220
 atcggcagct ccccaactt gttcacctc ctgccttga agccctct ctgctgctg 2280
 cggcttgaca agggctgcaa gcaacacag tatgaagacg ccaaggccta cgggttcaag 2340
 aacagctga tcatgtccc ggcagagagc gccggcaatg ggcgtgaca ctcatgtcg 2400
 ccaactcgg cctactacag atcccgaag gactgaacc ccaatgctgt cttgtggag 2460
 aacagacct accaactt cctggagcc atctgtgct tcccatggt ctactacatg 2520
 ggggtcttg tggacaagt ggacagctg ctgactgtg gcaatata tggcgacaac 2580
 ctgctctgtg tggacaagga gacacactg agcgccgagg aggaactatc ggcggagccc 2640

aagaccatg tcaactgtca gacctgttc cggctcttc ccagctctcag catcacca 2700
 gagctaccc acccttcaa catcgcttc atgcagttcc gcgcacaagg cagctactct 2760
 ttggctcttt caaactcaga aaaggggag cgagaatg gctcaacct ggccttcag 2820
 ctcctcgc cgttcgcgc cggcggtc ttcagatca gcatgttgg cacactgtc 2880
 taacgtctc ttgtgaagg ctacatgat accataccc ggcgtgctg ggcctgtgac 2940
 accagcgcg cgtgggcta cctctgtcc atgaatac cggaggcgga cctgtgata 3000
 cgaactgag gcgcctctt ccagaagtc tgcctcca gcgcgagat cccactggc 3060
 attacacga cagagagca cgtcttccc accctggag cccacgaact cagagcccg 3120
 tccagatct cgttgaact ggagactgt gaggacac ggaagtga gggcctcg 3180
 ggtcccgcg ctggcacgg agcagctcc caggcgccc acacggcgcg cgtgaccc 3240
 gcagacacc cactgctac gccaagagc ctgcaagtc cccggagct gagcgaag 3300
 ggcgcaag ccagcgccg ggcggggc cgaggtgga tcaagccag gcgcctcagc 3360
 ctgtaccgc cctctgagc cagagctc tccagctgg tgaagaacc catgaagcac 3420
 ctgggtgc ccaccacgg ctacagggac gtacaaatt taacggccag tatgtcatg 3480
 aatcggtlaa actgggata ttgcaagac gagatgaag accaccaga cactctcc 3540
 taagtctca taacctccc gccagacg aggtggag ccagtgatc tgttatctc 3600
 atcgtctcg taactggc tcaegtgc ccagctccc agagccgaa agcagctgc 3660
 agccaagc tgtgtctct caaccgag actcgagc agacacagct ctga 3714

<210> 55

<211> 3115

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7473418CB1

<400> 55

atggctcgg cgtcagacta tgttccaag tctaagctct tctgactct gtcgtcacc 60
 ccctctcgc tctgtccact cgttaactg atgcgcga agttgtcag gtcgactac 120
 gtaactatc taatggccat ttactgtgc acagaagta tccctggc ttgactctt 180
 ctatgcctg tctgtcttt ccactcttc cagattctg actcaggca ggtgtgtc 240
 cagtcagta aggaacaaa catgcttcc ctggcgccc tcatctagg cgtgctgtg 300
 gacgtctga actgcacaa gggatcgcc ctgcgcgc tccctgggt gggggccag 360
 ctcgcagcg tgatcgtgg gttctatggc gtcaagccc tccctccat gtgagtcaa 420
 aacagcgaa cagcgccat gatgtgcc atctggagg ccatltgca gcatgtgaa 480
 gccacaagc cagccacga gcccgctg gagctgtg acaaaggcaa gcccaaggag 540
 gtgcageta acagcgtgt gccacaca gggagtaag tgattttga aggcccaact 600
 ctgggcgag aggaagaca agacggag aggttgtga aggcatagc cctgtgact 660
 tgtaacggg cagatcagg gggcaccgc acctacgc ggaaggagc caactgtgtg 720
 ctcttgccc agatgaaga gttgttccc gtaagcag acctgtgaa ctltgtctc 780
 tggttgcat ttgcttccc caactgtc gtagctgc gttcgccg cgtgtgtgc 840
 gaatttgtt acatgagat caattttaa agtccctgg gctcgggct agagacaa 900
 aaaaacaga aggtcgccct caaggtctg cagagagat accggaagt gggggcctg 960
 tccctcggg agatacaagt gctgactgc tctctctgc ttgtctctc ttgttctcc 1020
 cgagccccg gcttcatgc cgtctggctg actgtgtcc ggttgaggaa aggaanaa 1080
 ceatttate cccctccc gctgattgg aagtaacc aggaanaag cccctggggc 1140
 atcgtgtgc tactggggg cggatttct ctggctaaag gatcggagg cctggggctg 1200
 tccgtgtgga tggggagca gatggccc ttgcagcag tgcgccggc agccatcac 1260
 ttgatttgt ccttgtcgt tgcctgttc actgtgttc actgtgca caagcaact gcccacac 1320
 acctgttcc tgcctatct tgcctcatg tctcgctca tggcctcaa tccgtgtac 1380
 ctcatgtgc cgttaccct gactgtccc ttgcttca tgtgtctgt ggcacacct 1440
 ccaaatgca tctgtttac ctatgggac ctcaagtt ctgactgtt gaaacagga 1500
 gtcaatga acatattcg agtcttgt gttgttttg ctgtcaaac atggggagcg 1560
 gacattatg acttgatca ttctctgac ttggctaat tgacatat tgactgtag 1620
 ggaagacc acagacacac acagacct taccctcc aggtactcg accttctg 1680
 cacacttgt acagatttt ggggttaca ccccaaat acccaacat gtccacac 1740
 caccaaac cagcaatgg gccacctt cctcaagc cagatgaga gatggacatg 1800
 ggcagtcga ggttaggtc agaaatga ggaacctc agtgggtgc tggaccatc 1860
 ttcccagc cttggcata tctctgag gaggcagg tagccaggg atcaggatgc 1920
 aggtctct agccctgc cctcaagat cccacaca gggctctgt tttaactgc 1980
 tctgtctagt atagttaa tgggaatcgg atccccgtt tgagagctaa gacaacac 2040

laccagctgc cagctgacct ccagctacac tggagagacc tcaagatcacc tctgtcacc 2100
tggagagagac acccagaccca gggagagagat ggcctgtacct gggagacacc ccaattgttg 2150
agctggagctg ggaatccagag cctctgtmag cctctctggag cctctctctcc ggcacacacc 2200
aagagagagag gaaatccagag tgcacacacc aagagagacc ctgtctgtcc acccaagctg 2250
cccccagagcc caactccagag ccttccatcc tgggttctgt tttaattcag tggccccctg 2340
tcccacaccg ccagagagacc tccatccacc cctgtgtccc ctgtgtctgt gttcaacatc 2400
tcccagagcc ccttgggtctg caccctctgg gcttccccct ttcccagag cagctccatc 2450
ttctctggag agctctcccc tgcctcccc aatatggatg gataccacc caaatatggag 2500
cagttatggc gctcgggggg ccagagtgag ctatctaggc cctccctcac ctggagacca 2550
gggtggagac agctgttaac tcccattgg tatggacct cagatctt calccctggc 2640
ttggagctcc tctgggttaa acctctcc ttgttacct cttccagag calgtctgtg 2700
ggacacagag gattcttgg altggagacc gcttggatg cttccagaa acaacacctg 2750
cccgctgggc aatctactg aagtaccg gtaacagac ccaagaggg gaaacaacc 2820
cagagagggc gtaggaatcc tcaagagacc cctgtttaga gaaagatag agagagagca 2880
aatctcttc ccccgcaagc ccaatttat tcttctctgt gctgtcttgg aagagagagca 2940
ttagaagaaa gattagagccc tggatggagc ttccagaggg aagagagagc altggagacct 3000
cagagaaacc catcagagagc agatagatg tctgttctct ttgggtggagc gattctcgagc 3060
cacactgtcc agctctgtcc ccaattaat gctctgtccc ccgctggaaa aaaaa 3115

<210> 56
<211> 2846
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 7474129CB1

<400> 56
tcccagagcc cctctgtacc gaaatggag ctctccggcc cctcagacac agtcccaccc 60
ggggagcttg gggccagagc altgctggat cccagagagc ggttggagca ggtctggacc 120
gacacaccca ggaatggagag cccacaccca ggaatggag cctctcttgg gaaagagagc 180
tctctcccc agtgggagac ctggctgtcc gcaagagagag agggccggcg agtccaccc 240
cccagagag agtactctgt gtaacttga tgaatggag tcccacagc ctccctcagag 300
tbatctggag gaaacccat ccaactccca cagccacagc gaaagcttgg ctggggctgg 360
ggagagggag aaaaagagag ggtcttggaa ggtcatcttc gaaagcttgc ctggggctgg 420
cgttggagag ttgttagatc tctgttggaa gttgagagc ctctcagagc gggccctga 480
tgaagatctg cctgactcc tcaatccca gttcggggc tccagacagc ggaagacctg 540
cctgtgttgg gctgtttaa acaatccccc caacacagc aagatctggc agatctgtcc 600
tgccttctgt gaaagagagc acaatctggc aggtcttacc aagagcagc agacagagga 660
ggcctatgaa gggcagagagc cgttcaaat cgcacatgag cggcggagagc ggtcttcaa 720
agcctctgtc atggccggcg gggccagagc caagcgagc gccaagggcg cctcttctcaa 780
cccacagatc caacagagag gcttctactc ggttggagc cccctggccc ttggagatgg 840
cacacacag cccggatltg tgcagctgt gattggagac gaggagagag acaatccacc 900
ggggagcca cggagagaaa acaatcttca cgccttgglt accgttggcg agaatctcaa 960
ggcgagagat gaaatttggga agcgatltc ctactgttgg ctactgttgg gttgcaactg 1020
ggagctggag acaacttcca acaacagatg cctcagcgcc ctgagcttgg ccgcaagat 1080
gggcaagggc gaaatcttga agtaacatc cagttcttgg altcaagaga aggggtctcg 1140
gaaacttggc aagaaatcca cagactggag ctgaatggc gttcatctc cctcttctga 1200
cctccacac gttgagaccca ccaagagaaa ctgaatggc gaaatctgt tctaacaaac 1260
gaaagtggag aagtttggca agtaacatc ccttctgtcc tctgtctt attcttctaa 1320
caactctccc ctgaacctcg tctcttacta cggcccccgg gaggagagag ccaatccgga 1440
ccccttggcc altgagacaa agatgggttg gcttggagtc ctggagtgga ttgttggct 1500
cacctgggac altgagatc ctgttaagaa gggcatlcc acctctcgcc tgaagacctg 1560
ggatctggag tcaatctctc cggatgtctg gttcaactt gttcttltta tccaagtgt 1620
gctgttgata ctgtgtctc tctgttact gtttgcctac taagagagc tggctctgt 1680
gctgtgtggc altggccctg gcttggcgaa calgtctcac taacggggc gtttccagtc 1740
cagtgcatat taatagttca tgaatcttgg gttcatlttg catgalttg taactctgt 1800
gtttgatata atcgtatltc tggatlttg ttgaatagc ttggcttggc taatggagaa 1860
gtttccaaa gaacaaagc actgagatc ctgaagcgcc ttcaagagc caggtctgga 1920
acctctcaag ctacacagag gcttgggtga cctgaacac cagagagatc caagatctcc 1980
catctcttct ctgtcttggc tcaatcacta tgtcatcttc accttgttc tctctctcaa 2040

catgtcatat gctctgagtg ggaagacttg ggaagagctc tccaaagaga gccaagcagc 2100
ctgtgtcttg cagagagacca ggaacatctt ggaatttggg aaaaatttacc caaatatggc 2150
ggagagagaa ttccgagatgg gaggagcttg caaagtctgg gaagatggatc tccgacttgg 2200
tttggagatc aatagatgaa agtggactga altggagagc cagcttctcc tccatcaaga 2250
aagacgggag ccttgaagc gaagagatc caacaaatc caaatltctc ccaagacaaa 2340
cagcaaaaac acctccagag calttgaaag agtccagaga ttcccggaaa cctcgtgtga 2400
gaaagggagc caagagcttg tbtgtcgctg cgttctcttg cgttccagagc gtagtctaccg 2450
actcttggag gaggagcttg gaggatggat gtagtccggc tccgttggcg taagagacaa 2500
gtccacacc gttctcagtg ctgaatgggt gttgaacttg agggagagatc gttcaatttga 2550
ctggatggga aacttccctg attttgtac ttggacaaa gctgtgtgaa acctaaagcc 2640
aagcaagctc ttggagcttg gagcttcccg aagtcggggc tgaagacctc gttttaaaca 2700
altgagatgc ggtcttggct ggaagatgg altggcggaa aggggagcagc gttcttggaa 2760
ggaagagag gagcttcttc tccgtcagagc ttcccctgtc agcccacac ccaagcccaa 2846
catgttccca tctctcttgc ttgtgac

<210> 57
<211> 906
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 7481414CB1

<400> 57
atgaagcttc acccttgcac ccaagcgccc atagacctca ctgctggcgcc agcagggggc 60
ggagcttgtg ttgtgacttg gaaaccttc gaaaccttc aagtaaagat gcaagatctc 120
cctcagcttg aaaaagagct tggcagatgc cttcttgaaa caatagaaa agtggagcac 180
cgttgccttc aagggagaa cagctctgca ctgtatagc altgacacaa gggatctctc 240
cgttcaact gtttctggct ttgcacacag ttgtcagaga aatgttgtag agtggagagc 300
aatgtgagc ttgcggagc ttgagcttgg agtcttggc actgtctgtc tggatlttgc 360
ggctgtgttc ttgcggcaa tgaatcttgg aagtgtcgcc ttgcagacac gttatggatg 420
aagatgttag gaaagatagc aaaaactat acaacaaatc gttctatggc taagatctac 480
ttcatgaag aggttctctc aggtcttat cgtgtactc cgtgcactc ttgtcagaga 540
aactctgctc alttcttca ctctgggggc ctctggggc taagaaaca gttcatatc ttgtacaa 600
ggggagatga agatltgaac aggtcctgtc ccttggatg taagtggagc ctltgtctgg 660
atcgttctc gttcttcaac altccagctg gactgtatg taagtgagag ccaagttctc 720
tctaatgttg gaaagcttgg agatltaac gaaaccttga taagtgttct gaaatgtga 780
ggagatagc cctgttatc ttgtttgaaa gcaaccttga ttccagagc ccttctcaat 840
gctgtctctc ttgtgttga ttgattacagc agaaagatga ttatgaatc gttggagaa 900
tactga

<210> 58
<211> 1840
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 7481461CB1

<400> 58
ggggcgagct ggggctggc ggccttggcg ctgcagagcc gcccggcgcc caggggcttc 60
ggcgcttcg ccttgagctc gttagcccg caggagagcc ggaagctctcc cggagagctc 120
cgtgttaact ggttcttccc ggttcttccc gggcttggag gggctccttc ggtcagcttg 180
gtctccacag gaggagcgcc actcggcgcg gggagagag ttggagcagc cgttccggcc 240
cggccacagc ggggagagct cggcgccggc cccggagacc tggagagagc cggagcgcc 300
ttctcagagc gttgagatgg agctggggcg gctggcgagc gggagagcc altagagccc 360
gaaaccccc gttggagagct ggggcttggc gttgattctg gctggcagatc gttgcaagc 420
gtctgtgttc ggtatccaga agccttggcg gttgtcttcc tgttcaatctc ctctccacat 480
cggctccaaa gacatgaca agatgttctc taagagagca ttggatgttgc cgtctccacat 540
gggagatgtc ttcttggc gccaatagc aagagcttcc aagagctatc ttgttggcg 600
gaaacagact gttctgggtg cgtctgttgg altgttggcg ctatgttca gttcttctgt 660

aagttccatc gagctctgt acctacta tggatcata ttgctgag gctgctctt 720
 tgcataccag ccttacttgg tcaatttgg acattattc aagaagcgc ttgacttgt 780
 gaattgcatt gcaactctgt gcagcaggt cttacaact ctgtctctt tgccttaag 840
 gttctcttt cggctgggt ttaacttga cacttttga gtctcttga tctctatgt 900
 tgtctcttt cggctgggt ttaacttga cacttttga gtctcttga tctctatgt 960
 tttaactttt gcaacttga aggtgagc ttaacttga cacttttga gtctcttga 1020
 tgcataccat gactcttga aggtgagc ttaacttga cacttttga gtctcttga 1080
 tgcataccat gactcttga aggtgagc ttaacttga cacttttga gtctcttga 1140
 tgcataccat gactcttga aggtgagc ttaacttga cacttttga gtctcttga 1200
 agcactcttc ttctcttga tggagatta tgccttgg ttgaagaag ttactata 1260
 tgcactcttc ttctcttga tggagatta tgccttgg ttgaagaag ttactata 1320
 tggggcctc atgctctgt gctcactat ggtctcttc gatggatgt tcttccat 1380
 tttgctctc atgctctgt gctcactat ggtctcttc gatggatgt tcttccat 1440
 tttgctctc atgctctgt gctcactat ggtctcttc gatggatgt tcttccat 1500
 tttgctctc atgctctgt gctcactat ggtctcttc gatggatgt tcttccat 1560
 tttgctctc atgctctgt gctcactat ggtctcttc gatggatgt tcttccat 1620
 tttgctctc atgctctgt gctcactat ggtctcttc gatggatgt tcttccat 1680
 tttgctctc atgctctgt gctcactat ggtctcttc gatggatgt tcttccat 1740
 tttgctctc atgctctgt gctcactat ggtctcttc gatggatgt tcttccat 1800
 tttgctctc atgctctgt gctcactat ggtctcttc gatggatgt tcttccat 1860
 tttgctctc atgctctgt gctcactat ggtctcttc gatggatgt tcttccat 1920

<210> 59
 <211> 5348
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> Incyte ID No: 7472541CB1

<400> 59

gtgacttga ggccttgcct tgcataact agataaact aganaaactg gaaattcat 60
 agnaaactg actatttga ccaagctgt ggaagaact acgcactt tcaagctgt 120
 tttcttca ctggagctgt tcaataggc agctcagct tcttcttga 180
 agnaaactg aacagctgt tgcatttga tcaataggc gaaactgt gaaactgt 240
 gaaatttga atgctctga atgctctga tcaataggc gaaactgt gaaactgt 300
 tttgctctc ggaagcga atgctctga tcaataggc ttaagctgt tcaagctgt 360
 gaaatttga atgctctga atgctctga tcaataggc ttaagctgt tcaagctgt 420
 tttgctctc ggaagcga atgctctga tcaataggc ttaagctgt tcaagctgt 480
 agnaaactg aacagctgt tgcatttga tcaataggc gaaactgt gaaactgt 540
 atacttca atgctctga atgctctga tcaataggc gaaactgt gaaactgt 600
 agnaaactg actatttga ccaagctgt ggaagaact acgcactt tcaagctgt 660
 gaaatttga atgctctga atgctctga tcaataggc ttaagctgt tcaagctgt 720
 tttgctctc ggaagcga atgctctga tcaataggc ttaagctgt tcaagctgt 780
 agnaaactg aacagctgt tgcatttga tcaataggc gaaactgt gaaactgt 840
 atacttca atgctctga atgctctga tcaataggc gaaactgt gaaactgt 900
 agnaaactg actatttga ccaagctgt ggaagaact acgcactt tcaagctgt 960
 tttgctctc ggaagcga atgctctga tcaataggc ttaagctgt tcaagctgt 1020
 agnaaactg aacagctgt tgcatttga tcaataggc gaaactgt gaaactgt 1080
 atacttca atgctctga atgctctga tcaataggc gaaactgt gaaactgt 1140
 agnaaactg actatttga ccaagctgt ggaagaact acgcactt tcaagctgt 1200
 tttgctctc ggaagcga atgctctga tcaataggc ttaagctgt tcaagctgt 1260
 agnaaactg aacagctgt tgcatttga tcaataggc gaaactgt gaaactgt 1320
 atacttca atgctctga atgctctga tcaataggc gaaactgt gaaactgt 1380
 agnaaactg actatttga ccaagctgt ggaagaact acgcactt tcaagctgt 1440
 tttgctctc ggaagcga atgctctga tcaataggc ttaagctgt tcaagctgt 1500
 agnaaactg aacagctgt tgcatttga tcaataggc gaaactgt gaaactgt 1560
 atacttca atgctctga atgctctga tcaataggc gaaactgt gaaactgt 1620
 agnaaactg actatttga ccaagctgt ggaagaact acgcactt tcaagctgt 1680
 tttgctctc ggaagcga atgctctga tcaataggc ttaagctgt tcaagctgt 1740
 agnaaactg aacagctgt tgcatttga tcaataggc gaaactgt gaaactgt 1800
 atacttca atgctctga atgctctga tcaataggc gaaactgt gaaactgt 1860
 agnaaactg actatttga ccaagctgt ggaagaact acgcactt tcaagctgt 1920

cccatcagg gccactacc gcaagggtct atggggcacc gtgaaagctc acagctctct 1980
 gtggcttca gcagctcat apaaaagat gtaactccag ataaaacct actgaccag 2040
 gtctgagc ctgctctgt ttgagacc ttgagacc ttgagacc caagctctc 2100
 tctcaca ctgctctcat tgcctgtc ttcctgtc taactctg caactctc 2160
 atggttca caaccaccg gccagcag aggtggatg atcaaaagt agtggaaat 2220
 gacatgtc taacttga attcaggcc ttgagaaa tcttgctt cacttatgt 2280
 aagaatct taacttga attcaggcc ttgagaaa tcttgctt cacttatgt 2340
 atctcatt cttatttgg ccaagctc accataac cctcaagc ggtctggg 2400
 agtctctt ggaagattca gcagcttc cagaagtga agctatag cctcagcag 2460
 tcaatcat ccaactgacc cctgacac gactcggg agacttag ggcacagtg 2520
 gcaacagc actcagatga gagatgat gactctgt gactcggg agacttag ggcacagtg 2580
 gactcgtt ttgagagc gactcgtg gactcggg acactcgg gactcgtt ttgagagc 2640
 ggtactct ttgagagc attgagcgc gactcgtg gactcggg acactcgg gactcgtt 2700
 ttgagagc ttgagagc gactcgtg gactcgtg gactcggg acactcgg gactcgtt 2760
 ttgagagc ttgagagc gactcgtg gactcgtg gactcggg acactcgg gactcgtt 2820
 ttgagagc gactcgtt gactcgtt gactcgtt gactcgtt gactcggg acactcgg 2880
 gttgagc accactgac ttgagagc ttgagagc ttgagagc ttgagagc ttgagagc 2940
 atctgagc ttgagagc ttgagagc ttgagagc ttgagagc ttgagagc ttgagagc 3000
 agaaaact ggcagcag ccaaaagc ctgactgt atgcaagc ggcagcag 3060
 acatagc ttgcaagc ggttgaag gaaagact tccggagc ggcagcag 3120
 ggcgtagc cctgagc cctgagc gactcgtt gactcgtt gactcgtt gactcgtt 3180
 catctgag atcaactac ctacttga gactcgtt gactcgtt gactcgtt gactcgtt 3240
 ggttccag atcagatgt cactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 3300
 ggttccag atcagatgt cactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 3360
 gactcgtt atactac atcagatgt cactcgtt gactcgtt gactcgtt gactcgtt 3420
 gactcgtt atcagatgt cactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 3480
 ttgagc cttcagc accactac cactcgtt gactcgtt gactcgtt gactcgtt 3540
 ttgagc atgagagc atgagc atgagc atgagc atgagc atgagc atgagc 3600
 ctgagatg cagatgt ttgagc atgagc atgagc atgagc atgagc atgagc 3660
 agtatag tcaagctt ggcagagc ttgagc atgagc atgagc atgagc atgagc 3720
 ggcagagc atgagc gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 3780
 ggcagagc atgagc gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 3840
 aggttctg ttgagc gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 3900
 tcaactac agagatgt cactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 3960
 ttgagc cactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4020
 ttgagc cactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4080
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4140
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4200
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4260
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4320
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4380
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4440
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4500
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4560
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4620
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4680
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4740
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4800
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4860
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4920
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 4980
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5040
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5100
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5160
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5220
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5280
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5340
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5400
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5460
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5520
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5580
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5640
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5700
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5760
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5820
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5880
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 5940
 gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt gactcgtt 6000

<210> 60
 <211> 5149
 <212> DNA
 <213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No: 6999183CB1

<400> 60

atgagagagag gacgacatgag cgtctgtcag caaacatggg cctctctcty caagaactgt 60
ctcaaaaat ggaagatgga aagacagacc tigtgtgaat ggtctcttcc attcttcttg 120
gtacgtcttc tgcacctatc ttctcccatc ttacatcagc ttcaatgaac tatccaatg 180
ttctcaatgg atctggagcg tgcagatagt tttaatgata ctcaatgata tcttccaatg 240
gacctctgat ccaaaatcgc ccaagagata atgaaaacag tggcttcagc ccaatctcta 300
atgcgttgga gaacatctat ggggtgtgct gatgaaaata gcatgagaaa atttgatagt 360
aaatcttcaa tagacagcat gagaagctc ttactgtata ccttctctca caattttgag 420
tttctctggg gaactggat ccccatgag agagacacat gaagacatct agctcaactg 480
caagcagggg atgaaaatat gaagtgtgga gttctcaggt tctgggagaa aggtcttcta 540
gcttctcaag ctgcactaa tgcgtctac atgaaaatcg caacaaatca tccagtgtg 600
gaacagctga tgcagtttc tgggttgaat atgaaatgat taacttttgg tgcacaaaga 660
gggtgtgcaa ctgatcttt catlctctc tgcattgat ctctctctac atttataaac 720
tatgtctcag tcaatgtatc acaaagaaaaga caatacatca cgtcaatgat gaaatagtg 780
ggactccagag agtcaatgct ctggtcttcc tgggtgttga tgcattgtcg ctctacact 840
atcatgggca cttaaatggc tcttatgtga aaatctgcac aaatctgctg cctgcagctg 900
tttgtatgg tcttccacct ctctctctc taatgcctg ctgttgatct tctgacttcc 960
ctgatggag tctgtataaa gaacctcttc cttaagcggt tggttgtgtc tcttacttcc 1020
tgacttttgg ggaactctgg atcccaagca ttctatacac atcttctcgc atttttga 1080
tgcactttgt gctctctag ccccttgcgc ttactgttg ggaatggcca gcttataat 1140
tggagctatg atctgaaatc taatggccac ttgacttct caaaaatcgc ataccctca 1200
atngctacac ttctcagtt ggttttggg acccttctgt atttggatg gaactatgat 1260
tttgaaaana ttctgcgcgc tgaatatgga cactgaltgt cctccctgt ttctctgaa 1320
tctgttcttt ggtttcaaca cggaaaggtc atcatgttgg tcccttgagaa tgaacagat 1380
tctgtaccta cactaatga ctgttttga caagctgtcc cagaattggt agtgaagaa 1440
gcatatgaa tcaaaatct taanaagaa tatcgagga agtctgtag agtagaagg 1500
ttgaaggttg tggttttga cataatgaa ggcagatca ctgcctctc tggtcacagt 1560
ggagctggaa aaactacct gttaaacata ctatgggtgt tgcctctcc aaactcagt 1620
tcaatgactg tctatatca caacttca agaatgtctc atatatgaaa tatcagcaag 1680
tcaacggatc ttgtcccaa accaaatg ctcaatgtat ttctacact gaagaagaa 1740
tcaaggtgt ttgttaaat aaagggat ttgcacatg aagtgtgaga agagtatg 1800
ctattggag aacggatcg tgaatggat ccccttcaaa ggcacggat atgaaatc 1860
ctgaagagag gaaatcaga cagatgat ctcttcaaga ccaagtttat aaatagagct 1920
gaactcttg cgaacagga ggtgttcata tcaatggga agcttgaatg tgcagacct 1980
tctctgtcc ctagaagaa atggtggcaga ggtcaacct taagtctga tctgaatga 2040
aggtgtgac cagaaglat aactacatg acttgaatg acatctctga tgcgaatg 2100
acagacaaa gttgaagaaa acttgatlat attttgctc tggaaagagc aaacaatt 2160
ccgaactct aagggatct tgaatagat tctaaacag gcatlgagga ttatgttgt 2220
tcaaaaana ctltgaatga ggtgtctcg aaatgaag gaataatcac tatgtatga 2280
tcaatgatg gaattctgg aaactatcaa actgtatggg caaaagaaat cagtggctg 2340
gttggactg acaaatgt gttctcttc cagaaacaaa ggaaaacat cagtggctg 2400
ggcgtctgga ggcagcaggt ctgtgcaat gcaaaagtc gttcctaaa gttaaagaa 2460
gaagaaaana ggcgtgggac tatatgatg ctlttgatg ttagtctat cctccaact 2520
ttgaaacac tatctcaga gtcatatcg aaagtatcc cgtggaaact gttcccaat 2580
acataatcct tctcaacag acacaacca caggtatccc tgaacctat actgttcatc 2640
aataagacag gttcaacat tgaatactt ttcatatc ttgaaggaga gaactatgt 2700
atgaaggtg atgaccttgg aactagaatg ggcacatgg acccatctta caatgtgtc 2760
atcaatgtg caggtatga aaagatcac agatttcaa tagcatgaa tacaagag 2820
ctgaattgtc ttctgtctc cctgtatgac attagaatg gactactg atttttat 2880
tctgtcagag acatctagac ttgacagag acatlttttg aagagatct gactcatag 2940
tatgttgaag gaattgaac ctctctctg atccatgg cagctctat cactcatcac 3000
atgcaatga gcaagatga ttgactaana aaaaagctc atccagctc acggaattca 3060
ggcctctaac ctctcgata ctgactaana aaaaagctc atccagctc cctgaactt 3120
ttgatctccc tgcataatga aatnatggat tatcttcta gcccaagaga gatataact 3180
ataatctaaa accgtttat tcaaatctc ttgatgtat gctatgtc atctctgt 3240
ttcttgaact atgtatctc atcaatctc cgaatggga gaaanaatg tgcattctg 3300
tcaattctc tctaatgt gttcatctc tgcataatg ctactgat ctactatct aatgaatc 3360
ggaattctg ggtcaattc tggcacaatg ttatctccc ccttcaact gatggacct 3420
ctaatctatc ttcttgagat ttctctctg tcaatgatc acttaagagc ttcaagatc 3480

84/85

gaattgtat accgtgacat gctaatcact taacttcaat ttctcaattc tctttcaat 3540
ctgcagatcc tagaaatgaa ctgcaggaag aaactaata gaaagatcc tggttcaga 3600
attctcaca gaagaacag tatlttcca aaccagaag agctctgaag agagagaga 3660
gatattcaga ttgaagaagt gaaacagtg gaactatg ctgcagaga ctltgatag 3720
acaccgttca tcatgtcag cgtctacag aagaatagc caggcaaaa gaataatgc 3780
tttctcaaaa gaaagaaac aattgccaa agaatgtct ctlttgtt taanaaggt 3840
gaattatag gactlttag acacatgga gctgttgaat gtacaactc taagatgaa 3900
acgtgagaca caaacacac tgaagacag gtgatltga aaggtgaagc tggaggtgaa 3960
ccccgggtc tcttgggtta ctgcctcag gagaatggc ttgtggccca cctgaacag 4020
agagacacac tggaggtgta cgttgcgtg aaggttcta gaaagggaga cgcataatc 4080
ggaacagag ggttatgta tgcctcaag ctgcagac agctgaaggg tccgttgag 4140
acctgttag agggataaa ggaagaatg tgcatttgc ttgaatcact ggggaacgg 4200
tcaatgttgc tcttgatga gccgtgacc ggaatggac ccgaagggca gcaagaatg 4260
tggcaggtga ttgcggccac ctltgaacg acgagaggg ggcctctct gaccaccca 4320
taactggagc agctggagcg ggtgtggac cgaatggca tcaatgtgc agaaatgag 4380
agatgatg gttcaatca acacttga agcaattg tcaaaagacta cctcttgag 4440
atgaatctga agaacctgc aaaaatgag cccctcagc cagaatcct gaggcttcc 4500
ccccaggtc ctcaagaga aagttctcc tcccttgatg tctataagt gctgtttgag 4560
gatgtggac ctltatcaga ggttctctc aaatagaaga tagttaaana gattttgag 4620
ctgagggagc aaagctctac agagtctac ctgagcag ttctctgga gcttccag 4680
gagcagagc tgggtgact tgaagagac ttgatcct cgttgaaggt gaaatccct 4740
ctgcagagc agcttgaag ctccaatcc ctatctt cttaatcct gtgacctt 4800
taagatgat atttataag cttaatatg cttaatatg agtctgaca atgtatg 4860
aaactatgc caatlatat cctcagtagt attcctaga ttgagacaa ggcagatga 4920
gtgagggcga tcaatagaga taagcttaag ccaatccat cagcttct ggcagacac 4980
aatccaatgt ttccatcgt ttatagtta aaatgatc tatatgaa ttgtcagc 5040
tctggagag gtcaatgga atgtcaatt tcttctctc gaacttcaa atattaaata 5100
tttcttatt ttlttgatla aagtataat taagacacc tatggactt 5149

85/85

THIS PAGE BLANK (USPTO)

This Page is inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURED OR ILLEGIBLE TEXT OR DRAWING
- ☒ SKEWED/SLANTED IMAGES
- ☐ COLORED OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REPERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images problems checked, please do not report the problems to the IFW Image Problem Mailbox

THIS PAGE BLANK (USPTO)